## World Academy of Science, Engineering and Technology International Journal of Mathematical and Computational Sciences Vol:14, No:12, 2020

## Effect of 16 Weeks Walking with Different Dosages on Psychosocial Function Related Quality of Life among 60 to 75 Years Old Men

Authors: Mohammad Ehsani, Elham Karimi, Hashem Koozechian

Abstract: Aim: The purpose of current semi-experimental study was a survey on effect of 16 week walking on psychosocial function related quality of life among 60 to 75 years old men. Methodology: For this reason, short from of health - related quality of life questionnaire (SF - 36) and Geriatric Depression Scale (GDS) had been distributed to the subjects at 2 times of pre - test and posttest. Statistical sample of current study was 60 to 75 years old men who placed at Kahrizak house and assessed by considering physically and medical background. Also factors of entrance to the intervention like age range, have satisfaction and have intent to participating in walking program, lack of having diabetic, cardiovascular, Parkinsonism diseases and postural, neurological, musculoskeletal disorders, lack of having clinical background like visual disorders or disordering on equilibrium system, lack of motor limitation, foot print disorders, having surgery and mental health had been determined and assessed. Finally after primary studies, 80 persons selected and categorized accidentally to the 3 experimental group (1, 2, 3 sessions per week, 30 min walking with moderate intension at every sessions) and one control group (without physical activity in period of 16 weeks). Data analysed by employing ANOVA, Pearson coefficient and Scheffe Post - Hoc tests at the significance level of p < 0.05. Results: Results showed that psychosocial function of men with 60 to 75 years old increase by influence of 16 week walking and increase of exercise sessions lead to more effectiveness of walking. Also there was no significant difference between psychosocial function of subjects within 1 session and 3 sessions experimental groups (p > 0.05). Conclusion: On the basis of results, we can say that doing regular walking with efficient and standard dosage for elderly people, can increase their quality of life. Furthermore, designing and action operation regular walking program for elderly men on the basis of special, logical and systematic pattern under the supervision of aware coaches have been recommended on the basis of results.

**Keywords:** walking, quality of life, psychosocial function, elders

Conference Title: ICSRD 2020: International Conference on Scientific Research and Development

Conference Location : Chicago, United States Conference Dates : December 12-13, 2020