

Effect of Lignocellulose-Degrading Bacteria Isolated from Termite Gut on the Nutritive Value of Wheat Straw as Ruminant Feed

Authors : Ayoub Azizi-Shotorkhoft, Tahereh Mohammadabadi, Hosein Motamedi, Morteza Chaji, Hasan Fazaeli

Abstract : This study was conducted to investigate nutritive value of wheat straw processed with termite gut symbiotic bacteria with lignocellulosic-degrading potential including *Bacillus licheniformis*, *Ochrobactrum intermedium* and *Microbacterium paludicola* in vitro. These bacteria were isolated by culturing termite guts contents in different culture media containing different lignin and lignocellulosic materials that had been prepared from water-extracted sawdust and wheat straw. Results showed that incubating wheat straw with all of three isolated bacteria increased ($P<0.05$) acid-precipitable polymeric lignin (APPL) compared to control, and highest amount of APPL observed following treatment with *B. licheniformis*. Highest and lowest ($P<0.05$) in vitro gas production and ruminal organic matter digestibility were obtained when treating wheat straw with *B. licheniformis* and control, respectively. However, other fermentation parameters such as b (i.e., gas production from the insoluble fermentable fractions at 144h), c (i.e., rate of gas production during incubation), ruminal dry matter digestibility, metabolizable energy, partitioning factor, pH and ammonia nitrogen concentration were similar between experimental treatments ($P>0.05$). It is concluded that processing wheat straw with isolated bacteria improved its nutritive value as ruminants feed.

Keywords : termite gut bacteria, wheat straw, nutritive value, ruminant

Conference Title : ICANMB 2016 : International Conference on Animal Nutrition, Mating and Breeding

Conference Location : Istanbul, Türkiye

Conference Dates : July 21-22, 2016