

Screening of Risk Phenotypes among Metabolic Syndrome Subjects in Adult Pakistani Population

Authors : Muhammad Fiaz, Muhammad Saqlain, Abid Mahmood, S. M. Saqlan Naqvi, Rizwan Aziz Qazi, Ghazala Kaukab Raja

Abstract : Background: Metabolic Syndrome is a clustering of multiple risk factors including central obesity, hypertension, dyslipidemia and hyperglycemia. These risk phenotypes of metabolic syndrome (MetS) prevalent world-wide, Therefore we aimed to identify the frequency of risk phenotypes among metabolic syndrome subjects in local adult Pakistani population. Methods: Screening of subjects visiting out-patient department of medicine, Shaheed Zulfiqar Ali Bhutto Medical University, Islamabad was performed to assess the occurrence of risk phenotypes among MetS subjects in Pakistani population. The Metabolic Syndrome was defined based on International Diabetes Federation (IDF) criteria. Anthropometric and biochemical assay results were recorded. Data was analyzed using SPSS software (16.0). Results: Our results showed that dyslipidemia (31.50%) and hyperglycemia (30.50%) was most population specific risk phenotypes of MetS. The results showed the order of association of metabolic risk phenotypes to MetS as follows hyperglycemia>dyslipidemia>obesity >hypertension. Conclusion: The hyperglycemia and dyslipidemia were found be the major risk phenotypes among the MetS subjects and have greater chances of deceloping MetS among Pakistani Population.

Keywords : dyslipidemia, hypertention, metabolic syndrome, obesity

Conference Title : ICMSED T 2018 : International Conference on Metabolic Syndrome, Endocrine Disorders and Treatments

Conference Location : Bangkok, Thailand

Conference Dates : February 08-09, 2018