

The Game of Synchronized Quadromineering

Alessandro Cincotti

Abstract—In synchronized games players make their moves simultaneously rather than alternately. Synchronized Quadromineering is the synchronized version of Quadromineering, a variants of a classical two-player combinatorial game called Domineering. Experimental results for small $m \times n$ boards (with $m + n < 15$) and some theoretical results for general $k \times n$ boards (with $k = 4, 5, 6$) are presented. Moreover, some Synchronized Quadromineering variants are also investigated.

Keywords—Combinatorial games, Synchronized games, Quadromineering.

I. INTRODUCTION

THE game of Domineering is a typical two-player game with perfect information, proposed around 1973 by Göran Andersson [2], [10], [11]. The two players, usually denoted by Vertical and Horizontal, take turns in placing dominoes (2×1 tile) on a checkerboard. Vertical is only allowed to place its dominoes vertically and Horizontal is only allowed to place its dominoes horizontally on the board. Dominoes are not allowed to overlap and the first player that cannot find a place for one of its triominoes loses. After a time the remaining space may separate into several disconnected regions, and each player must choose into which region to place a domino. Berlekamp [1] solved the general problem for $2 \times n$ board for odd n . The 8×8 board and many other small boards were recently solved by Breuker, Uiterwijk and van den Herik [4] using a computer search with a good system of transposition tables. Subsequently, Lachmann, Moore, and Rapaport solved the problem for boards of width 2, 3, 5, and 7 and other specific cases [12]. Finally, Bullock solved the 10×10 board [5].

The game of Triomineering was proposed in 2004 by Blanco and Fraenkel [3]. In Triomineering Vertical and Horizontal alternate in tiling with a straight triomino (3×1 tile) on a checkerboard. Blanco and Fraenkel calculated Triomineering and values for boards up to 6 squares and small rectangular boards.

The game of Quadromineering is a further extension of Domineering where Vertical and Horizontal alternate in tiling with a straight quadromino (4×1 tile) on a checkerboard.

II. SYNCHRONIZED GAMES

For the sake of self containment, we recall the previous results concerning synchronized games. Initially, the concept of synchronism was introduced in the games of Cutcake [6], Maundy Cake [7], Domineering [8], and Triomineering [9] in order to study combinatorial games where players make their moves simultaneously.

The author is with the School of Information Science, Japan Advanced Institute of Science and Technology, 1-1 Asahidai, Nomi, Ishikawa 923-1292 Japan, (phone: +81-761-51-1293; email: cincotti@jaist.ac.jp).

TABLE I
THE POSSIBLE OUTCOMES IN SYNCHRONIZED QUADROMINEERING.

| | Horizontal ls | Horizontal ds | Horizontal ws |
|---------------|-----------------|-----------------|-----------------|
| Vertical ls | $G = VHD$ | $G = HD$ | $G = H$ |
| Vertical ds | $G = VD$ | $G = D$ | - |
| Vertical ws | $G = V$ | - | - |

As a result, in the synchronized versions of these games there exist no zero-games (fuzzy-games), i.e., games where the winner depends exclusively on the player that makes the second (first) move. Moreover, there exists the possibility of a draw, which is impossible in a typical combinatorial game. In this work, we continue to investigate synchronized combinatorial games by focusing our attention on Quadromineering.

In the game of Synchronized Quadromineering, a general instance and the legal moves for Vertical and Horizontal are defined exactly in the same way as defined for the game of Quadromineering. There is only one difference: Vertical and Horizontal make their legal moves simultaneously, therefore, quadrominoes are allowed to overlap if they have a 1×1 tile in common. We note that 1×1 overlap is only possible within a simultaneous move. At the end, if both players cannot make a move, then the game ends in a draw, else if only one player can still make a move, then he/she is the winner.

For each player there exist 3 possible outcomes:


- 1) The player has a *winning strategy* (ws) independently of the opponent's strategy, or
- 2) The player has a *drawing strategy* (ds), i.e., he/she can always get a draw in the worst case, or
- 3) The player has a *losing strategy* (ls), i.e., he/she does not have a strategy for winning or for drawing.

Table I shows all the possible cases. It is clear that if one player has a winning strategy, then the other player has neither a winning strategy nor a drawing strategy. Therefore, the cases $ws-ws$, $ws-ds$, and $ds-ws$ never happen. As a consequence, if G is an instance of Synchronized Quadromineering, then we have 6 possible legal cases:

- 1) $G = D$ if both players have a drawing strategy, and the game will always end in a draw under perfect play, or
- 2) $G = V$ if Vertical has a winning strategy, or
- 3) $G = H$ if Horizontal has a winning strategy, or
- 4) $G = VD$ if Vertical can always get a draw in the worst case, but he/she could be able to win if Horizontal makes a wrong move, or
- 5) $G = HD$ if Horizontal can always get a draw in the worst case, but he/she could be able to win if Vertical makes a wrong move, or
- 6) $G = VHD$ if both players have a losing strategy and the outcome is totally unpredictable.


III. EXAMPLES OF SYNCHRONIZED QUADROMINEERING

The game




always ends in a draw, therefore $G = D$.

In the game



Vertical has a winning strategy moving in the central column, therefore $G = V$.


In the game



if Vertical moves in the first column we have two possibilities




therefore, either Vertical wins or the game ends in a draw. Symmetrically, if Vertical moves in the third column we have two possibilities



therefore, either Vertical wins or the game ends in a draw. It follows $G = VD$.

In the game



each player has 4 possible moves. For every move of Vertical, Horizontal can win or draw (and sometimes lose); likewise, for every move by Horizontal, Vertical can win or draw (and sometimes lose). As a result it follows that $G = VHD$.

IV. RESULTS FOR SYNCHRONIZED QUADROMINEERING

Table II shows the results obtained using an exhaustive search algorithm for $n \times m$ boards with $n + m \leq 14$.

Theorem 1: Let $G = [n, 5]$ be a rectangle of Synchronized Quadromineering with $n \geq 20$. Then Vertical has a winning strategy.

TABLE II
OUTCOMES FOR RECTANGLES IN SYNCHRONIZED QUADROMINEERING.

| | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|----|---|---|---|---|---|---|----|
| 4 | D | V | V | V | D | V | V |
| 5 | H | D | V | V | H | H | |
| 6 | H | H | D | V | H | | |
| 7 | H | H | H | D | | | |
| 8 | D | V | V | | | | |
| 9 | H | V | | | | | |
| 10 | H | | | | | | |






Fig. 1. Synchronized Quadromineering played on $n \times 5$ rectangular board.

Proof: In the beginning, Vertical will always move into the central column of the board, i.e., $(k, c), (k + 1, c), (k + 2, c), (k + 3, c)$ where $k \equiv 1 \pmod{4}$, as shown in Fig. 1. When Vertical cannot move anymore into the central column, let us imagine that we divide the main rectangle into 4×5 sub-rectangles starting from the top of the board (by using horizontal cuts). Of course, if $n \not\equiv 0 \pmod{4}$, then the last sub-rectangle will be of size $1 \times 5, 2 \times 5$ or 3×5 , and Horizontal will be able to make respectively one more move, two more moves or three more moves. We can classify all these sub-rectangles into 6 different classes:

- Class A. Vertical is able to make four more moves in each sub-rectangle of this class.



- Class B. Vertical is able to make one more move in each sub-rectangle of this class. For example



- Class C. Horizontal is able to make three more moves in each sub-rectangle and Vertical is able to make at least $\lceil |C|/2 \rceil$ moves where $|C|$ is the number of sub-rectangles

TABLE III
THE 6 CLASSES FOR 4×6 SUB-RECTANGLES.

| Class | vq | hq | vm | hm | Example |
|-------|----|----|--------|--------|---------|
| A | 1 | 0 | $5 A $ | 0 | |
| B | 1 | 1 | $2 B $ | 0 | |
| C | 0 | 1 | $ C $ | $3 C $ | |
| D | 0 | 2 | 0 | $2 D $ | |
| E | 0 | 3 | 0 | $ E $ | |
| F | 0 | 4 | 0 | 0 | |

$moves(V) \leq moves(H)$ using the data in Table III

$$5|A| + 2|B| + |C| \leq 3|C| + 2|D| + |E| + 3$$

and applying Equation 1

$$3|A| + 2|B| + 2|D| + 5|E| + 8|F| \leq 3$$

which is false because

$$|A| + |B| + |C| + |D| + |E| + |F| = \lfloor n/4 \rfloor$$

and by hypothesis $n \geq 12$. We note that if $|A| = 0$ then $|C| = 0$, $|D| = 0$, $|E| = 0$, $|F| = 0$, and $|B| \geq 3$. Moreover, if $|A| = 1$ then $|C| = 1$, $|D| = 0$, $|E| = 0$, $|F| = 0$, and $|B| \geq 1$. So $moves(V) \leq moves(H)$ does not hold and consequently $moves(V) > moves(H)$. ■

By symmetry the following two theorems hold.

Theorem 3: Let $G = [5, n]$ be a rectangle of Synchronized Quadromineering with $n \geq 20$. Then Horizontal has a winning strategy.

Theorem 4: Let $G = [6, n]$ be a rectangle of Synchronized Quadromineering with $n \geq 12$. Then Horizontal has a winning strategy.





Fig. 3. Synchronized Quadromineering played on $n \times 4$ rectangular board.


Theorem 5: Let $G = [n, 4]$ be a rectangle of Synchronized Quadromineering. If $n \equiv 0 \pmod{4}$, then Vertical has a drawing strategy.

Proof: In the beginning, Vertical will always move into the first column of the board, i.e., (k, b) , $(k+1, b)$, $(k+2, b)$, and $(k+3, b)$ where $k \equiv 1 \pmod{4}$, as shown in Fig. 3. When Vertical cannot move anymore into the central column, let us imagine that we divide the main rectangle into 4×4 sub-rectangles starting from the top of the board (by using horizontal cuts). We can classify all these sub-rectangles into 6 different classes:


- Class A. Vertical is able to make three more moves in each sub-rectangle of this class.




- Class B. Neither Vertical nor Horizontal are able to make another move in the sub-rectangles of this class. For example




- Class C. Horizontal is able to make three more moves in each sub-rectangle of this class. For example




- Class D. Horizontal is able to make two more moves in each sub-rectangle of this class. For example



- Class E. Horizontal is able to make one more move in each sub-rectangle of this class. For example



- Class F . In each sub-rectangle of this class Horizontal has already made four moves.



We show that when Vertical cannot move anymore into the first column, he/she can make a number of moves greater or equal to Horizontal, i.e., $moves(H) \leq moves(V)$. We denote with $|A|$ the number of sub-rectangles in the class A , with $|B|$ the number of sub-rectangles in the class B , and so on. We observe that $|A| = |C| + 2|D| + 3|E| + 4|F|$ because both Vertical and Horizontal have placed the same number of triominoes.

$$\begin{aligned} moves(H) &= 3|C| + 2|D| + |E| \\ &= 3|A| - 4|D| - 8|E| - 12|F| \\ &\leq 3|A| \\ &= moves(V) \end{aligned}$$

■

By symmetry the following theorem holds.


Theorem 6: Let $G = [4, n]$ be a rectangle of Synchronized Quadromineering. If $n \equiv 0 \pmod{4}$, then Horizontal has a drawing strategy.

V. SYNCHRONIZED QUADROMINEERING VARIANTS

In this section, we present results obtained using an exhaustive search algorithm for two Synchronized Quadromineering variants where quadrominoes have different shape from 1×4 and 4×1 tile.

A. First variant

In this variant, Vertical can place



and Horizontal can place





Table IV shows the results for $n \times m$ boards with $n + m \leq 12$.

B. Second variant

In this variant, Vertical can place



and Horizontal can place

TABLE IV
OUTCOMES FOR RECTANGLES IN SYNCHRONIZED QUADROMINEERING (FIRST VARIANT).

| | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
|---|----|-----|-----|-----|----|----|---|
| 3 | D | D | D | VD | D | D | D |
| 4 | D | VHD | D | HD | HD | HD | |
| 5 | D | D | VHD | HD | HD | | |
| 6 | HD | VD | VD | VHD | | | |
| 7 | D | VD | VD | | | | |
| 8 | D | VD | | | | | |
| 9 | D | | | | | | |

TABLE V
OUTCOMES FOR RECTANGLES IN SYNCHRONIZED QUADROMINEERING (SECOND VARIANT).

| | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
|---|---|-----|----|----|----|----|---|
| 3 | D | D | D | D | D | D | D |
| 4 | D | VHD | D | HD | HD | HD | |
| 5 | D | D | D | HD | HD | | |
| 6 | D | VD | VD | D | | | |
| 7 | D | VD | VD | | | | |
| 8 | D | VD | | | | | |
| 9 | D | | | | | | |




Table V shows the results for $n \times m$ boards with $n + m \leq 12$.

REFERENCES

- [1] E. R. Berlekamp, "Blockbusting and Domineering," *Journal of Combinatorial Theory Ser. A*, vol. 49, pp. 67-116, 1988.
- [2] E. R. Berlekamp, J. H. Conway, R. K. Guy, *Winning way for your mathematical plays*, A K Peters, 2001.
- [3] S. A. Blanco, A. S. Fraenkel, "Triomineering, Tridomineering, and L-Tridomineering," *Technical Report MCS04-10, Department of Computer Science and Applied Mathematics, The Weizmann Institute of Science*. Available: <http://wisdomarchive.wisdom.weizmann.ac.il:81/archive/00000367/>.
- [4] D. M. Breuker, J. W. H. M. Uiterwijk, H. J. van den Herik, "Solving 8×8 Domineering," *Theoretical Computer Science*, vol. 230, pp. 195-206, 2000.
- [5] N. Bullock, "Domineering: Solving Large Combinatorial Search Space," *ICGA Journal*, vol. 25, no. 2, pp. 67-84, 2002.
- [6] A. Cincotti, H. Iida, "The Game of Synchronized Cutcake," in *Proc. of the IEEE Symposium on Computational Intelligence and Games*, Honolulu, 2007, pp. 374-379.
- [7] A. Cincotti, H. Iida, "The Game of Synchronized Maundy Cake," in *Proc. of the 7th Annual Hawaii International Conference on Statistics, Mathematics and Related Fields*, Honolulu, 2008, pp. 422-429.
- [8] A. Cincotti, H. Iida, "The Game of Synchronized Domineering," in *Proc. of the Conference on Computers and Games 2008*, Beijing, 2008, pp. 241-251.
- [9] A. Cincotti, S. Komori, H. Iida, "The game of Synchronized Triomineering and Synchronized Tridomineering," *International Journal of Computational and Mathematical Sciences*, vol. 2, pp. 143-148, 2008.
- [10] J. H. Conway, *On Numbers and Games*, A K Peters, 2001.
- [11] M. Gardner, "Mathematical games," *Scientific American* vol. 230, pp. 106-108, 1974.
- [12] M. Lachmann, C. Moore, I. Rapaport, "Who Wins Domineering on Rectangular Boards," in R. J. Nowakowski (ed.) *More Games of No Chance*, vol. 42, pp. 307-315, Cambridge University Press 2002.