

# Periodic Solutions for a Food Chain System with Monod–Haldane Functional Response on Time Scales

Kejun Zhuang, Hailong Zhu

**Abstract**—In this paper, the three species food chain model on time scales is established. The Monod–Haldane functional response and time delay are considered. With the help of coincidence degree theory, existence of periodic solutions is investigated, which unifies the continuous and discrete analogies.

**Keywords**—Food chain system, periodic solution, time scales, coincidence degree.

## I. INTRODUCTION

**R**ESearch on food chain system has been a hotspot in population dynamics recently. Dynamical behavior of these models governed by differential equations and difference equations has been extensively studied in [1–5].

Motivated by the so-called Monod–Haldane function proposed in [6], we can get the following non-autonomous food chain model with time delays,

$$\begin{cases} \dot{u}_1(t) = u_1(t)[r_1(t) - d_1(t)u_1(t) - \frac{m_{12}(t)u_2(t)}{a_1(t)+b_1(t)u_1(t)+u_1^2(t)}], \\ \dot{u}_2(t) = u_2(t)[-r_2(t) + \frac{m_{21}u_1(t-\tau)}{a_1(t)+b_1(t)u_1(t-\tau)+u_1^2(t-\tau)} \\ - d_2(t)u_2(t) - \frac{m_{23}(t)u_3(t)}{a_2(t)+b_2(t)u_2(t)+u_2^2(t)}], \\ \dot{u}_3(t) = u_3(t)[-r_3(t) + \frac{m_{32}(t)u_2(t-\sigma)}{a_2(t)+b_2(t)u_2(t-\sigma)+u_2^2(t-\sigma)} \\ - d_3(t)u_3(t)], \end{cases} \quad (1)$$

where  $u_1(t)$ ,  $u_2(t)$  and  $u_3(t)$  stand for the population density of prey, predator and top-predator at time  $t$ , respectively. All coefficients are positive continuous functions.  $m_{i,i+1}(t)$  is the capture rate of the predator,  $m_{i+1,i}(t)$  is a measure of the food quality that the prey provided for conversion into predator birth, where  $i = 1, 2$ .

However, if the populations have non-overlapping generations, the discrete model governed difference equations is more appropriate :

$$\begin{cases} u_1(n+1) = u_1(n) \exp[r_1(n) - d_1(n)u_1(n) \\ - \frac{m_{12}(n)u_2(n)}{a_1(n)+b_1(n)u_1(n)+u_1^2(n)}], \\ u_2(n+1) = u_2(n) \exp[\frac{m_{21}u_1(n-\tau)}{a_1(n)+b_1(n)u_1(n-\tau)+u_1^2(n-\tau)} \\ - r_2(n) - d_2(n)u_2(n) - \frac{m_{23}(n)u_3(n)}{a_2(n)+b_2(n)u_2(n)+u_2^2(n)}], \\ u_3(n+1) = u_3(n) \exp[\frac{m_{32}(n)u_2(n-\sigma)}{a_2(n)+b_2(n)u_2(n-\sigma)+u_2^2(n-\sigma)} \\ - r_3(n) - d_3(n)u_3(n)], \end{cases} \quad (2)$$

Kejun Zhuang is with the Institute of Applied Mathematics, School of Statistics and Applied Mathematics, Anhui University of Finance and Economics, Bengbu 233030, P.R.China, e-mail: zhkj123@163.com

Hailong Zhu is with the Institute of Applied Mathematics, School of Statistics and Applied Mathematics, Anhui University of Finance and Economics, Bengbu 233030, P.R.China, e-mail: mmyddk@163.com

where all the coefficients are positive periodic sequences.

Enlightened by the idea of Stefan Hilger [7], to unify the continuous and discrete dynamic systems, we consider the following dynamic system on time scales

$$\begin{cases} x^\Delta(t) = r_1(t) - d_1(t)e^{x(t)} - \frac{m_{12}(t)e^{y(t)}}{a_1(t)+b_1(t)e^{x(t)}+e^{2x(t)}}, \\ y^\Delta(t) = -r_2(t) + \frac{m_{21}(t)e^{x(t-\tau)}}{a_1(t)+b_1(t)e^{x(t-\tau)}+e^{2x(t-\tau)}} - d_2(t)e^{y(t)} \\ - \frac{m_{23}(t)e^{z(t)}}{a_2(t)+b_2(t)e^{y(t)}+e^{2y(t)}}, \\ z^\Delta(t) = -r_3(t) + \frac{m_{32}(t)e^{y(t-\sigma)}}{a_2(t)+b_2(t)e^{y(t-\sigma)}+e^{2y(t-\sigma)}} - d_3(t)e^{z(t)}, \end{cases} \quad (3)$$

where all coefficients are positive  $\omega$ -periodic functions. Set  $u_1(t) = e^{x(t)}$ ,  $u_2(t) = e^{y(t)}$ ,  $u_3(t) = e^{z(t)}$ , then (3) can be reduced to (1) and (2) when  $\mathbb{T} = \mathbb{R}$  and  $\mathbb{T} = \mathbb{Z}$ , respectively.

The aim of this paper is to study the periodicity of three-species food chain system on time scales. We would like to mention that there are several papers on periodicity in dynamic systems on time scales by using the coincidence degree theory, see [8–11]. The remainder of the paper is organized as follows. In the following section, some preliminary results about calculus on time scales and the continuation theorem are stated. Next, the sufficient conditions for the existence of periodic solutions are explored.

## II. PRELIMINARIES

For convenience, we first present the useful lemma about time scales and the continuation theorem of the coincidence degree theory; more details can be found in [12–14].

A time scale  $\mathbb{T}$  is an arbitrary nonempty closed subset of real numbers  $\mathbb{R}$ . Throughout this paper, we assume that the time scale  $\mathbb{T}$  is unbounded above and below, such as  $\mathbb{R}$ ,  $\mathbb{Z}$  and  $\bigcup_{k \in \mathbb{Z}} [2k, 2k+1]$ . The following definitions and lemmas about time scales are from [12].

**Lemma 2.1.** ([13]) Let  $t_1, t_2 \in I_\omega$  and  $t \in \mathbb{T}$ . If  $g : \mathbb{T} \rightarrow \mathbb{R} \in C_{rd}(\mathbb{T})$  is  $\omega$ -periodic, then

$$g(t) \leq g(t_1) + \frac{1}{2} \int_k^{k+\omega} |g^\Delta(s)| \Delta s$$

and

$$g(t) \geq g(t_2) - \frac{1}{2} \int_k^{k+\omega} |g^\Delta(s)| \Delta s,$$

the constant factor  $\frac{1}{2}$  is the best possible.

For simplicity, we use the following notations throughout this paper. Let  $\mathbb{T}$  be  $\omega$ -periodic, that is  $t \in \mathbb{T}$  implies  $t + \omega \in \mathbb{T}$ ,  
 $k = \min\{\mathbb{R}^+ \cap \mathbb{T}\}$ ,  $I_\omega = [k, k + \omega] \cap \mathbb{T}$ ,  $g^L = \inf_{t \in \mathbb{T}} g(t)$ ,

$$g^M = \sup_{t \in \mathbb{T}} g(t), \quad \bar{g} = \frac{1}{\omega} \int_{I_\omega} g(s) \Delta s = \frac{1}{\omega} \int_k^{k+\omega} g(s) \Delta s,$$

where  $g \in C_{rd}(\mathbb{T})$  is an  $\omega$ -periodic real function, i.e.,  $g(t + \omega) = g(t)$  for all  $t \in \mathbb{T}$ .

Next, we state the Mawhin's continuation theorem, which is a main tool in the proof of our theorem.

**Lemma 2.2.** [14] (Continuation Theorem) Let  $L$  be a Fredholm mapping of index zero and  $N$  be  $L$ -compact on  $\bar{\Omega}$ . Suppose

- (a) for each  $\lambda \in (0, 1)$ , every solution  $u$  of  $Lu = \lambda Nu$  is such that  $u \notin \partial\Omega$ ;
- (b)  $QN u \neq 0$  for each  $u \in \partial\Omega \cap \ker L$  and the Brouwer degree  $\deg\{JQN, \Omega \cap \ker L, 0\} \neq 0$ .

Then the operator equation  $Lu = Nu$  has at least one solution lying in  $\text{Dom } L \cap \bar{\Omega}$ .

### III. MAIN RESULTS

**Theorem 3.1.** If

$$m_{32}^L e^{L_2} > r_3^M (a_2^M + b_2^M e^{M_2} + e^{2M_2})$$

holds, where

$$L_2 = \ln \frac{a_2^L r_3^L}{m_{32}^M} - \frac{m_{21}^M \omega}{b_1^L}$$

and

$$M_2 = \ln \frac{r_1^M (a_1^M + \frac{b_1^M r_1^M}{d_1^M} + \frac{2r_1^M}{d_1^M})}{m_{12}^L} + \frac{m_{21}^M \omega}{b_1^L},$$

then system (3) has at least one  $\omega$ -periodic solution.

**Proof** Let  $X = Z = \{(x, y, z)^T \in C(\mathbb{T}, \mathbb{R}^3) : x(t + \omega) = x(t), y(t + \omega) = y(t), z(t + \omega) = z(t), \forall t \in \mathbb{T}\}$ ,  $\|(x, y, z)^T\| = \max_{t \in I_\omega} |x(t)| + \max_{t \in I_\omega} |y(t)| + \max_{t \in I_\omega} |z(t)|$ ,  $(x, y, z)^T \in X$  (or in  $Z$ ). Then  $X$  and  $Z$  are both Banach spaces when they are endowed with the above norm  $\|\cdot\|$ . Let

$$N \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} N_1 \\ N_2 \\ N_3 \end{bmatrix},$$

where  $N_1 = r_1(t) - d_1(t)e^{x(t)} - \frac{m_{12}(t)e^{y(t)}}{a_1(t)+b_1(t)e^{x(t)}+e^{2x(t)}}$ ,  
 $N_2 = -r_2(t) + \frac{m_{21}(t)e^{x(t-\tau)}}{a_1(t)+b_1(t)e^{x(t-\tau)}+e^{2x(t-\tau)}} - d_2(t)e^{y(t)} - \frac{m_{23}(t)e^{z(t)}}{a_2(t)+b_2(t)e^{y(t)}+e^{2y(t)}}$ ,  $N_3 = -r_3(t) + \frac{m_{32}(t)e^{y(t-\sigma)}}{a_2(t)+b_2(t)e^{y(t-\sigma)}+e^{2y(t-\sigma)}} - d_3(t)e^{z(t)}$ .

$$L \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x^\Delta \\ y^\Delta \\ z^\Delta \end{bmatrix}, \quad P \begin{bmatrix} x \\ y \\ z \end{bmatrix} = Q \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \frac{1}{\omega} \int_k^{k+\omega} x(t) \Delta t \\ \frac{1}{\omega} \int_k^{k+\omega} y(t) \Delta t \\ \frac{1}{\omega} \int_k^{k+\omega} z(t) \Delta t \end{bmatrix}.$$

Obviously,  $\ker L = \mathbb{R}^3$ ,  $\text{Im } L = \{(x, y, z)^T \in Z : \bar{x} = \bar{y} = \bar{z} = 0, t \in \mathbb{T}\}$ ,  $\dim \ker L = 3 = \text{codim Im } L$ . Since  $\text{Im } L$  is closed in  $Z$ , then  $L$  is a Fredholm mapping of index zero. It is

easy to show that  $P$  and  $Q$  are continuous projections such that  $\text{Im } P = \ker L$  and  $\text{Im } L = \ker Q = \text{Im}(I - Q)$ . Furthermore, the generalized inverse (of  $L$ )  $K_P : \text{Im } L \rightarrow \ker P \cap \text{Dom } L$  exists and is given by

$$K_P \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} \int_k^t x(s) \Delta s - \frac{1}{\omega} \int_k^{k+\omega} \int_k^t x(s) \Delta s \Delta t \\ \int_k^t y(s) \Delta s - \frac{1}{\omega} \int_k^{k+\omega} \int_k^t y(s) \Delta s \Delta t \\ \int_k^t z(s) \Delta s - \frac{1}{\omega} \int_k^{k+\omega} \int_k^t z(s) \Delta s \Delta t \end{bmatrix}.$$

Thus

$$QN \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{\omega} \int_k^{k+\omega} (r_1(t) - d_1(t)e^{x(t)} - \frac{m_{12}(t)e^{y(t)}}{a_1(t)+b_1(t)e^{x(t)}+e^{2x(t)}}) \Delta t \\ \frac{1}{\omega} \int_k^{k+\omega} (-r_2(t) + \frac{m_{21}(t)e^{x(t-\tau)}}{a_1(t)+b_1(t)e^{x(t-\tau)}+e^{2x(t-\tau)}} - d_2(t)e^{y(t)} - \frac{m_{23}(t)e^{z(t)}}{a_2(t)+b_2(t)e^{y(t)}+e^{2y(t)}}) \Delta t \\ \frac{1}{\omega} \int_k^{k+\omega} (-r_3(t) + \frac{m_{32}(t)e^{y(t-\sigma)}}{a_2(t)+b_2(t)e^{y(t-\sigma)}+e^{2y(t-\sigma)}} - d_3(t)e^{z(t)}) \Delta t \end{bmatrix},$$

and

$$K_P(I - Q)N \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$= \begin{bmatrix} \int_k^t x(s) \Delta s - \frac{1}{\omega} \int_k^{k+\omega} \int_k^t x(s) \Delta s \Delta t - (t - k - \frac{1}{\omega} \int_k^{k+\omega} (t - k) \Delta t) \bar{x} \\ \int_k^t y(s) \Delta s - \frac{1}{\omega} \int_k^{k+\omega} \int_k^t y(s) \Delta s \Delta t - (t - k - \frac{1}{\omega} \int_k^{k+\omega} (t - k) \Delta t) \bar{y} \\ \int_k^t z(s) \Delta s - \frac{1}{\omega} \int_k^{k+\omega} \int_k^t z(s) \Delta s \Delta t - (t - k - \frac{1}{\omega} \int_k^{k+\omega} (t - k) \Delta t) \bar{z} \end{bmatrix}.$$

Clearly,  $QN$  and  $K_P(I - Q)N$  are continuous. According to the Arzela-Ascoli theorem, it is not difficult to show that  $K_P(I - Q)N(\bar{\Omega})$  is compact for any open bounded set  $\Omega \subset X$  and  $QN(\bar{\Omega})$  is bounded. Thus,  $N$  is  $L$ -compact on  $\bar{\Omega}$ .

Now, we shall search an appropriate open bounded subset  $\Omega$  for the application of the continuation theorem, Lemma 2.2. For the operator equation  $Lu = \lambda Nu$ , where  $\lambda \in (0, 1)$ , we have

$$\begin{cases} u_1^\Delta(t) = \lambda(r_1(t) - d_1(t)e^{x(t)} - \frac{m_{12}(t)e^{y(t)}}{a_1(t)+b_1(t)e^{x(t)}+e^{2x(t)}}), \\ u_2^\Delta(t) = \lambda(-r_2(t) + \frac{m_{21}(t)e^{x(t-\tau)}}{a_1(t)+b_1(t)e^{x(t-\tau)}+e^{2x(t-\tau)}} - d_2(t)e^{y(t)} - \frac{m_{23}(t)e^{z(t)}}{a_2(t)+b_2(t)e^{y(t)}+e^{2y(t)}}), \\ u_3^\Delta(t) = \lambda(-r_3(t) + \frac{m_{32}(t)e^{y(t-\sigma)}}{a_2(t)+b_2(t)e^{y(t-\sigma)}+e^{2y(t-\sigma)}} - d_3(t)e^{z(t)}). \end{cases} \quad (4)$$

Assume that  $(u_1, u_2, u_3)^T \in X$  is a solution of system (4) for a certain  $\lambda \in (0, 1)$ . Integrating (4) on both sides from  $k$  to  $k + \omega$ , we obtain

$$\begin{cases} \int_k^{k+\omega} [d_1(t)e^{x(t)} + \frac{m_{12}(t)e^{y(t)}}{a_1(t)+b_1(t)e^{x(t)}+e^{2x(t)}}] \Delta t = \bar{r}_1 \omega, \\ \int_k^{k+\omega} [r_2(t) + d_2(t)e^{y(t)} + \frac{m_{23}(t)e^{z(t)}}{a_2(t)+b_2(t)e^{y(t)}+e^{2y(t)}}] \Delta t \\ = \int_k^{k+\omega} \frac{m_{21}(t)e^{x(t-\tau)}}{a_1(t)+b_1(t)e^{x(t-\tau)}+e^{2x(t-\tau)}} \Delta t, \\ \int_k^{k+\omega} [r_3(t) + d_3(t)e^{z(t)}] \Delta t \\ = \int_k^{k+\omega} \frac{m_{32}(t)e^{y(t-\sigma)}}{a_2(t)+b_2(t)e^{y(t-\sigma)}+e^{2y(t-\sigma)}} \Delta t. \end{cases} \quad (5)$$

Since  $(x, y, z)^T \in X$ , there exist  $\xi_i, \eta_i \in I_\omega$ ,  $i = 1, 2, 3$ , such that

$$\begin{aligned} x(\xi_1) &= \min_{t \in I_\omega} \{x(t)\}, & x(\eta_1) &= \max_{t \in I_\omega} \{x(t)\}, \\ y(\xi_2) &= \min_{t \in I_\omega} \{y(t)\}, & y(\eta_2) &= \max_{t \in I_\omega} \{y(t)\}, \\ z(\xi_3) &= \min_{t \in I_\omega} \{z(t)\}, & z(\eta_3) &= \max_{t \in I_\omega} \{z(t)\}. \end{aligned} \quad (6)$$

From (4) and (5), we have

$$\begin{aligned} \int_k^{k+\omega} |x^\Delta(t)| \Delta t &\leq 2\bar{r}_1\omega, \\ \int_k^{k+\omega} |y^\Delta(t)| \Delta t &\leq 2\frac{m_{21}^M\omega}{b_1^L}, \\ \int_k^{k+\omega} |z^\Delta(t)| \Delta t &\leq 2\frac{m_{32}^M\omega}{b_2^L}. \end{aligned}$$

By the first equation of (5) and (6),

$$d_1(\xi_1)e^{x(\xi_1)} < r_1(\xi_1),$$

that is

$$x(\xi_1) < \ln \frac{r_1^M}{d_1^L}.$$

From the second equation of (5), we have

$$r_2(\eta_2) < \frac{m_{21}(\eta_2)e^{x(\eta_2-\tau)}}{a_1(\eta_2)}$$

and

$$x(\eta_1) \geq x(\eta_2 - \tau) > \ln \frac{r_2^L a_1^L}{m_{21}^M}.$$

According to Lemma 2.1, we have

$$\begin{aligned} x(t) &\leq x(\xi_1) + \frac{1}{2} \int_k^{k+\omega} |x^\Delta(t)| \Delta t \\ &\leq \ln \frac{r_1^M}{d_1^L} + \bar{r}_1\omega \triangleq M_1, \end{aligned}$$

$$\begin{aligned} x(t) &\geq x(\eta_1) - \frac{1}{2} \int_k^{k+\omega} |x^\Delta(t)| \Delta t \\ &\geq \ln \frac{r_2^L a_1^L}{m_{21}^M} - \bar{r}_1\omega \triangleq L_1. \end{aligned}$$

From the first equation of (5) and (6), we can obtain

$$\frac{m_{12}(\xi_1)e^{y(\xi_1)}}{a(\xi_1) + b_1(\xi_1)e^{x(\xi_1)} + e^{2x(\xi_1)}} < r_1(\xi_1),$$

and

$$y(\xi_2) < y(\xi_1) < \ln \frac{r_1^M(a_1^M + \frac{b_1^M r_1^M}{d_1^L} + \frac{2r_1^M}{d_1^L})}{m_{12}^L},$$

then

$$\begin{aligned} y(t) &\leq y(\xi_2) + \frac{1}{2} \int_k^{k+\omega} |y^\Delta(t)| \Delta t \\ &< \ln \frac{r_1^M(a_1^M + \frac{b_1^M r_1^M}{d_1^L} + \frac{2r_1^M}{d_1^L})}{m_{12}^L} + \frac{m_{21}^M\omega}{b_1^L} \\ &\triangleq M_2. \end{aligned}$$

From the third equation of (5), we have

$$\frac{m_{32}(\xi_3)}{a_2(\xi_3)} e^{y(\xi_3-\sigma)} > r_3(t),$$

this reduces to

$$y(\eta_2) \geq y(\xi_3 - \sigma) > \ln \frac{a_2^L r_3^L}{m_{32}^M},$$

then

$$\begin{aligned} y(t) &\geq y(\eta_2) - \frac{1}{2} \int_k^{k+\omega} |y^\Delta(t)| \Delta t \\ &\geq \ln \frac{a_2^L r_3^L}{m_{32}^M} - \frac{m_{21}^M\omega}{b_1^L} \triangleq L_2. \end{aligned}$$

According to the first equation of (5), we have

$$\frac{m_{32}(\xi_3)}{b_2(\xi_3)} > d_3(\xi_3)e^{z(\xi_3)},$$

then

$$\begin{aligned} z(t) &\leq z(\xi_3) + \frac{1}{2} \int_k^{k+\omega} |z^\Delta(t)| \Delta t \\ &\leq \ln \frac{m_{32}^M}{b_2^L d_3^L} + \frac{m_{32}^M\omega}{b_2^L} \triangleq M_3. \end{aligned}$$

Besides, we have

$$d_3^M e^{z(\eta_3)} > \frac{m_{32}^L e^{L_2}}{a_2^M + b_2^M e^{M_2} + e^{2M_2}} - r_3^M,$$

then

$$\begin{aligned} z(t) &\geq z(\eta_3) - \frac{1}{2} \int_k^{k+\omega} |z^\Delta(t)| \Delta t \\ &\geq \ln \frac{\frac{m_{32}^L e^{L_2}}{a_2^M + b_2^M e^{M_2} + e^{2M_2}} - r_3^M}{d_3^M} - \frac{m_{32}^M\omega}{b_2^L} \\ &\triangleq L_3. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \max_{t \in [k, k+\omega]} |x(t)| &\leq \max\{|M_1|, |L_1|\} \triangleq R_1, \\ \max_{t \in [k, k+\omega]} |y(t)| &\leq \max\{|M_2|, |L_2|\} \triangleq R_2, \\ \max_{t \in [k, k+\omega]} |z(t)| &\leq \max\{|M_3|, |L_3|\} \triangleq R_3. \end{aligned}$$

Clearly,  $R_1, R_2$  and  $R_3$  are independent of  $\lambda$ . Let  $R = R_1 + R_2 + R_3 + R_0$ , where  $R_0$  is taken sufficiently large such that for for the following algebraic equations:

$$\begin{cases} \bar{r}_1 - \bar{d}_1 e^x - \frac{1}{\omega} \int_\kappa^{\kappa+\omega} \frac{m_{12}(t)e^y}{a_1(t)+b_1(t)e^x+e^{2x}} \Delta t = 0, \\ -\bar{r}_2 + \frac{1}{\omega} \int_\kappa^{\kappa+\omega} \frac{m_{21}(t)e^x}{a_1(t)+b_1(t)e^x+e^{2x}} \Delta t - \bar{d}_2 e^y = 0, \\ -\frac{1}{\omega} \int_\kappa^{\kappa+\omega} \frac{\bar{m}_{23}e^z}{a_2(t)+b_2(t)e^y+e^{2y}} \Delta t = 0, \\ -\bar{r}_3 + \frac{1}{\omega} \int_\kappa^{\kappa+\omega} \frac{m_{32}(t)e^y}{a_2(t)+b_2(t)e^y+e^{2y}} \Delta t - \bar{d}_3 e^z = 0, \end{cases} \quad (7)$$

every solution  $(x^*, y^*, z^*)^T$  of (7) satisfies  $\|(x^*, y^*, z^*)^T\| < R$ . Now, we define  $\Omega = \{(x, y, z)^T \in X : \|(x, y, z)^T\| < R\}$ . Then it is clear that  $\Omega$  verifies the requirement (a) of Lemma 2.2. If  $(u_1, u_2, u_3)^T \in \partial\Omega \cap \ker L = \partial\Omega \cap \mathbb{R}^3$ , then  $(x, y, z)^T$  is

a constant vector in  $\mathbb{R}^3$  with  $\|(x, y, z)^T\| = |x| + |y| + |z| = R$ , so we have

$$QN \begin{bmatrix} x \\ y \\ z \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

By the assumption in Theorem 3.1 and the definition of topological degree, a direct calculation yields  $\deg(JQN, \Omega \cap \ker L, 0) \neq 0$ . We have verified that  $\Omega$  satisfies all requirements of Lemma 2.2; therefore, system (3) has at least one  $\omega$ -periodic solution in  $\text{Dom } L \cap \Omega$ . This completes the proof.

#### ACKNOWLEDGMENT

This work was supported by the Anhui Provincial Natural Science Foundation under Grant 1208085QA11, Natural Science Foundation of the Higher Education Institutions of Anhui Province under Grant KJ2012B002 and KJ2012B004.

#### REFERENCES

- [1] Jean-Pierre Francoise, Jaume Llibre, Analytical study of a triple Hopf bifurcation in a tritrophic food chain model, *Applied Mathematics and Computation*, 217(2011), 7146–7154.
- [2] B.Mukhopadhyay, R.Bhattacharyya, A stage-structured food chain model with stage dependent predation: Existence of codimension one and codimension two bifurcations, *Nonlinear Analysis: RWA*, 12(2011), 3056–3072.
- [3] Changyong Xu, Meijuan Wang, Permanence for a delayed discrete three-level food-chain model with Beddington-DeAngelis functional response, *Applied Mathematics and Computation*, 187(2007), 1109–1119.
- [4] Rui Xu, Lansun Chen, Feilong Hao, Periodic solutions of a discrete time Lotka-Volterra type food-chain model with delays, *Applied Mathematics and Computation*, 171(2005), 91–103.
- [5] Weiming Wang, Hailing Wang, Zhenqing Li, The dynamic complexity of a three-species Beddington-type food chain with impulsive control strategy, *Chaos, Solitons and Fractals*, 32(2007), 1772–1785.
- [6] J.F. Andrews, A mathematical model for the continuous culture of microorganisms utilizing inhibitory substrates, *Biotechnol Bioengr.* 10(1986), 707–723.
- [7] Stefan Hilger, Analysis on measure chains—a unified approach to continuous and discrete calculus, *Results Math.*, 18(1990), 18–56.
- [8] Martin Bohner, Meng Fan, Jiming Zhang, Periodicity of scalar dynamic equations and applications to population models, *J. Math. Anal. Appl.* 330(2007), 1–9.
- [9] Yu Tong, Zhenjie Liu, Zhiying Gao, Yonghong Wang, Existence of periodic solutions for a predator-prey system with sparse effect and functional response on time scales, *Commun. Nonlinear Sci. Numer. Simulat.*, 17(2012), 3360–3366.
- [10] Kejun Zhuang, Periodic solutions for a three species predator-prey system on time scales, *Int. J. Dynamical Systems and Differential Equations*, 3(2011), 3-11.
- [11] Martin Bohner, Meng Fan, Jiming Zhang, Existence of periodic solutions in predator-prey and competition dynamic systems, *Nonl. Anal: RWA*, 7(2006), 1193–1204.
- [12] Martin Bohner, Allan Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Boston: Birkhäuser, 2001.
- [13] Bingbing Zhang, Meng Fan, A remark on the application of coincidence degree to periodicity of dynamic equations on time scales, *J. Northeast Normal University(Natural Science Edition)*, 39(2007), 1–3. (in Chinese)
- [14] R. E. Gaines, J. L. Mawhin, *Coincidence Degree and Nonlinear Differential Equations*, Lecture Notes in Mathematics, Berlin: Springer-Verlag, 1977.