

Molecular Identification of ESBL Genes *bla*_{GES-1}, *bla*_{VEB-1}, *bla*_{CTX-M}, *bla*_{OXA-1}, *bla*_{OXA-4}, *bla*_{OXA-10} and *bla*_{PER-1} in *Pseudomonas* *aeruginosa* Strains Isolated from Burn Patients by PCR, RFLP and Sequencing Techniques

Fereshteh Shacheraghi, Mohammad Reza Shakibaie, Hanieh Noveiri

Abstract—Fourty one strains of ESBL producing *P.aeruginosa* which were previously isolated from burn patients in Kerman University general hospital, Iran were subjected to PCR, RFLP and sequencing in order to determine the type of extended spectrum β -lactamases (ESBL), the restriction digestion pattern and possibility of mutation among detected genes. DNA extraction was carried out by phenol chloroform method. PCR for detection of *bla* genes was performed using specific primer for each gene. Restriction Fragment Length Polymorphism (RFLP) for ESBL genes was carried out using *EcoRI*, *NheI*, *PvuII*, *EcoRV*, *DdeI*, and *PstI* restriction enzymes. The PCR products were subjected to direct sequencing of both the strands for identification of the ESBL genes. The *bla*_{CTX-M}, *bla*_{VEB-1}, *bla*_{PER-1}, *bla*_{GES-1}, *bla*_{OXA-1}, *bla*_{OXA-4} and *bla*_{OXA-10} genes were detected in the (n=1) 2.43%, (n=41)100%, (n=28) 68.3%, (n=10) 24.4%, (n=29) 70.7%, (n=7)17.1% and (n=38) 92.7% of the ESBL producing isolates respectively. The RFLP analysis showed that each ESBL gene has identical pattern of digestion among the isolated strains. Sequencing of the ESBL genes confirmed the genuinity of PCR products and revealed no mutation in the restriction sites of the above genes. From results of the present investigation it can be concluded that *bla*_{VEB-1} and *bla*_{CTX-M} were the most and the least frequently isolated ESBL genes among the *P.aeruginosa* strains isolated from burn patients. The RFLP and sequencing analysis revealed that same clone of the *bla* genes were indeed existed among the antibiotic resistant strains.

Keywords—ESBL genes, PCR, RFLP, Sequencing, *P.aeruginosa*

I. INTRODUCTION

Pseudomonas aeruginosa is ranking second among gram negative hospital acquiring pathogens and one of leading cause of burn infections reported to the National Nosocomial Infection Surveillance System [1, 2]. The idea of eradication of *P. aeruginosa* from burn patients through intense antimicrobial therapy may lead to significant selection of

resistance strains in burn unit of the hospitals [3]. One of the important features of these strains is resistant to multiple clinically important antibiotics like third generation of cephalosporins, imipenem and aztronam [4]. Many *P.aeruginosa* strains produces different class of extended spectrum β -lactamases (ESBLs) that enable bacterium to stand against extended -spectrum cephalosporins, such as cefotaxime, ceftriaxone and ceftazidime and have been reported with increasing frequency [5, 6]. ESBL mediate resistance to cephalosporin antibiotics and were first discovered in Europe in the early 1980s. They have become a widespread problem, particularly in *Klebsiella pneumoniae*, and increasingly in non-typhoid *Salmonella* species. The OXA-type ESBLs have been found mainly in *P. aeruginosa* isolates from Turkey and France [7].

Traditionally, ESBL enzymes have been derivatives of TEM and SHV parent enzymes. The last year, however, has seen an explosion of developments in ESBLs of non-TEM, non-SHV lineage in Europe. The CTX-M type ESBLs have become particularly widespread [8]. Bert *et al.*, [9] detected *bla*_{PSE} and *bla*_{OXA} gene variants using PCR. The genotypes were distinguished by restriction of PCR products with endonucleases recognizing sites involved in point mutations. Jiang *et al.*, [10] studied a total of 75 clinical isolates of *P. aeruginosa*. Thirty-four of 36 multidrug-resistant *P. aeruginosa* clinical isolates were positive for ESBLs and *bla*_{VEB-3} was the most prevalent ESBL gene reported by the authors. Antibiotic susceptibility tests and PCR amplification of genes encoding class A (*bla*_{PSE-1}, *bla*_{PER-1}, *bla*_{VEB-1}, *bla*_{TEM}, *bla*_{SHV}, *bla*_{CTX-M} and *bla*_{GES-1}) and class D β -lactamases (*bla*_{OXA-groupI}, *bla*_{OXA-groupII} and *bla*_{OXA-groupIII}) in *P.aeruginosa* were carried out by Lee *et al.*, (11). In 64 (25.4%) isolates, structural genes for PSE-1 (6.3%), OXA-10 (13.1%), OXA-4 (4.3%), OXA-30 (2.0%), OXA-2 (2.3%) and OXA-17 (0.4%) were found, their distribution varied between provinces. None harboured *bla*_{PER-1}, *bla*_{VEB-1}, *bla*_{TEM}, *bla*_{SHV}, *bla*_{CTX-M} and *bla*_{GES-1}. Similarly, PCR and sequence analysis revealed the presence of the *bla*_{CTX-M-1}, *bla*_{SHV-1} and *bla*_{TEM-116} genes in the *P. aeruginosa* and *bla*_{CTX-M-1} and *bla*_{SHV-1} in the *Stenotrophomonas maltophilia* strains [12]. Mirsalehian *et al.*, [13] studied the prevalence of ESBLs and antimicrobial susceptibilities of *P. aeruginosa* isolated from burn patients in Tehran, Iran. It was found that 50 (74.62%), 33 (49.25%) and

F. Shacheraghi & H. Noveiri are with Department of Microbiology, Pasture Institute of Iran, Tehran, Iran.
MR. Shakibaie is with Department of Microbiology, Kerman University of Medical Sciences, Kerman, Iran.
Corresponding author: MR. Shakibaie Department of Microbiology, Kerman University of Medical Sciences, End of 22-Bahman BLVD, 76167-14111, Kerman, Iran. Work Phone: +98-341- 3221660-64. Mobile Phone: +98-913-140-8226, Fax: +98-341- 3221676. e-mail: mr_shakibaie@kmu.ac.ir

21 (31.34%) strains among 67 ESBL-producing strains amplified *bla*_{OXA-10}, *bla*_{PER-1} and *bla*_{VEB-1} respectively. Woodford *et al.*, [14] studied *P. aeruginosa* isolates producing VEB-type ESBL in the United Kingdom. In one UK centre, a VEB-1 producing strain was isolated. This strain was resistant to all beta-lactams, aminoglycosides and ciprofloxacin, remaining susceptible only to colistin (MICs ≤ 1 mg/L). Two other *P. aeruginosa* isolates co-producing both VEB and VIM enzymes were received from two other UK hospitals; one isolate represented inter-hospital spread of the O15 strain and the second was distinct. Existence of SHV-type ESBL genes in *P. aeruginosa* by PCR-restriction fragment length polymorphism have been reported by Blagui *et al.* [15]. Restriction of PCR products by *DdeI* and *BsrI* revealed the same restriction pattern with the *bla*_{SHV-1} positive control.

In pervious study, we isolated 41 strains of ESBL producing *P.aeruginosa* from burn infected patients in Kerman University of Medical Sciences general hospital, Iran [4]. In present study we tried to identify the ESBL genes, the restriction digestion patterns and possibility of mutations in restriction sites among detected *bla* genes by PCR, RFLP and sequencing techniques.

II. MATERIAL AND METHODS

Bacterial sources

120 strains of *P.aeruginosa* were isolated from burn infected patients in burn unit of the Kerman University general hospital, Iran within one year period. Identification of the isolates was done according to standard microbiology procedures [4]. The ESBL production was detected among 41 strains by phenotypic confirmatory test and double disc synergy methods as previously published by Shakibaie *et al.* [4]. The standard microbial cultures for detection of ESBL genes were including *P.aeruginosa* KOAS containing *bla*_{PER-1} obtained from Prof. Nordman, institutes pasture France. *K. pneumoniae* 7881, *K. pneumoniae* CHU-BICETRE containing *bla*_{CTX-M} and *bla*_{GES-1} respectively provided by Prof. Nordman, France. *P.aeruginosa* ATCC 27853, *E.coli* ATCC25922 ESBL sensitive strains and *P. aeruginosa* strains containing *bla*_{VEB-1}, *OXA-1* *OXA-4*, and *bla*_{OXA-10} were available in the stock collection of Institute pasture of Iran [16].

Antibiotic sensitivity tests

The antibiotic sensitivity of the above strains was carried out by disc diffusion break point assay and MIC was determined by agar dilution method in Muller-Hinton agar (MHA) as described previously [4].

DNA extraction

One ml of 24 hours grown *P.aeruginosa* cultures in Triplicate Soy Broth (TSB) medium (Merck, Germany) were transferred into 1.5ml sterile Eppendorff microfuge tubes and centrifuged at 10.000g for 10 minutes. The pellets were dissolved in 600µl of lysis buffer (NaCl 1M, Tris-HCl 1M, EDTA 0.5M), 20µl SDS (25%), 3µl of proteinase -K (20mg/ml) and incubated at 60°C for 1 hour. After the lysis, 620µl of phenol/chloroform/isoamylalcohol (25:24:1

Volume/Volume) were added to the above solutions, carefully vortexed, and centrifuged at 12.000g for 10 minutes. The supernatants were transferred to other sterile microfuge tubes. 1ml of 95% cold ethanol was added and allowed to stand for 1hour in refrigeration condition (4°C). DNA was then precipitated in each tube by centrifugation at 12.000g for 10 minutes. The precipitated DNA was dissolved in 50µl of 10mM Tris EDTA - buffer (TE) containing 10µl of RNase -A as described by Sambrook *et al.*, (28) and used for further investigation.

PCR reaction

The primer sequence for ESBL genes is showed in Table-1. The PER-1 sequence derived from *Pseudomonas aeruginosa* KOAS Producing PER-1 (Pasteur Institute of France), VEB-1 *Pseudomonas aeruginosa* 10.2 (24), GES-1 *K. pneumoniae*, Prf. P. Nordmann CHU Bicetre-France and CTX-M *K.pneumoniae* 7881 kindly provided by P. Nordmann.

A typical 25µl PCR reaction mixture for every primer set was consisted of 1X- PCR reaction buffer (Fermentase, Lithuania), 1.5 µm MgCl₂ (25mM), 0.7 µm of each dNTP (10mM), 0.7µl of each primer, 1unit of Taq DNA polymerase 5U/µl (Fermentase, Lithuania) and 0.5 µl of 10µg DNA template. Amplification was carried out in a thermocycler (Eppendorf Mastercycler®, Massachusetts, USA). Agarose gel electrophoresis (1.0%) of PCR products was carried out in horizontal bed apparatus using 1mM Tris-Borate- EDTA (TBE) buffer (pH-7.2) at 90V for 1hour and the DNA bands were then stained with 0.5µg/ml ethidium bromide (Sigma USA) for 10 minutes. The gels were washed twice with D/W and observed under U.V. gel documentation (UV DOC, England) at 280nm. 1000-100bp DNA ladder was used to confirm the size of each specific *bla* gene. Simultaneously, a positive control was run for each ESBL gene.

Restriction Fragment Length Polymorphism (RFLP)

The RFLPs of ESBL genes were carried out using *EcoRI*, *NheI*, *PvuII*, *EcoRV*, *DdeI*, and *PstI* restriction enzymes respectively. The enzymes were purchased from Fermentas Company Ltd. 1.5µl of each enzyme and DNA mixtures were added to 1µl restriction buffer and kept at 37°C. Digestions were completed within 3 hours and the mixture then loaded into 1.5% agarose gel concentration. The electrophoresis was conducted for each RFLP set in TBE-buffer at 60V for two hours. The gel was then stained with ethidium bromide solution (0.5µg/ml) and observed under UV light gel documentation system as described above.

Sequencing of the PCR products

DNA sequencing for all 41 strains was performed for identification of detected *bla* genes using primers as shown in Table-1. The PCR products of above genes were further purified with PCR purification Kits (Fermentas) and subjected to direct sequencing of both the strands performed by the Macrogen Company (Seoul, Korea) as described previously [12]. The nucleotide and deduced amino acid sequences were analyzed with CROMASPRO-2 and MEGA- 4 softwares.

TABLE I
THE PRIMERS AND SEQUENCES USED FOR AMPLIFICATION WITH
RESPECT TO ESBL GENES

Primer Name	5' – Sequence - 3'	Detected gene	Molecular Weight	Reference
VEB-1(F)	CGACTTCATTCCCGATG C	bla _{VEB}	643bp	21
VEB-1(R)	GGACTCTGCAACAATAAC GC			
GES-1(F)	ATGCGCTTCATTACGCAC	bla _{GES}	643bp	22
GES-1(R)	CTATTGTCCGTGCTCAGG			
CTX-M(F)	CGCTTTCGATGTGCAG	bla _{CTX-M}	550bp	23
CTX-M(R)	ACCGCGATATCGTTGGT			
OXA-1(F)	AGCCGTAAATAAAGCC C	bla _{OXA-1}	882bp	24
OXA-1(R)	CTTGATTGAAGGGTGGGC G			
OXA-4(F)	TCAACAGATATCTACTG TT	bla _{OXA-4}	216bp	25
OXA-4(R)	TTTATCCCATTTGAATATG GT			
OXA-10(F)	TCAACAAATCGCCAGAGA AG	bla _{OXA-10}	277bp	26
OXA-10(R)	TCCCACACAGAAAAACC A			
PER-1(F)	AATTTGGGCTTAGGGCAG AA	bla _{PER}	925bp	27
PER-1(R)	ATGAATGTCATTATAAAA GC			

F= Forward primer
R= Reverse primer

III. RESULTS

The results of PCR of the *bla* genes of the *P.aeruginosa* strains isolated from one of the Kerman University of Medical Sciences general hospital in Iran are shown in Figure 1. The *bla_{VEB-1}* (643bp) was the most frequent ESBL gene and isolated from almost all [100% (n=41)] of the ESBL producing strains, while, *bla_{OXA-10}* (227bp) was detected in 92.7% (n=38) of the isolates. The other ESBL genes were detected among ESBL producing populations including *bla_{CTX-M}* (550bp) *bla_{PER-1}* (927bp), *bla_{GES-1}* (864bp), *bla_{OXA-1}* (909bp), *bla_{OXA-4}* (216bp) with frequency 2.43% (n=1), 68.3% (n=28), 24.4% (n=10), 70.7% (n=29) and 17.1% (n=7) respectively as shown in Figure 1. *bla_{CTX-M}* gene was detected in only one isolate that showed MIC₂ to aztronam. Restriction digestion pattern of *bla_{VEB-1}* PCR product by *EcoRI* enzyme revealed identical digestion pattern among the strains [two bands with 152 and 491bp] while, digestion of *bla_{GES-1}* gene by *NheI* resulted in two bands [207 and 657bp]. Similarly, digestion of *bla_{OXA-1}* and *bla_{OXA-4}* with restriction enzyme *EcoRV* resulted in two bands with 124 and 785 bp and 9 & 207bp respectively. Digestion of *bla_{CTX-M}* gene with *PvuII* resulted in three bands with 170, 180 and 297bp. The results were further confirmed by sequencing technique (Fig. 2). In order to increase the accuracy of the results, the sequencing of both forward and reverse strands of the ESBL genes were carried out and revealed that the same sequencing pattern with each *bla* positive controls. Figure 2 shows the conserved region for the restriction sequence *bla_{VEB-1}* gene compared with positive control. This was obtained by comparing with blast data and the chromatograms and predicted amino acid sequences of the above genes. The sequencing of *bla* genes also confirmed the RFLP results and suggested that there were no any mutations in the restriction sites of these genes. It therefore might be

suggested that same clone of each ESBL gene has been spread among the *P.aeruginosa* isolated strains by mobile genetic elements like transposons or integrons. This is due to selective pressure (presence of antibiotics) exerted in the hospital environment on the bacteria.



Fig. 1. Agarose gel electrophoresis of PCR products of ESBL genes *bla_{CTX-M}*, *bla_{VEB-1}*, *bla_{PER-1}*, *bla_{GES-1}*, *bla_{OXA-1}*, *bla_{OXA-4}* and *bla_{OXA-10}* detected among *P.aeruginosa* isolated from burn patients. M= Molecular weight marker (100-1000bp). Positive control was also run alongside the tests as shown in the right side of each gene

5'-NAGCNANNGCNNATTGCTTTAGCCGTTTTGTCTGAGATAGATAAAGGGAATCTTTCTTTTGAACAAAAATAGAGATTACCCCTCAAGACCTTTTGCCTAAAATGTGGAGTCCGATTAAAGAGGAATTCCCTAATGGAACAACCTTTGACGATTGAACAAATACTAAATTATACAGTATCAGAGAGCGACAATATTGGTTGTGATATTTTGCTAAAATTAATCGGAGGAACCTGATTCTGTTCAAAAATCTTGAATGCTAATCATTCTACTGATATTCAATCAAAGCAAACGAAGAACAATGCACAAGGATTGGAATACCCAATATCAAATTTGGCAACCCCAACAGCGATGAACAACTGTTAATAGATACTTATAATAAAGAACCATTACTTTCTAAAAAAGTTATGATTTTATTTGGAAAAATTATGAGAGAAACAACAACAGGAAGTAACCGATTAAAAGGACAATTACCAAAGAATAACAATTGTTGCTCATAAACAGGGACTTCCGGAATAAATATGGAAATGTCAGCAGCCACTAATGATGTTGGGGTAATTACTTTACCGAATGGACAATTAATTTTATAAGCGTATTGTTGGCGAAAAGATGTCACAAAAATCCCATTTTGTAGATTTGTCCAGGATATGTGCTGTCAAGTAGAAAAATTTCTGTAATGCTCTCTATATTTCTCTGACAAAAGGACACTGTTCAACCCTTGGGACATCCATTCTCGCTATAAGNGNACNGGGGGAAN 3'-




Fig. 2. Sequences and Chromatogram of *bla_{VEB-1}* of *P.aeruginosa* strains isolated from burn patients in Kerman University general hospital. The nucleotide sequences was analyzed with CROMASPRO-2 and MEGA 4 softwares and confirmed by blast system in internet

IV. DISCUSSION

Chromosomal or plasmid mediated antibiotic resistance is common place in *P.aeruginosa* isolated from burn infected patients in different hospitals in Iran [1]. In this research, the frequency of the ESBL genes among ESBL producing *P.aeruginosa* burn isolates were studied. The *bla_{VEB}* gene was detected in almost all ESBL producing isolate. This was accordance with other results in Middle East and Iran. A retrospective survey was conducted to characterize beta-

lactamases in a collection of 43 ceftazidime-resistant *P. aeruginosa* isolates recovered from patients with bloodstream infections hospitalized at a Brazilian teaching hospital between January and December 2005 (17). It was found that nine isolates (20.9%) produced an ESBL, either GES-1 (n = 7, 16.3%) or CTX-M-2 (n = 2, 4.6%).

Drug susceptibility testing and PCR assay were used to determine the antibiotic susceptibility patterns and prevalence of genes encoding five different extended ESBLs (PER, VEB, SHV, GES, and TEM) among 600 isolates of *P. aeruginosa* cultured from patients at two hospitals in Tehran (16). The frequency of *bla*_{VEB}, *bla*_{SHV}, *bla*_{PER}, *bla*_{GES}, and *bla*_{TEM} among the ESBL isolates (MIC ≥ 16) were 24%, 22%, 17%, 0%, and 9%, respectively. Isolates containing *bla*_{VEB} were resistant to almost all tested antibiotics except imipenem. However, the frequency of the *bla*_{OXA} group of ESBL in our study was higher as compared with other authors, while only one isolate carried the gene for *bla*_{CTX-M}. 41 *P. aeruginosa* strains were isolated with ESBLs from several wards were collected over 9 months in 2003 and 2004 in a hospital in Warsaw, Poland (18). The isolates were recovered from patients with multiple types of infections, mostly respiratory tract and postoperative wound infections. All 41 isolates produced the PER-1 ESBL, originally observed in Turkey but recently also identified in several countries in Europe and the Far East. The *bla*_{PER-1} gene resided within the Tn1213 composite transposons, which was chromosomally located. The PER, VEB, GES, and IBC beta-lactamases, have been found mainly in *P. aeruginosa* and at a limited number of geographic sites (16, 19). Similarly, in our study, the ESBL gene VEB-1 exhibited similar pattern of digestion in all 41 isolates. Therefore, it might be suggested that the gene was resided on class 1 integron. Prevalence of Class-A ESBL in Clinical Isolates of *Acinetobacter baumannii* and *P. aeruginosa* were studied by Oh *et al* [20]. It was found that the most prevalent class A ESBL genotype in *Acinetobacter baumannii* isolates was *bla*_{PER-1} (n=6), and *bla*_{SHV-12} gene was also found in one *P. aeruginosa* isolate.

V. CONCLUSION

From results of present investigation, it can be concluded that *bla*_{VEB-1} and *bla*_{CTX-M} genes were the most and the least frequently isolated ESBL gene among the *P. aeruginosa* strains detected from burn infected patients while, *bla*_{OXA-10} was the second most frequently isolated gene. They exhibited similar pattern of digestion of PCR products when digested with *EcoRI*, *DdeI* and *PvuII* restriction enzymes. The sequencing analysis further confirmed the results of PCR and revealed no mutation in the restriction sites of the above genes. The results also confirmed that these ESBL genes evolved from same clone and spread through the *P. aeruginosa* population by selective pressure of antibiotics that prescribed in this region.

ACKNOWLEDGMENT

Our sincere thanks to the authority of Kerman University of Medical Sciences, Kerman, Iran for awarding grant (number

18/1387) to Dr. Shakibaie MR and Microbiology Laboratory, Pasture Institute of Iran for providing Lab. Facilities for this research.

REFERENCES

- [1] Shakibaie M.R. Plasmid mediated metal and antibiotic resistance in *Pseudomonas aeruginosa* strains isolated from burn patients. Medical Journal of the Islamic Republic of Iran 2002; 16:159-163.
- [2] Talbot GH, Bradley J, Edwards JE Jr, Gilbert D, Scheld M, Bartlett JG. Bad bugs need drugs: an update on the development pipeline from the antimicrobial availability task force of the Infectious Diseases Society of America. Clin Infect Dis 2006; 42:657-68. (Erratum in Clin Infect Dis 2006; 42:1065).
- [3] Navon-Venezia S, Ben-Ami R, Carmeli Y. Update on *Pseudomonas aeruginosa* and *Acinetobacter baumannii* infections in the healthcare setting. Curr Opin Infect Dis 2005; 18:306-13.
- [4] Shakibaie MR, Shahcheraghi F, Hashemi A, Saeed Adeli N. Detection of TEM, HSV and PER type extended spectrum beta- lactamases genes among clinical strains of *Pseudomonas aeruginosa* isolated from burnt patients at Shafa – hospital, Kerman, Iran. Iranian J Basic Medical Sciences 2008; 11:104-111.
- [5] Jones RN. Resistance patterns among nosocomial pathogens: trends over the past few years. Chest 2001; 119(2 suppl):S397-404.
- [6] Weld Hagen G, Poirel L, Nordmann P. Ambler class A extended-spectrum β -lactamase in *Pseudomonas aeruginosa*: novel development and clinical impact. Antimicrob Agents Chemother 2003; 47:2385-92.
- [7] Bradford PA. Extended-spectrum β -lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. Clin Microbiol Rev 2001; 14:933-51.
- [8] Paterson L. Extended-spectrum beta-lactamases: the European experience. Current Opinion in Infectious Diseases 2001; 14: 697-701.
- [9] Bert F, Branger C, Lambert-Zechovsky N. Identification of PSE and OXA β -lactamase genes in *Pseudomonas aeruginosa* using PCR-restriction fragment length polymorphism. Antimicrob Chemother 2002; 50: 11-18.
- [10] Jiang X, Zhang Z, Li M, Zhou D, Ruan F, Lu E. Detection of Extended-Spectrum β -Lactamases in clinical isolates of *Pseudomonas aeruginosa*. Antimicrob Agents and Chemother 2006; 50: 2990-2995.
- [11] Lee S, Park YJ, Kim M, Lee HK, Han K, Kang CS, et al. Prevalence of Ambler class A and D β -lactamases among clinical isolates of *Pseudomonas aeruginosa* in Korea. J Antimicrob Chemother 2005; 56:122-7.
- [12] Al Naiemi, N, Duim, B, Bart, A. A CTX-M extended-spectrum β -lactamase in *Pseudomonas aeruginosa* and *Stenotrophomonas maltophilia*. J Med Microbiol 2006; 55:1607-1608.
- [13] Mirsalehian A, Feizabadi M, Nakhjavani F, Jabalameli F, Goli H, Kalantari N. Detection of VEB-1, OXA-10 and PER-1 genotypes in extended-spectrum β -lactamase-producing *Pseudomonas aeruginosa* strains isolated from burn patients. Burns 2010; 36: 70- 74.
- [14] Woodford N, Zhang J, Kaufmann ME, Yarde S, Tomas Mdel M, Faris C, Vardhan MS, Dawson S, Cotterill SL, Livermore DM. Detection of *Pseudomonas aeruginosa* isolates producing VEB-type extended-spectrum beta-lactamases in the United Kingdom. Antimicrob Chemother 2008; 62:1265-8.
- [15] Blagui S, Achour W, Abdeladhim A, Ben Hassen A. Identification of SHV-type extended spectrum beta-lactamase genes in *Pseudomonas aeruginosa* by PCR-restriction fragment length polymorphism and insertion site restriction-PCR. Pathol Biol (Paris) 2009; 57(5): 420-4.
- [16] Shahcheraghi F, Nikbin VS, Feizabadi MM. Prevalence of ESBLs genes among multidrug-resistant isolates of *Pseudomonas aeruginosa* isolated from Patients in Tehran. Microb Drug Resist 2009; 15:37-9.
- [17] picão rc, poirel l, gales ac, nordmann p. diversity of beta-lactamases produced by ceftazidime-resistant *pseudomonas aeruginosa* isolates causing bloodstream infections in brazil. antimicrob agents chemother 2009; 53:3908-13.
- [18] Empel J, Filczak K, Mrówka A, Hryniewicz W, Livermore DM, Gniadkowski M. Outbreak of *Pseudomonas aeruginosa* i nfections with PER-1 extended-spectrum beta-lactamase in Warsaw, Poland: further evidence for an international clonal complex. J Clin Microbiol 2007; 45:2829-34.

- [19] Livermore DM. Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare? Clin Infect Dis 2002; 34:634-40.
- [20] Oh SJ, Lee SU, Hwang HY, Bae K, et al. Prevalence of Class A Extended-Spectrum β -Lactamases in Clinical Isolates of *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. Korean J Lab Med 2006; 26(1):14-20.
- [21] Poirel L, Rotimi VO, Mokaddas ME, Karim A, and Nordmann P. VEB-1-like extended-spectrum β -lactamases in *Pseudomonas aeruginosa*, Kuwait. Emerg Infect Dis 2001; 7:468- 470.
- [22] Poirel L, Weldhagen GF, Naas T, Champs C, et al. GES-2, a class A β -lactamase from *Pseudomonas aeruginosa* with increased hydrolysis of imipenem. Antimicrob Agents Chemother 2001; 45:2598-2603.
- [23] Ahmed AM, Nakano H, Shimamoto T. The first characterization of extended-spectrum β -lactamase producing *Salmonella* in Japan. Antimicrob Chemother 2004; 54:283-284.
- [24] Aubert D, Poirel L, Chevalier J, et al. *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2001; 45: 1615–1620.
- [25] Kenji Marumo, Takeda A, Nakamura Y and Nakay K. Detection of OXA-4 β -lactamase in *Pseudomonas aeruginosa* isolates by genetic methods. Antimicrob Chemother 1999; 43: 187-193.
- [26] Bert F, Branger C, Zechovsky NL. Identification of PSE and OXA β -lactamase genes in *Pseudomonas aeruginosa* using PCR-restriction fragment length polymorphism. Antimicrob Chemother 2002; 50:11–8.
- [27] Claeys G, Verschraegen G, Baere T, and Vaneechoutte M. PER-1 β -lactamase producing *Pseudomonas aeruginosa* in an intensive care unit. Antimicrob Chemother 2000; 45:924-925.
- [28] Sambrook J, Fritsch EF, Maniatis T. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press 1989.