

# New exact three-wave solutions for the (2+1)-dimensional asymmetric Nizhnik-Novikov-Veselov system

Fadi Awawdeh and O. Alsayyed

**Abstract**—New exact three-wave solutions including periodic two-solitary solutions and doubly periodic solitary solutions for the (2+1)-dimensional asymmetric Nizhnik–Novikov–Veselov (ANNV) system are obtained using Hirota’s bilinear form and generalized three-wave type of ansatz approach. It is shown that the generalized three-wave method, with the help of symbolic computation, provides an effective and powerful mathematical tool for solving high dimensional nonlinear evolution equations in mathematical physics.

**Keywords**—Soliton Solution; Hirota Bilinear Method; ANNV System.

## I. INTRODUCTION

IN last decades, increasing attention has been paid to the study of the solution theory in many natural sciences particularly in almost all branches of physics like the fluid dynamics, plasma physics, field theory, nonlinear optics and condensed matter physics.

Integrable models play a prominent role in theoretical physics. The reason is not only the direct phenomenological interest of some of them, but also the fact that they often provide some deep insights into the mathematical structure of theory in which they arise. Up to now, the (1+1)-dimensional integrable models are well understood due to many systematic methods such as the inverse scattering transformation [9], the Darboux transformation, Hirota’s bilinear method [3], [10], [11], [12], [13], [15], Bäcklund transformation method [18], Painlevé expansion method [8], tanh function method [16], homogenous balance method [19], variable separation approach [6] and the three-wave approach [2], [5]. However, studies of the (2+1)-dimensional cases are fewer in number and such systems are being actively investigated from different viewpoints.

In this paper, we consider the following (2+1)-dimensional asymmetric Nizhnik–Novikov–Veselov (ANNV) equation:

$$u_t + u_{xxx} + 3 \left[ u \left( \int u_x dy \right) \right]_x = 0. \quad (1)$$

Fadi Awawdeh is with the Mathematics Department, Hashemite University, Zarqa 13115, Jordan (e-mail: awawdeh@hu.edu.jo).

O. Alsayyed is with the Mathematics Department, Hashemite University, Zarqa 13115, Jordan (e-mail: o.alsayyed@yahoo.com).

Boiti et al. [4] was first derived Eq. (1) by using the weak Lax pair. It may be considered as a model for an incompressible fluid where  $u$  and  $v$  are the components of the (dimensionless) velocity [8]. The spectral transformation for this system has been investigated in [1], [4]. Moreover, Hu et al. used the Darboux transformations to find the variable separation solutions of this equation in [14], Wazwaz used the Hirota bilinear method to derive N-soliton solutions of this completely integrable equation [17], Dai and Wang used the exp-function method for a new general variable separation solutions [7] and the Bäcklund transformation for the ANNV equation was derived by using the extended homogeneous balance method by Zhang in [20].

## II. ANNV SYSTEM

The ANNV equation can be written as

$$\begin{cases} u_t + u_{xxx} + 3[uv]_x = 0 \\ u_x = v_y. \end{cases} \quad (2)$$

To solve the ANNV system (2), we substitute the dependent variable transformation

$$u = 2(\ln f)_{xy}, \quad v = 2(\ln f)_{xx}, \quad (3)$$

where  $f(x, y, t)$  is an unknown real function. Substituting (3) into system (2), we can get

$$\begin{aligned} f \left( \frac{\partial^2 f}{\partial y \partial t} + \frac{\partial^4 f}{\partial y \partial x^3} \right) - \frac{\partial f}{\partial y} \left( \frac{\partial f}{\partial t} + \frac{\partial^3 f}{\partial x^3} \right) + \\ 3 \left( \frac{\partial^2 f}{\partial y \partial x} \frac{\partial^2 f}{\partial x^2} - \frac{\partial f}{\partial x} \frac{\partial^3 f}{\partial y \partial x^2} \right) = 0 \end{aligned}$$

or equivalently in bilinear form

$$D_y (D_t + D_x^3) f \cdot f = 0, \quad (4)$$

where the Hirota  $D$ -operator is defined by [10]

$$D_x^m D_t^n f \cdot g = \left( \frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^m \left( \frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^n f(x, t) g(x', t') \Big|_{x'=x, t'=t}.$$

We propose a novel test function of extended three-soliton method

$$f(x, y, t) = e^{\xi_1} + d_1 \cos(\xi_2) + d_2 \cosh(\xi_3) + d_3 e^{-\xi_1} \quad (5)$$

where

$$\xi_i = a_i x + b_i y + c_i t, \quad i = 1, 2, 3 \quad (6)$$

and  $a_i, b_i, c_i, d_i$  are some constants to be determined later.

Substituting (5) into Eq. (4), and equating all the coefficients of  $\sin(a_2 x + b_2 y + c_2 t)$ ,  $\cos(a_2 x + b_2 y + c_2 t)$ ,  $\sinh(a_3 x + b_3 y +$

$c_3 t$ ),  $\cosh(a_3 x + b_3 y + c_3 t)$  and any product of them to zero, we get the set of algebraic equations for  $a_i, b_i, c_i, d_i$  ( $i = 1, 2, 3$ )

$$\begin{aligned} c_2 b_1 - b_1 a_3^2 + b_2 c_1 + 3b_1 a_2 a_1^2 - 3b_2 a_1 a_2^2 + b_2 a_1^3 &= 0, \\ c_1 b_1 - c_2 b_2 - 3b_2 a_2 a_1^2 + b_2 a_2^3 - 3b_1 a_1 a_2^2 + b_1 a_1^3 &= 0, \\ c_3 b_2 + c_2 b_3 - 3b_2 a_3 a_2^2 - b_3 a_2^3 + 3b_3 a_2 a_2^2 + b_2 a_3^3 &= 0, \\ c_3 b_3 - c_2 b_2 - 3b_3 a_3 a_2^2 + b_2 a_2^3 - 3b_2 a_2 a_2^2 + b_3 a_3^3 &= 0, \\ c_1 b_3 + c_3 b_1 + 3b_1 a_3 a_1^2 + b_3 a_1^3 + 3b_3 a_1 a_3^2 + b_1 a_3^3 &= 0, \\ c_1 b_1 + c_3 b_3 + 3b_3 a_3 a_1^2 + b_1 a_1^3 + 3b_1 a_1 a_3^2 + b_3 a_3^3 &= 0, \\ 4c_1 b_1 d_3 - c_2 b_2 d_1^2 + 4b_2 d_1^2 a_2^3 + c_3 b_3 d_2^2 + & \\ 16b_1 d_3 a_1^3 + 4b_3 a_3^3 d_2^2 &= 0 \end{aligned}$$

Solving the resulting equations simultaneously with the help of symbolic computation, we obtain the following:

Case 1:

$$a_1 = a_2 = c_1 = c_2 = b_3 = 0, \quad c_3 = -a_3^3, \quad (7)$$

where  $a_3, b_1, b_2, c_3, d_i$  ( $i = 1, 2, 3$ ) are some free constants.

Substituting (5) into Eq. (3) with (7) and (6), we obtain the periodic two-solitary solution as follows

$$u(x, y, t) = \frac{2a_3 d_2 \sinh(a_3 x - a_3^3 t) \times (d_1 b_2 \sin(b_2 y) - 2b_1 \sqrt{d_3} \sinh(b_1 y - \theta))}{(2\sqrt{d_3} \cosh(b_1 y - \theta) + d_2 \cosh(a_3 x - a_3^3 t) + d_1 \cos(b_2 y))^2} \quad (8)$$

$$v(x, y, t) = \frac{2a_3^2 d_2 \times (d_2 + d_1 \cos(b_2 y) \cosh(a_3 x - a_3^3 t) + 2\sqrt{d_3} \cosh(b_1 y - \theta) \cosh(a_3 x - a_3^3 t))}{(2\sqrt{d_3} \cosh(b_1 y - \theta) + d_2 \cosh(a_3 x - a_3^3 t) + d_1 \cos(b_2 y))^2} \quad (9)$$

where  $\theta = \frac{1}{2} \ln d_3$  and  $d_3 > 0$ . Moreover, when  $\theta = \frac{1}{2} \ln(-d_3)$  and  $d_3 < 0$

$$u(x, y, t) = \frac{2a_3 d_2 \sinh(a_3 x - a_3^3 t) \times (d_1 b_2 \sin(b_2 y) - 2b_1 \sqrt{-d_3} \cosh(b_1 y - \theta))}{(2\sqrt{-d_3} \sinh(b_1 y - \theta) + d_2 \cosh(a_3 x - a_3^3 t) + d_1 \cos(b_2 y))^2} \quad (10)$$

$$v(x, y, t) = \frac{2a_3^2 d_2 \times (d_2 + d_1 \cos(b_2 y) \cosh(a_3 x - a_3^3 t) + 2\sqrt{-d_3} \sinh(b_1 y - \theta) \cosh(a_3 x - a_3^3 t))}{(2\sqrt{-d_3} \sinh(b_1 y - \theta) + d_2 \cosh(a_3 x - a_3^3 t) + d_1 \cos(b_2 y))^2} \quad (11)$$

Case 2:

$$b_1 = b_2 = c_3 = a_3 = 0, \quad c_1 = -a_1^3, \quad c_2 = -a_2^3, \quad (12)$$

where  $a_1, a_2, b_3, d_i$  ( $i = 1, 2, 3$ ) are some free constants.

Substituting (5) into Eq. (3) with (12) and (6), we obtain the periodic two-solitary solution

$$u(x, y, t) = \frac{2b_3 d_2 \sinh(b_3 y) \times (a_2 d_1 \sin(a_2 x + a_2^3 t) - 2a_1 \sqrt{d_3} \sinh(a_1 x - a_1^3 t - \theta))}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} \quad (13)$$

$$v(x, y, t) = \frac{2(d_1^2 a_2^2 - 4d_3 a_1^2) + 4d_1 \sqrt{d_3} (a_2^2 - a_1^2) \cosh(a_1 x - a_1^3 t - \theta) \cos(a_2 x + a_2^3 t)}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} + \frac{2d_2 (2a_1^2 \sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) - d_1 a_2^2 \cos(a_2 x + a_2^3 t)) \cosh(b_3 y)}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} + \frac{8a_1 a_2 d_1 \sqrt{d_3} \sinh(a_1 x - a_1^3 t - \theta) \sin(a_2 x + a_2^3 t)}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} \quad (14)$$

where  $\theta = \frac{1}{2} \ln d_3$  and  $d_3 > 0$ . In the case when  $\theta = \frac{1}{2} \ln(-d_3)$  and  $d_3 < 0$ , we get that

$$u(x, y, t) = \frac{2b_3 d_2 \sinh(b_3 y) \times (a_2 d_1 \sin(a_2 x + a_2^3 t) - 2a_1 \sqrt{-d_3} \cosh(a_1 x - a_1^3 t - \theta))}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} \quad (15)$$

$$v(x, y, t) = \frac{2(d_1^2 a_2^2 - 4d_3 a_1^2) + 4d_1 \sqrt{-d_3} (a_2^2 - a_1^2) \sinh(a_1 x - a_1^3 t - \theta) \cos(a_2 x + a_2^3 t)}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} + \frac{2d_2 (2a_1^2 \sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) - d_1 a_2^2 \cos(a_2 x + a_2^3 t)) \cosh(b_3 y)}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} + \frac{8a_1 a_2 d_1 \sqrt{-d_3} \cosh(a_1 x - a_1^3 t - \theta) \sin(a_2 x + a_2^3 t)}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cosh(b_3 y))^2} \quad (16)$$

Case 3: Using the transformation  $b_3 = iB_3$ , where  $B_3$  is real, in Eqs. (13)-(14) and Eqs. (15)-(16), we get the doubly periodic solitary solutions

$$u(x, y, t) = \frac{2B_3 d_2 \sin(B_3 y) \times (2a_1 \sqrt{d_3} \sinh(a_1 x - a_1^3 t - \theta) - a_2 d_1 \sin(a_2 x + a_2^3 t))}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} \quad (17)$$

$$v(x, y, t) = \frac{2(d_1^2 a_2^2 - 4d_3 a_1^2) + 4d_1 \sqrt{d_3} (a_2^2 - a_1^2) \cosh(a_1 x - a_1^3 t - \theta) \cos(a_2 x + a_2^3 t)}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} + \frac{2d_2 (2a_1^2 \sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) - d_1 a_2^2 \cos(a_2 x + a_2^3 t)) \cos(B_3 y)}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} + \frac{8a_1 a_2 d_1 \sqrt{d_3} \sinh(a_1 x - a_1^3 t - \theta) \sin(a_2 x + a_2^3 t)}{(2\sqrt{d_3} \cosh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} \quad (18)$$

when  $d_3 > 0$  and when  $d_3 < 0$

$$u(x, y, t) = \frac{2B_3 d_2 \sin(B_3 y) \times (2a_1 \sqrt{-d_3} \cosh(a_1 x - a_1^3 t - \theta) - a_2 d_1 \sin(a_2 x + a_2^3 t))}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} \quad (19)$$

$$v(x, y, t) = \frac{2(d_1^2 a_2^2 - 4d_3 a_1^2) + 4d_1 \sqrt{-d_3} (a_2^2 - a_1^2) \sinh(a_1 x - a_1^3 t - \theta) \cos(a_2 x + a_2^3 t)}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} + \frac{2d_2 (2a_1^2 \sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) - d_1 a_2^2 \cos(a_2 x + a_2^3 t)) \cos(B_3 y)}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} + \frac{8a_1 a_2 d_1 \sqrt{-d_3} \cosh(a_1 x - a_1^3 t - \theta) \sin(a_2 x + a_2^3 t)}{(2\sqrt{-d_3} \sinh(a_1 x - a_1^3 t - \theta) + d_1 \cos(a_2 x + a_2^3 t) + d_2 \cos(B_3 y))^2} \quad (20)$$

To our knowledge, these solutions (8)–(20) have not been reported in other literatures. All the solutions of the (2+1)-dimensional ANN system obtained in this paper include three independent variables. In these solutions the arbitrary constants imply that (1) has abundant local physical structures.

### III. CONCLUSION

By choosing different ansatz of extended three-soliton type, we obtain a new type of three-wave solution, periodic type of three-wave solutions, including periodic two-solitary-wave, doubly periodic solitary-wave of (2+1)-dimensional ANN system. Actually, our present short paper investigates different mechanical features of these wave solutions. It is worth noting that this is merely a beginning work, and we can obtain richer exact solutions by a more general ansatz of extended three-soliton type.

### REFERENCES

- [1] M.J. Ablowitz and P.A. Clarkson, Solitons, Nonlinear Evolution Equations and Inverse Scattering, Lecture Notes Series, vol. 149. Cambridge University Press, Cambridge (1991).
- [2] F. Awawdeh, New exact solitary wave solutions of the Zakharov-Kuznetsov equation in the electron-positron-ion plasmas, Appl. Math. Comput. 218 (2012) 7139-7143.

- [3] F. Awawdeh, H.M. Jaradat, S. Al-Shara', Applications of a simplified bilinear method to ion-acoustic solitary waves in plasma, *Eur. Phys. J. D.*, doi:10.1140/epjd/e2011-20518-0, 2012.
- [4] M Boiti, J. Leon, M. Manna and F Pempinelli, On the spectral transform of a Korteweg-de Vries equation in two spatial dimensions, *Inverse Probl.* 2 (1986) 271.
- [5] Z. Dai, S. Lin, Ha. Fu and X. Zeng, Exact three-wave solutions for the KP equation, *Appl. Math. Comput.* 216 (2010) 1599-1604.
- [6] C. Dai, S. Wu and X. Cen, New Exact Solutions of the (2+1)-Dimensional Asymmetric Nizhnik-Novikov-Veselov System, *Int J Theor Phys* 47 (2008) 1286–1293.
- [7] C. Dai and Y. Wang, New variable separation solutions of the (2+1)-dimensional asymmetric Nizhnik-Novikov-Veselov system, *Nonlinear Analysis* 71 (2009) 1496-1503.
- [8] P.G. Estevez, S. Leble, A wave equation in (2+ 1): Painlevé analysis and solutions, *Inverse Problems* 11 (1995) 925.
- [9] C.S. Gardner, J.M. Green, M.D. Kruskal, R.M. Miura, Method for solving the Korteweg–de Vries equation, *Phys. Rev. Lett.* 19 (1967) 1095–1097.
- [10] R. Hirota, Direct methods in soliton theory, in: R.K. Bullough, P.J. Caudrey (Eds.), *Solitons*, Springer, Berlin, 1980.
- [11] R. Hirota, Exact solution of the Korteweg–de Vries equation for multiple collisions of solitons, *Phys. Rev. Lett.* 27 (1971) 1192–1194.
- [12] R. Hirota, Y. Ohta, Hierarchies of coupled soliton equations. I, *J. Phys. Soc. Japan* 60 (1991) 798–809.
- [13] R. Hirota, J. Satsuma, A variety of nonlinear network equations generated from the Bäcklund transformation for the Toda lattice, *Suppl. Progr. Theoret. Phys.* 59 (1976) 64–100.
- [14] H.C. Hu, X.Y. Tang, S.Y. Lou and Q.P. Liu, *Chaos Solitons Fractals* 22 (2004) 327.
- [15] H.M. Jaradat, S. Al-Shara', F. Awawdeh, M. Alquran, Variable coefficient equations of the Kadomtsev-Petviashvili hierarchy: multiple soliton solutions and singular multiple soliton solutions, *Phys. Scr.* 85 (2012) 035001.
- [16] E.J. Parkes, B.R. Duffy, An automated tanh-function method for finding solitary wave solutions to non-linear evolution equations, *Commun. Comput. Phys.* 98 (1996) 288–300.
- [17] A. Wazwaz, Multiple soliton solutions for the (2+1)-dimensional asymmetric Nizhnik-Novikov-Veselov equation, *Nonlinear Analysis* 72 (2010) 1314-1318.
- [18] G. Yu and H. Tam, A vector asymmetrical NNV equation: Soliton solutions, bilinear Bäcklund transformation and Lax pair, *J. Math. Anal. Appl.* 344 (2008) 593–600.
- [19] F. Zhaosheng, Comment on “On the extended applications of homogeneous balance method”, *Appl. Math. Comput.* 158 (2) (2004) 593–596.
- [20] J. Zhang, Homogeneous balance method and chaotic and fractal solutions for the Nizhnik–Novikov–Veselov equation, *Physics Letters A* 313 (2003) 401–407.