

Comparison of Indoor and Outdoor Air Quality in Children Homes at Prenatal Period and One Year Old

S. Lakestani, B. Karakas, S. Acar Vaizoglu, B. Guciz Dogan, C. Guler, B. Sekerel, A. Taner, and G. Gullu

Abstract—Indoor air (VOCs) samples were collected simultaneously from variety of indoors (e.g. living rooms, baby's rooms) and outdoor environments which were voluntarily selected from the houses in which pregnant residents live throughout Ankara. This is the first comprehensive study done in Turkey starting from prenatal period and continued till the babies had one year old. VOCs levels were measured over 76 homes. Air samples were collected in Tenax TA sorbent filled tubes with active sampling method and analyzed with Thermal Desorber and Gas Chromatography/Mass spectrometry (TD-GC/MS). At the first sampling period in the baby's rooms maximum concentration of toluene was measured about $240.77\mu\text{g.m}^{-3}$ and in the living rooms maximum concentration of naphthalene was $180.24\mu\text{g.m}^{-3}$. At the second sampling period in the baby's rooms maximum concentration of toluene was measured about $144.97\mu\text{g.m}^{-3}$ and in the living rooms maximum concentration of naphthalene was $247.89\mu\text{g.m}^{-3}$. Concentration of TVOCs in the first period was generally higher than the second period.

Keywords—Indoor Air, Volatile Organic Compounds (VOCs), Gas Chromatography.

I. INTRODUCTION

VOLATILE Organic Compounds (VOCs) are a large group of carbon-based chemicals that easily evaporate at room temperature. They have been associated with a number of health problems that include cancer [1], [2] and the induction of acute and chronic health pathologies [3].

There are hundreds of VOCs in the air, which increases the complexion of indoor air pollution. VOCs emitted from various sources such as building materials [4]-[9] household materials [10] and combusted materials [11]-[14]. VOCs are widely used in many household products and are emitted by paints [15]-[17] adhesives [18] waxes, solvents, detergents, woods [19]-[20] and items containing them, including carpets

S. Lakestani, B. Karakas, and G. Gullu are with the Faculty of Engineering, Environmental Engineering Department; Hacettepe University, 06800 Beytepe, Ankara Turkey (e-mail: sanaz@hacettepe.edu.tr, bilgekarakas@hacettepe.edu.tr, ggullu@hacettepe.edu.tr).

S. Acar Vaizoglu, B. Guciz Dogan, and C.Guler are with the Faculty of Medicine, Department of Public Health; Hacettepe University, 06100 Sıhhiye, Ankara Turkey (e-mail: sacar@hacettepe.edu.tr, bdogan@hacettepe.edu.tr, cguler@hacettepe.edu.tr).

B. Sekerel is with Faculty of Medicine, Pediatric Allergy and Asthma Unit; Hacettepe University, 06100 Sıhhiye, Ankara Turkey (e-mail: bsekerel@yahoo.com).

A. Taner is with TOBB ETÜ Hospital, Microbiology and Clinical Microbiology; Yasam Caddesi No: 5 06510 Söğütözü, Ankara Turkey (e-mail: ataner@tobbetuhastanesi.edu.tr).

[21] vinyl flooring [22], [23] air-conditioners [24] newspapers [25] printers and photocopiers [26].

Combined indoor/outdoor air quality measurements have shown that there exist significant VOC sources indoors.

People spend 80–90% of their time in enclosed spaces (e.g., houses, office buildings, and schools). Several researches have demonstrated that babies and children are more sensitive to environmental pollution than adults. Babies spend about 95% of their time in indoor environment. The time spend in indoor might be more during winter times. The health effects of indoor air pollutants are not fully understood, but indoor air quality has been linked with a wide array of health outcomes including deficits in lung function, chronic respiratory disease, lung cancer, heart disease, developmental disorders, and damage to the brain, nervous system, liver, or kidneys [27], [28].

Benzene, toluene, ethyl benzene, ortho, meta, and para-xylene (BTEX) are mono aromatic hydrocarbons belonging to the family of volatile organic compounds (VOCs) [29]. Ambient air concentration of some of these compounds, most notably benzene, is regulated by law due to their toxicity [30]. Benzene (C₆H₆), Toluene (C₇H₈) and xylene (C₈H₁₀) are found in the vapor of products such as gasoline, oils, paints, glues, inks, plastics, and rubber, where they are used as solvents. These three pollutants also enter into the composition of detergents, explosives, pharmaceuticals, foams and dyes.

II. MATERIAL AND METHOD

A. Study Site and Sampling Strategy

The study undertook at the 76 houses in which pregnant woman and their babies live in. The sampling was conducted in spring, during two periods from May 2011 and from May 2012. Indoor and outdoor levels of volatile organic compounds (VOC) were determined. The measurements are done in two periods, during the third three-master of pregnancy, and when the babies are one year old in different areas of Ankara.

The questionnaire about building conditions and household characteristics (e.g., type of heating fuel) and number of occupants in homes was completed by the participants during the sampling period.

B. Sampling

The air sampling tubes was placed in the middle of the sampled rooms in the 50cm above the floor on a horizontal surface.

Previously to sampling or calibration, all sampling tubes were conditioned at 280°C by forcing pure nitrogen gas through the tubes at a flow rate of 75ml min⁻¹ for 5h and then analyzed to certify that no target pollutants were present.

Samples of VOCs were collected by drawing air through a stainless steel tube containing 500mg of Tenax TA using a personal air sampling pump (SKC Pump) at a flow rate of 80-85ml min⁻¹ for a period of 45min.

C. Chemical Analysis

7 VOCs were selected for chemical analysis in this study. The description of sampling and species analysis methods is shown in Table I.

TABLE I
SAMPLING AND SPECIES ANALYSIS

sampler	Active Tenax tube
Target compounds (7 compounds)	Benzene, Toluene, m,p,o-Xylene, Ethyl benzene, Naphthalene
Sampling method	Active sampling for 45min
Desorption method	Solvent desorption (Tenax disulfide) Thermal Desorber-Gas chromatography/Mass Spectroscopy
Analytical method (TD-GC/MS)	Column: DB-1 (Agilent J&W Columns, part number 123-1063) ID: 0,32mm, length:60m, film 1µm

Air samples were collected in Tenax TA sorbent filled tubes with active sampling method and analyzed with Thermal Desorber (Tekmar Dohrmann AERO Trap6000)-Gas chromatography (Agilent 6890N)/Mass Spectroscopy (Agilent 5975C) Detector (TD-GC/MS). The sum of the concentration of the 7 target VOCs was defined as TVOC.

III. RESULTS AND DISCUSSION

A. Indoor Concentrations

Monitoring for indoor air quality was conducted in the rooms in which the infant spends the most time for 76 homes in Ankara. Tables II and III summarize the mean, median and TVOCs levels in each home in the May 2011.

TABLE II
THE MEAN MEAN± STD, MEDIAN, MAXIMUM, MINIMUM, AND TVOCs¹ IN MAY 2011 (PRENATAL PERIOD)

Area		Living Room I (µg.m ⁻³)			
Sample	N	AVE	MAX	MIN	MED
Benzene	59	1.27±1.71	10.89	0.18	0.72
Toluene	58	23.56±22.41	91.65	0.36	13.59
Eth.benzene	59	4.36±9.49	68.05	0.01	2.10
m.p.xylene	59	6.63±6.24	35.07	0.21	4.73
o.xylene	59	5.90±7.23	45.82	0.23	3.77
Naphthalene	59	22.45±34.22	180.24	0.54	10.62
TVOCs ¹	59	63.77±52.44	224.18	2.42	50.45

Area		Living + Baby's Room I ² (µg.m ⁻³)			
Sample	N	AVE	MAX	MIN	MED
Benzene	10	1.12±1.12	4.07	0.14	0.75
Toluene	10	39.04±50.78	142.38	4.55	17.57
Eth.benzene	10	1.99±1.44	4.69	0.37	1.55
m.p.xylene	10	5.94±3.92	13.86	1.03	5.25
o.xylene	10	2.60±1.77	5.19	0.55	2.15
Naphthalene	10	11.23±12.88	42.42	1.84	5.63
TVOCs ¹	10	61.93±64.87	207.78	9.64	35.77

1. Total Volatile Organic Compounds (µg.m⁻³)

2. living room and baby's room is the same

Mean of toluene 39.04µg.m⁻³ at houses without baby's room and ethyl benzene 4.36µg.m⁻³ in the living (Table II), m+p+o-xylene 7.96 and 6.88µg.m⁻³ and naphthalene 25.18µg.m⁻³ in the baby's rooms were the highest concentration (Table III).

TABLE III
THE MEAN MEAN± STD, MEDIAN, MAXIMUM, MINIMUM, AND TVOCs¹ IN MAY 2011 (PRENATAL PERIOD)

Area		Baby's Room I (µg.m ⁻³)			
Sample	N	AVE	MAX	MIN	MED
Benzene	59	1.05±1.21	7.90	0.07	0.63
Toluene	59	36.12±43.09	240.77	0.22	23.37
Ethbenzene	58	3.28±2.89	17.01	0.06	2.93
m.p.xylene	59	7.96±11.05	80.36	0.20	6.42
o.xylene	59	6.88±6.42	36.91	0.09	6.17
Naphthalene	59	25.18±30.83	165.92	0.05	14.08
TVOCs ¹	59	80.42±64.75	264.31	2.08	57.78

Area		Outdoor I (µg.m ⁻³)			
Sample	N	AVE	MAX	MIN	MED
Benzene	76	1.68±2.01	17.06	0.03	1.26
Toluene	76	22.08±44.89	296.18	0.28	6.72
Eth.benzene	76	1.65±1.70	11.56	0.04	1.15
m.p.xylene	76	4.16±5.32	34.51	0.05	2.50
o.xylene	76	1.62±2.20	12.84	0.06	0.89
Naphthalene	76	5.69±10.52	72.95	0.16	2.97
TVOCs ¹	76	36.88±57.63	383.00	0.97	16.69

1. Total Volatile Organic Compounds (µg.m⁻³)

Tables IV and V summarize the mean, median and TVOCs levels in each home in the May 2012.

TABLE IV
THE MEAN MEAN± STD, MEDIAN, MAXIMUM, MINIMUM, AND TVOCs¹ IN
MAY 2012 (BABIES ONE YEAR OLD)

Area		Living Room II ($\mu\text{g}\cdot\text{m}^{-3}$)			
Sample	N	AVE	MAX	MIN	MED
Benzene	67	2.93±1.85	8.49	1.04	2.36
Toluene	67	28.15±31.54	137.35	0.11	12.24
Eth.benzene	67	7.14±7.30	51.13	0.97	5.42
m,p.xylene	67	10.74±10.89	62.99	0.78	8.19
o.xylene	67	9.32±9.54	48.44	0.79	6.12
Naphthalene	67	25.94±41.13	247.90	2.10	12.92
TVOCs ¹	67	84.21±64.53	308.86	6.87	64.74

Area		Living + Baby's Room II ($\mu\text{g}\cdot\text{m}^{-3}$)			
Sample	N	AVE	MAX	MIN	MED
Benzene	7	2.30±0.35	3.00	2.05	2.10
Toluene	7	12.37±18.36	50.62	0.56	2.87
Eth.benzene	7	3.77±2.12	7.50	1.32	3.60
m,p.xylene	7	5.72±4.57	14.02	0.78	5.19
o.xylene	7	4.55±3.11	9.02	0.73	4.20
Naphthalene	7	19.59±26.92	78.22	2.16	9.06
TVOCs ¹	7	48.30±52.07	161.43	8.54	34.70

1. Total Volatile Organic Compounds ($\mu\text{g}\cdot\text{m}^{-3}$)
2. living room and baby's room is the same

TABLE V
THE MEAN± STD, MEDIAN, MAXIMUM, MINIMUM, AND TVOCs¹ IN MAY
2012 (BABIES ONE YEAR OLD)

Area		Baby's Room II ($\mu\text{g}\cdot\text{m}^{-3}$)			
Sample	N	AVE	MAX	MIN	MED
Benzene	69	3.40±2.23	9.74	1.00	2.37
Toluene	69	27.69±35.53	144.98	0.14	9.98
Eth.benzene	69	7.46±10.77	79.30	1.31	4.80
m,p.xylene	69	9.62±10.79	58.59	0.75	7.19
o.xylene	69	8.21±9.77	62.54	0.80	5.82
Naphthalene	69	21.94±33.02	204.20	1.60	11.41
TVOCs ¹	69	78.32±61.97	291.58	7.28	52.74

Area		Outdoor II ($\mu\text{g}\cdot\text{m}^{-3}$)			
Sample	N	AVE	MAX	MIN	MED
Benzene	76	3.14±2.20	15.65	1.88	2.29
Toluene	76	11.2±16.58	92.33	0.12	5.29
Ethbenzene	76	3.03±1.73	11.42	1.27	2.63
m,p.xylene	76	3.71±2.98	15.76	0.76	2.99
o.xylene	76	3.04±2.31	12.76	0.70	2.40
Naphthalene	76	5.41±7.74	53.25	2.04	3.27
TVOCs ¹	76	29.53±26.35	158.45	6.80	20.30

1. Total Volatile Organic Compounds ($\mu\text{g}\cdot\text{m}^{-3}$)

In Table IV and V mean of toluene $28.15\mu\text{g}\cdot\text{m}^{-3}$, m+p+o-xylene 11.23 and $9.56\mu\text{g}\cdot\text{m}^{-3}$ and naphthalene $25.9\mu\text{g}\cdot\text{m}^{-3}$ in the living rooms were highest. Warmer temperatures during the summer time will obviously increase the evaporation of VOCs from vehicle fuel tanks.

Concentration of toluene and naphthalene at baby's room in first sampling periods were higher than other VOCs. Household painting and paint, varnish and lacquer removal, tobacco smoke, and consumer products such as adhesives, floor polish, inks, coatings and solvent-thinned products may contain toluene.

In indoor air, emissions from cooking, tobacco smoking, or moth repellants are expected to be the predominant sources of naphthalene. Available data are inadequate to establish a causal association between exposure to naphthalene and cancer in humans. Adequately scaled epidemiologic studies designed to examine a possible association between naphthalene exposure and cancer was not located.

The results clearly show that baby's rooms are not as safe as we think. Open the windows at least twice a day for 15

minutes at a time to help flush out stuffy air (but if it's cold outside, make sure the child is not in the room or is dressed appropriately).

Concentration of Total Volatile Organic Compounds (TVOCs) at the first sampling were higher than the second period, because in the first sampling they had bought new furniture for baby's room.

m+p+o-xylene levels in homes with smokers were significantly higher than that in homes without smokers. Xylenes are also emitted from building materials such as carpet adhesives, vinyl cove adhesive, latex caulk, latex paint, and various moldings. Environmental tobacco smoke is also a common indoor source of xylenes [33].

Our results are relevant because they are the first reference values of VOCs exposure levels in two periods be measured in Ankara-Turkey.

Finally, the identification of the determinants that influence both indoor and outdoor pollution levels is vital for designing programs for controlling exposure to household pollution.

Parents can impact the air their infants breathe in several ways. Air conditioners or humidifiers are classic examples, but they may be too harsh for an infant's system or may be inappropriate for the climatic conditions, depending on where the family lives.

B. Indoor and Outdoor Ratio

The indoor (I) to outdoor (O) ratio of the air concentrations of the different compounds (I/O ratio) is a useful tool to assess the importance of indoor versus outdoor sources better than the absolute concentrations. When the I/O ratio is over 1, the presence of indoor sources is considered to be significant. The I/O ratios for each compound are presented in table VI and VII.

TABLE VI
INDOOR AND OUTDOOR RATIOS IN MAY 2011

	B1/L1	L+B1/O1	B1/O1	L1/O1
benzene	0.83	1.77	1.66	2.01
Toluene	1.53	1.77	1.64	1.07
Ethylbenzene	0.75	1.21	1.99	2.65
m,p.xylene	1.20	1.43	1.91	1.59
o.xylene	1.17	1.60	4.24	3.63
Naphthalene	1.12	1.97	4.43	3.95

The indoor levels of VOCs were higher than those registered outdoors (I/O ratios > 1) indicating that the sources of these compounds are located at the indoor environments. In this case, only benzene and ethyl benzene presented lower than 1, and then B1/L1 ratios, so only these compounds are related to outdoor sources.

TABLE VII
INDOOR AND OUTDOOR RATIOS IN MAY 2012

	B2/L2	L+B2/O2	B2/O2	L2/O2
Benzene	1.12	0.76	1.16	1.04
Toluene	0.99	1.38	2.49	2.53
Ethylbenzene	1.01	1.24	2.50	2.46
m,p.xylene	0.86	1.41	2.61	3.02
o.xylene	0.89	1.35	2.81	3.15
Naphthalene	0.87	3.66	4.19	4.81

In Table VII toluene, m+p+o-xylene and naphthalene ratios in baby's room to living room were lower than 1, then these VOCs sources were from living room.

C. Questionnaire Survey and Results

The questionnaire containing 30 questions about building conditions, residential life-style and indoor situations was completed by the participants during the sampling period. The questionnaire also asked about the distance of the residence from a street with continuous traffic, type of cooking energy, type of heating used in the home, the parents' smoking habits, whether the parents smoked at home.

In addition, the questionnaire asked about any construction or home improvement projects that had been carried out in the home during the child's first year of life (construction, painting, both). Other questions included how often the house was cleaned (more than once a week, once a week, less than once a week). In tables VIII-XI in different rooms, you can see the statistically significant differences between BTEX and factors.

The possible influence of the building age and proximity to the street, cigarette smoking, number of the family member, number of people during sampling, cooking energy, heating energy, naphthalene tablets and formaldehyde, temperature, relative humidity, CO₂ and CO on the indoor VOCs concentrations in homes was discussed (Table VIII, IX).

TABLE VIII
EFFECT OF FACTORS ON BTEX

	PS	BAY	NFM	HE	CE
Benzene	+	+			+
Toluene					
Ethyl benzene					
o-Xylene					
m-p-Xylene			+		
TVOCs			+	+	
	SAH	NT	VC	NPDS	
Benzene			+		
Toluene		+	+	+	
Ethyl benzene	+		+		
o-Xylene	+		+		
m-p-Xylene	+				
TVOCs			+	+	

+: Statistically significant differences ($p < 0.05$), PS : Proximity to the Street, BAY: Building Age _yr, NFM: Number of the Family Members, HE: Heating Energy, CE: Cooking Energy, SAH: Smokers Area at Home, NT: Naphthalene Tablets, VC: Ventilation Condition, NPDS : Number of people during sampling.

TABLE X
COMPARISON OF INDOOR BTEX CONCENTRATIONS WITH OTHER COUNTRIES ($\mu\text{g.M}^{-3}$)

	INMA (Valencia, Spain) ¹ 2006–2007	Shield (Minneapolis, USA) Spring ² 1999–2000	This Study (Turkey) May 2011 Living room	This Study (Turkey) May 2012 Living room
	Median	Median	Median	Median
Benzene	1.04	1.5	0.63	2.36
Toluene	10.1	3.1	23.37	12.24
Ethyl benzene	1.3	0.9	2.93	5.42
o-Xylene	1.9	1.0	6.42	8.19
m-p-Xylene	1.3	2.9	6.17	6.12

[29]¹, [31]²

Thus, the indoor levels of most VOCs in Ankara regions were higher than the other countries. This difference in methodology may have contributed to the higher levels found in our study.

TABLE IX

EFFECT OF FACTORS ON BTEX

	HCHO	CO ₂	CO	T (°C)	R	H
Benzene	+			+		
Toluene		+		+		+
Ethyl benzene	+					+
o-Xylene	+		+			+
m-p-Xylene	+		+			+
TVOCs				+		

+: Statistically significant differences ($p < 0.05$), HCHO: Formaldehyde, R_H: Relative Humidity.

Concentration of benzene was higher in homes near to the street and newly built homes and in homes that use gas energy for cooking.

Number of the family members in homes and number of people during sampling may affect the indoor VOCs levels because of the variations of indoor activities, living habits and home volumes [32]. For example, a home with four people frequently used cleaning products and air fresheners when the occupants were at home. On the other hand, a home with one resident had very few indoor activities. Number of people during sampling when >4 in homes, toluene concentration was the highest.

D. Comparison of Our Study and Other Studies

The number of studies that have measured personal BTEX levels in children is scarce. Table X compares VOCs concentrations in Ankara's homes with those in other countries. It is worth to emphasize that the results were obtained from the measurements in using the same sampling and analytical methods.

Our results thus contribute important information with respect to the BTEX exposure levels of Turkish children.

Moreover, we have been able to identify certain sources and factors associated with this exposure, which should facilitate the planning and implementation of measures to reduce it.

ACKNOWLEDGMENTS

This study was supported by grants obtained by The Scientific and Technological Research Council of Turkey (TÜBİTAK, Project No: 110Y082). We are grateful to all the mothers who are taking part in the Project, especially those who have allowed us access to their homes.

REFERENCES

[1] H. Hellén, H. Hakola, T. Laurila, V. Hiltunen, T. Koskentalo. Aromatic hydrocarbon and methyl tert-butyl ether measurements in ambient air of Helsinki (Finland) using diffusive samplers *The Science of the Total Environment* 298, pp. 5–64, 2002.

[2] T. Ohura, T. Amagai, Y. Senga, M. Fusaya. Organic air pollutants inside and outside residences in Shimizu, Japan: levels, sources and risks *The Science of the Total Environment*, 366, pp. 485–499, 2006.

[3] K.H. Park, W.K. Jo. Personal volatile organic compound (VOC) exposure of children attending elementary schools adjacent to industrial complex *Atmospheric Environment* 38 pp. 1303–1312, 2004.

[4] F. Haghighat, C.S. Lee, W.S. Ghaly. Measurement of diffusion coefficients of VOCs for building materials: review and development of a calculation procedure, *Indoor Air* Vol. 12(2): 81-91, 2002.

[5] F. Haghighat, C.S. Lee, W.S. Ghaly. A study on VOC source and sink behavior in porous building materials - analytical model development and assessment, *Indoor Air* 15(3), 183-196, 2005.

[6] A.S. Claeson, M. Sandstrom, A.-L. Sunesson,. Volatile organic compounds (VOCs) emitted from materials collected from buildings affected by microorganisms, *J. Environ. Monit* 9, 240-245, 2007.

[7] J. Nicolle, V. Desauziers, P. Mocho. Solid phase microextraction sampling for a rapid and simple on-site evaluation of volatile organic compounds emitted from building materials, *J. Chromatogr.* 1208(1-2), 10-15, 2008.

[8] K.H. Han, J.S. Zhang, P. Wargocki, H.N. Knudsen, B. Guo. Determination of material emission signatures by PTR-MS and their correlations with odor assessments by human subjects, *Indoor Air* 20(4), 341-354, 2010.

[9] C. Jia, S. Batterman, C. Godwin, S. Charles, J.Y. Chin. Sources and migration of volatile organic compounds in mixed-use buildings, *Indoor Air* 20(5), 357-369, 2010.

[10] K.D. Kwon, W.K. Jo, H.J. Lim, W.S. Jeong. Volatile pollutants emitted from selected liquid household products, *Environ. Sci. Pollut. Res. Int* 15(6), 521-526, 2008.

[11] W. Liu, J. Zhang, J.H. Hashim, J. Jalaudin, Z. Hashim, B.D. Goldstein,. Mosquito coil emissions and health implications, *Environ. Health Perspect* 111(12), 1454-1460, 2003.

[12] Q. Ye, Development of solid-phase microextraction followed by gas chromatography-mass spectrometry for rapid analysis of volatile organic chemicals in mainstream cigarette smoke, *J. Chromatogr* 1213(2), 239-244, 2008.

[13] H. Fromme, S. Dietrich, D. Heitmann, H. Dressel, J. Diemer, T. Schulz, R.A. Jörres, K. Berlin, W. Völkel. Indoor air contamination during a waterpipe (narghile) smoking session, *Food Chem. Toxicol* 47(7), 1636-1641, 2009.

[14] E. Kabir, K.H. Kim. An investigation on hazardous and odorous pollutant emission during cooking activities, *J. Hazard. Mater* 188(1-3), 443-454, 2011.

[15] A. Afshari, B. Lundgren, L.E. Ekberg. Comparison of three small chamber test methods for the measurement of VOC emission rates from paint, *Indoor Air* Vol. 13(2), 156-165, 2003.

[16] G. Wieslander, D. Norbäck. Ocular symptoms tear film stability, nasal patency, and biomarkers in nasal lavage in indoor painters in relation to emissions from waterbased paint, *Int. Arch. Occup. Environ. Health* Vol. 83(7): 733-741, 2010.

[17] Y.M. Chang, W.H. Hu, W.B. Fang, S.S. Chen, C.T. Chang, H.W. Ching. A study on dynamic volatile organic compound emission characterization of water-based paints, *J. Air Waste Manag. Assoc.* Vol. 61(1), 35-45, 2011.

[18] O. Wilke, O. Jann, D. Brödner. VOC- and SVOC-emissions from adhesives, floor coverings and complete floor structures, *Indoor Air* Vol. 14 Suppl 8: 98-107, 2004.

[19] L.K. Jensen, A. Larsen, L. Mølhave, M.K. Hansen, B. Knudsen. Health evaluation of volatile organic compound (VOC) emissions from wood and wood-based materials, *Arch. Environ. Health.* Vol. 56(5), 419-432, 2001.

[20] L. Kirkeskov, T. Witterseh, L.W. Funch, E. Kristiansen, L. Mølhave, M.K. Hansen, B. Knudsen. Health evaluation of volatile organic compound (VOC) emission from exotic wood products, *Indoor Air* Vol. 19(1), 45-57, 2009.

[21] A. Katsoyiannis, P. Leva, D. Kotzias. VOC and carbonyl emissions from carpets: a comparative study using four types of environmental chambers, *J. Hazard Mater.* Vol. 152(2), 669-676, 2008.

[22] S.S. Cox, J.C. Little, A.T. Hodgson. Measuring concentrations of volatile organic compounds in vinyl flooring, *J. Air Waste Manag. Assoc.* Vol. 51(8), 1195-1201, 2001.

[23] Cox S.S., Little J.C. & Hodgson A.T. Predicting the emission rate of volatile organic compounds from vinyl flooring, *Environ. Sci. Technol.* Vol. 36(4), 709-714, 2002.

[24] K.W. Tham, M.S. Zuraimi, S.C. Sekhar. Emission modelling and validation of VOCs' source strengths in air-conditioned office premises, *Environ. Int.* Vol. 30(8), 1075-1088, 2004.

[25] M. Caselli, G. de Gennaro, M.R. Saracino, M. Tutino. Indoor contaminants from newspapers: VOCs emissions in newspaper stands, *Environ. Res.* Vol. 109(2), 149-157, 2009.

[26] C.W. Lee, Y.T. Dai, C.H. Chien, D.J. Hsu. Characteristics and health impacts of volatile organic compounds in photocopy centers, *Environ. Res.* 100(2), 139-149, 2006.

[27] C.J. Weschler. Chemistry in indoor environments: 20 years of research. *Indoor Air* 2011, 21, 205-218.

[28] L. Wang, K.E. Pinkerton. Air pollutant effects on fetal and early postnatal development. *Birth Defects Res. Part C, Embryo Today: Rev.* 81, 144-154, 2007.

[29] A. Esplugues, F. Ballester, M. Estarlich, S. Llop, V.F. Leonarte, E. Mantilla, C. Iñiguez. Indoor and outdoor air concentrations of BTEX and determinants in a cohort of One-Year old children in valencia, Spain. *Science Of The Total Environment.* 409, 63–69, 2010.

[30] European Commission (2009) Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. Available online at: <http://ec.europa.eu/environment/air/legis.htm>.

[31] J.L. Adgate, T.R. Church, A.D. Ryan, G. Ramachandran, A.L. Fredrickson, T.H. Stock, et al. Outdoor, indoor, and personal exposure to VOCs in children. *Environ Health Perspect* 112, 1386–92, 2004.

[32] A.L. Clobes, G.P. Ananth, A.L. Hood, J.A. Schroeder, K.A. Lee. (1992) Human activities as sources of volatile organic compounds in residential environments, *Ann. NY Acad. Sci.*, 641, 79–86 doi:10.1111/j.1749-6632.1992.tb16534.x.

[33] A.T. Hodgson, J.D. Wooley. Assessment of Indoor Concentrations, Indoor Sources and Source Emissions of Selected Volatile Organic Compounds. Final Report, Contract No. A933□063, California Air Resources Board, Research Division, 179 p, 1991.



Sanaz LAKESTANI was born in Iran, on September, 1975. LAKESTANI received a Bs degree in Chemistry in 1997 from Shahid Beheshti University, Tehran-IRAN. In 2001 received a master's degree in Marine Chemistry from Islamic Azad University North Tehran, Tehran-IRAN. Since 2008 she is as a PhD student in Department of Environmental Engineering at Hacettepe University, Ankara-TURKEY. She currently studies under the supervision of Prof. Dr. Gulen Gullu on the indoor and outdoor air quality.

She is PhD student and works as a project assistant in Hacettepe University, department of environmental engineering, Ankara- Turkey. She worked as researcher and development manager at laboratory of chemical factory between 2001- 2007. Now she works in project named "Investigation of relationship between lower respiratory and allergic symptoms and indoor air pollution in children from prenatal period to two years of age". In this project she has measured VOCs (TD-GC-MS/MS) of indoor and outdoor environments where infants live.



B. Karakas comes from Gemerek, Sivas and born in October, 1985. She has got her Bs degree in the department of Environmental Engineering in 2008 from Ondokuz Mayıs

University, Samsun-TURKEY. She is an Ms student candidate in Hacettepe University, Department of Environmental Engineering, Ankara-TURKEY. She currently studies under the supervision of Prof. Dr. Gulen Gullu on the indoor and outdoor air quality. She is also working on a project under Prof. Dr. Gulen Gullu investigating of relationship between lower respiratory and allergic symptoms and indoor air pollution in children from prenatal period to two years of age.

She is working as a project specialist at AECOM Turkey-is a global provider of professional technical and management support services to a broad range of markets, including transportation, facilities, environmental, energy, water and government.

Ms. Karakas have been a member of the chamber of TMMB Environmental Engineers.

Gulen GULLU was born in Turkey on July; 1965. Gullu received a Bs degree in 1987 and a master's degree in 1989 and a PhD degree in 1996 from Middle East Technical University, Department of Environmental Engineering, Ankara-TURKEY.

She worked as Associate Professors, in 1999 at Hacettepe University, in Environmental Engineering Department. She became a Professor in 2006. She currently is the head of the Department of Environmental Engineering at Hacettepe University, Ankara- TURKEY. She works on indoor and outdoor air quality control.