

# Maximum Norm Analysis of a Nonmatching Grids Method for Nonlinear Elliptic Boundary Value Problem $-\Delta u = f(u)$

Abida Harbi

*Abstract*—We provide a maximum norm analysis of a finite element Schwarz alternating method for a nonlinear elliptic boundary value problem of the form  $-\Delta u = f(u)$ , on two overlapping sub domains with non matching grids. We consider a domain which is the union of two overlapping sub domains where each sub domain has its own independently generated grid. The two meshes being mutually independent on the overlap region, a triangle belonging to one triangulation does not necessarily belong to the other one. Under a Lipschitz assumption on the nonlinearity, we establish, on each sub domain, an optimal  $L^\infty$  error estimate between the discrete Schwarz sequence and the exact solution of the boundary value problem.

*Keywords*—Error estimates, Finite elements, Nonlinear PDEs, Schwarz method.

## I. INTRODUCTION

THE Schwarz alternating method can be used to solve elliptic boundary value problems on domains which consist of two or more overlapping subdomains. The solution is approximated by an infinite sequence of functions which results from solving a sequence of elliptic boundary value problems in each of the subdomains. Extensive analysis of Schwarz alternating method for nonlinear elliptic boundary value problems can be found in [8], [9], [10] and [11] and the references therein. In this paper, we are interested in the error analysis in the maximum norm for a class of nonlinear elliptic boundary value problems in the context of overlapping nonmatching grids: we consider a domain which is the union of two overlapping subdomains where each subdomain has its own triangulation. Quite a few works on maximum norm error analysis of overlapping nonmatching grids methods for elliptic problems are known in the literature cf., e.g., [1], [2], [4], [6] and [12]. To prove the main result of this paper, we proceed as in [2] and [6]. More precisely, we develop an approach which combines a geometrical convergence result due to [8] and a lemma which consists of estimating the error in the maximum norm between the continuous and discrete Schwarz iterates. The optimal convergence order is then derived making use of standard finite element  $L^\infty$ -error estimate for linear elliptic equations. In the present paper, the proof of this lemma stands on a Lipschitz continuous dependency with respect to both the boundary condition and the source term for linear elliptic equations see [6]. Now, we give an outline of the paper. In

Section 2 we state a continuous alternating Schwarz sequences and define their respective finite element counterparts in the context of nonmatching overlapping grids. Section 3 is devoted to the  $L^\infty$ -error analysis of the method.

mds  
 March 30, 2013

## II. PRELIMINARIES

We begin by laying down some definitions and classical results related to linear elliptic equations.

### A. Linear Elliptic Equations

Let  $\Omega$  be a bounded polyhedral domain of  $\mathbb{R}^2$  or  $\mathbb{R}^3$  with sufficiently smooth boundary  $\partial\Omega$ . We consider the bilinear form

$$a(u, v) = \int_{\Omega} (\nabla u \cdot \nabla v) dx \quad (1)$$

the linear form

$$(f, v) = \int_{\Omega} f(x) \cdot v(x) dx, \quad (2)$$

the right hand side  $f$ , a regular function and the space

$$V^{(g)} = \{v \in H^1(\Omega) \text{ such that } v = g \text{ on } \partial\Omega\}, \quad (3)$$

where  $g$  is a regular function defined on  $\partial\Omega$ . We consider the linear elliptic equation: Find  $\zeta \in V^{(g)}$  such that:

$$a(\zeta, v) + c(\zeta, v) = (f, v), \forall v \in V^{(g)}, \quad (4)$$

where  $c \in \mathfrak{R}$   $c > 0$ , such that

$$c \geq \beta > 0 \quad (5)$$

Let  $V_h$  be the space of finite elements consisting of continuous piecewise linear functions  $v$  vanishing on  $\partial\Omega$  and  $s, s = 1, 2, \dots, m(h)$  be the basis function of  $V_h$ . The discrete counterpart of (4) consists of finding  $\zeta_h \in V_h^{(g)}$  such that

$$a(\zeta_h, v) + c(\zeta_h, v) = (f, v), \forall v \in V_h^{(g)} \quad (6)$$

where

$$V_h^{(g)} = \{v \in V_h : v = \pi_h(g) \text{ on } \partial\Omega\} \quad (7)$$

and  $\pi_h$  is an interpolation operator on  $\partial\Omega$ .

**Theorem 2.1** (cf.[13]) Under suitable regularity of the solution of problem (4), there exists a constant  $C$  independent of  $h$  such that

$$\|\zeta - \zeta_h\|_{L^\infty(\Omega)} \leq Ch^2 |\log h|. \quad (8)$$

**Lemma 2.1** (cf.[11]) Let  $w \in H_1(\Omega) \cap C(\Omega)$  satisfy  $a(w, \Phi) + c(w, \Phi) \geq 0$  for all non-negative  $\Phi \in H_0^1(\Omega)$  and  $w \geq 0$  on  $\partial\Omega$ . Then  $w \geq 0$  on  $\bar{\Omega}$ .

The proposition below establishes a Lipschitz continuous dependency of the solution with respect to the data. Let  $(f; g); (\tilde{f}, \tilde{g})$  be a pair of data, and  $\zeta = \sigma(f, g); \tilde{\zeta} = \sigma(\tilde{f}, \tilde{g})$  the corresponding solutions to (4).

**Proposition 2.1** (cf.[6]) Under conditions of the preceding lemma, we have:

$$\|\zeta - \tilde{\zeta}\|_{L^\infty(\Omega)} \leq \max\left\{\left(\frac{1}{\beta}\right)\|f - \tilde{f}\|_{L^\infty(\Omega)}; \|g - \tilde{g}\|_{L^\infty(\partial\Omega)}\right\}. \quad (9)$$

**Remark 2.1** Lemma 2.1 stays true in the discrete case. Indeed, assume that the discrete maximum principle (d.m.p) hold [5], [7] that is the matrix resulting from the finite element discretization is an M-Matrix. Then we have:

**Lemma 2.2** Let  $w \in V_h$  satisfy  $a(w, s) + c(w, s) \geq 0$  for all  $s = 1, 2, \dots, m(h)$  and  $w \geq 0$  on  $\partial\Omega$ . Then  $w \geq 0$  on  $\bar{\Omega}$ .

Let  $(f, g); (\tilde{f}, \tilde{g})$  be a pair of data, and  $\zeta_h = \sigma_h(f, g); \tilde{\zeta}_h = \sigma_h(\tilde{f}, \tilde{g})$  the corresponding solutions to (6);

**Proposition 2.2** Let the d.m.p hold. Then, under conditions of lemma 2.2, we have

$$\|\zeta_h - \tilde{\zeta}_h\|_{L^\infty(\Omega)} \leq \max\left\{\left(\frac{1}{\beta}\right)\|f - \tilde{f}\|_{L^\infty(\Omega)}; \|g - \tilde{g}\|_{L^\infty(\partial\Omega)}\right\}. \quad (10)$$

### III. SCHWARZ ALTERNATING METHODS FOR NONLINEAR ELLIPTIC BOUNDARY VALUE PROBLEM

Consider the nonlinear elliptic boundary value problem

$$-\Delta u = f(u) \text{ in } \Omega, \quad (11)$$

$$u = 0 \text{ on } \partial\Omega \quad (12)$$

where  $f(\cdot)$  is a nondecreasing nonlinearity. Thanks to [3], problem (11) has a unique solution. Let us also assume that  $f(\cdot)$  is a Lipschitz continuous on  $\mathfrak{R}$ ; that is

$$|f(x) - f(y)| \leq k|x - y|, \forall x, y \in \mathfrak{R} \quad (13)$$

The problem (11) is equivalent to

$$-\Delta u + cu = F(u) \text{ in } \Omega \quad (14)$$

$$u = 0 \text{ on } \partial\Omega \quad (15)$$

where the functional  $F$  is defined by:  $F(u) = f(u) + cu$  and its Lipschitz constant is  $K = k + c$ . Evidently we have

$$\beta \leq c < K. \quad (16)$$

The related weak form of (14) is defined by

$$a(u, v) + c(u, v) = (F, v), \forall v \in H_0^1(\Omega), \quad (17)$$

**Remark 2.2** The main idea of this work consists to applied the multiplicative Schwarz algorithm to the problem (14) equivalent to (11). We decompose  $\Omega$  into two overlapping

smooth subdomains  $\Omega_1$  and  $\Omega_2$  such that:  $\Omega = \Omega_1 \cup \Omega_2$ . We denote by  $\partial\Omega_i$  the boundary of  $\Omega_i$  and  $\Gamma_i = \partial\Omega_i \cap \Omega_j$ , We assume that the intersection of  $\Gamma_i$  and  $\Gamma_j, i \neq j$  is empty. Let

$$V_i^{(w_j)} = \{v \in H^1(\Omega_i) \text{ such that } v = w_j \text{ on } \Gamma_i\} \quad (18)$$

We associate with problem (17) the following system: Find  $(u_1, u_2) \in V_1^{(u_2)} \times V_2^{(u_1)}$  solution of

$$a_1(u_1, v) + c(u_1, v) = (F(u_1), v), \forall v \in V_1^{(u_2)}, \quad (19)$$

$$a_2(u_2, v) + c(u_2, v) = (F(u_2), v), \forall v \in V_2^{(u_1)}, \quad (20)$$

where

$$a_i(u, v) = \int_{\Omega_i} (\nabla u \cdot \nabla v) dx; \quad (21)$$

$$u_i = u/\Omega_i, i = 1, 2. \quad (22)$$

#### A. The Continuous Schwarz Sequences

Let  $u^0$  be an initialization in  $C^0(\Omega)$  (i.e. continuous functions vanishing on  $\partial\Omega$ ) such that

$$a(u^0, v) + c(u^0, v) = (F, v), \forall v \in H_0^1(\Omega). \quad (23)$$

$u_2^0 = u^0/\Omega_2$ , we respectively define the alternating Schwarz sequences  $(u_1^{n+1})$  on  $\Omega_1$  such that  $u_1^{n+1} \in V_1^{(u_2^n)}$  solves  $n \geq 0$

$$a_1(u_1^{n+1}, v) + c(u_1^{n+1}, v) = (F(u_1^{n+1}), v) \forall v \in V_1^{(u_2^n)}; \quad (24)$$

and  $(u_2^{n+1})$  on  $\Omega_2$  such that  $u_2^{n+1} \in V_2^{(u_1^{n+1})}$  solves  $n \geq 0$

$$a_2(u_2^{n+1}, v) + c(u_2^{n+1}, v) = (F(u_2^{n+1}), v) \forall v \in V_2^{(u_1^{n+1})}; \quad (25)$$

The following is a geometrical convergence of the Schwarz sequences.

**Theorem 2.2** (cf. [8] pp. 51-63) The sequences  $(u_1^{n+1}); (u_2^{n+1}); n \geq 0$  produced by the Schwarz alternating method converge geometrically to the solution  $(u_1, u_2)$  of the system (19) (20). More precisely, there exist two constants  $k_1, k_2 \in ]0; 1[$  which depend on  $(\Omega_1, \Gamma_2)$  and  $(\Omega_2, \Gamma_1)$  respectively, such that for all  $n \geq 0$ ;

$$\|u_1 - u_1^{n+1}\|_{L^\infty(\Omega_1)} \leq k_1^n k_2^n \|u^0 - u\|_{(\Gamma_1)} \quad (26)$$

$$\|u_2 - u_2^{n+1}\|_{L^\infty(\Omega_2)} \leq k_1^{n+1} k_2^n \|u^0 - u\|_{(\Gamma_2)}. \quad (27)$$

#### B. The Discretization

For  $i = 1, 2$ , let  $\tau^{h_i}$  be a standard regular and quasi-uniform finite element triangulation in  $\Omega_i$ ;  $h_i$ , being the mesh size. The two meshes being mutually independent on  $\Omega_1 \cap \Omega_2$ , a triangle belonging to one triangulation does not necessarily belong to the other. We consider the following discrete spaces:

$$V_{h_i} = \{v \in C(\bar{\Omega}_i) \cap H_0^1(\Omega_i) \text{ such that } v/K \in P_1, \forall K \in \tau^{h_i}\} \quad (28)$$

and for every  $w \in C(\bar{\Gamma}_i)$ , we set:

$$V_{h_i}^{(w)} = \{v \in V_{h_i} : v = 0 \text{ on } \partial\Omega_i \cap \partial\Omega; v = \pi_{h_i}(w) \text{ on } \Gamma_i\} \quad (29)$$

where  $\pi_{h_i}$  denote the interpolation operator on  $\Gamma_i$ .

1) *The Discrete Maximum Principle:* ([5], [7]) We assume that the respective matrices resulting from the discretizations of problems (24) and (25) are M-matrices. Note that, as the two meshes  $h_1$  and  $h_2$  are independent over the overlapping subdomains, it is impossible to formulate a global approximate problem which would be the direct discrete counterpart of problem (14).

### C. The Discrete Schwarz Sequences

Now, we define the discrete counterparts of the continuous Schwarz sequences defined in (24) and (25). Indeed, let  $u_{h_1}^0$  be the discrete analog of  $u^0$ , defined in (23), let  $u_{h_1}^{n+1} \in V_{h_1}^{(u_{h_2}^n)}$  such that

$$a_1(u_{h_1}^{n+1}, v) + c(u_{h_1}^{n+1}, v) = (F(u_{h_1}^{n+1}), v) : \forall v \in V_{h_1}^{(u_{h_2}^n)}; n \geq 0 \quad (30)$$

and  $u_{h_2}^{n+1} \in V_{h_2}^{(u_{h_1}^{n+1})}$  such that

$$a_2(u_{h_2}^{n+1}, v) + c(u_{h_2}^{n+1}, v) = (F(u_{h_2}^{n+1}), v) : \forall v \in V_{h_2}^{(u_{h_1}^{n+1})}; n \geq 0 \quad (31)$$

## IV. $L^\infty$ - ERROR ANALYSIS

This section is devoted to the proof of the main result of the present paper. To that end we begin by introducing two discrete auxiliary sequences and prove a fundamental lemma.

### A. Two Auxiliary Schwarz Sequences

For  $w_{h_2}^0 = u_{h_2}^0$ , we define the sequences  $(w_{h_1}^{n+1})$  such that  $u_{h_1}^{n+1} \in V_{h_1}^{(u_{h_2}^n)}$  solves

$$a_1(w_{h_1}^{n+1}, v) + c(w_{h_1}^{n+1}, v) = (F(u_{h_1}^{n+1}), v) : \forall v \in V_{h_1}^{(u_{h_2}^n)} \quad (32)$$

and  $(w_{h_2}^{n+1})$  such that  $w_{h_2}^{n+1} \in V_{h_2}^{(u_{h_1}^{n+1})}$  solves

$$a_1(w_{h_2}^{n+1}, v) + c(w_{h_2}^{n+1}, v) = (F(u_{h_2}^{n+1}), v) : \forall v \in V_{h_2}^{(u_{h_1}^{n+1})} \quad (33)$$

It is then clear that  $w_{h_1}^{n+1}$  and  $w_{h_2}^{n+1}$  are the finite element approximation of  $u_1^{n+1}$  and  $u_2^{n+1}$  defined in (24), (25), respectively. Then, as  $F(\cdot)$  is continuous,  $\|F(u_i^n)\|_\infty \leq C$  ( $C$  independent of  $n$ ) and, therefore, making use of standard maximum norm estimates for linear elliptic problems, we have

$$\|u_i^n - w_{h_i}^n\|_{L^\infty(\Omega_i)} \leq Ch_i^2 |\log h_i|. \quad (34)$$

**Notation 3.1** From now on, we shall adopt the following notations:

$$\|\cdot\|_1 = \|\cdot\|_{L^\infty(\Omega_1)}, \|\cdot\|_2 = \|\cdot\|_{L^\infty(\Omega_2)} \quad (35)$$

### B. The Main Results

The following lemma will play a key role in proving the main result of this paper.

**Lemma 3.1** Let  $u_i^{n+1}$  the continuous Schwarz sequence and  $u_{h_i}^{n+1}$  its discrete counterpart then we have

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \sum_{i=1}^{n+1} \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^n \|u_2^i - w_{h_2}^i\|_2 \quad (36)$$

$$\|u_2^{n+1} - u_{h_2}^{n+1}\|_2 \leq \sum_{i=0}^{n+1} \|u_2^i - w_{h_2}^i\|_2 + \sum_{i=1}^{n+1} \|u_1^i - w_{h_1}^i\|_1 \quad (37)$$

**Proof** Let us now prove (36) and (37) simultaneously by induction. Indeed for  $n = 1$ , using the Proposition 2.2, we have in domain 1

$$\begin{aligned} \|u_1^1 - u_{h_1}^1\|_1 &\leq \|u_1^1 - w_{h_1}^1\|_1 + \|w_{h_1}^1 - u_{h_1}^1\|_1 \\ &\leq \|u_1^1 - w_{h_1}^1\|_1 \\ &\quad + \max\left\{\left(\frac{1}{\beta}\right)\|F(u_1^1) - F(u_{h_1}^1)\|_1; |u_2^0 - u_{h_2}^0|_2\right\} \end{aligned}$$

then

$$\begin{aligned} \|u_1^1 - u_{h_1}^1\|_1 &\leq \|u_1^1 - w_{h_1}^1\|_1 \\ &\quad + \max\left\{\left(\frac{1}{\beta}\right)\|F(u_1^1) - F(u_{h_1}^1)\|_1; |u_2^0 - u_{h_2}^0|_2\right\} \end{aligned}$$

thus

$$\begin{aligned} \|u_1^1 - u_{h_1}^1\|_1 &\leq \|u_1^1 - w_{h_1}^1\|_1 \\ &\quad + \max\{\rho\|u_1^1 - u_{h_1}^1\|_1; \|u_2^0 - u_{h_2}^0\|_2\} \end{aligned}$$

where

$$\rho = \frac{K}{\beta} > 1 \quad (38)$$

We then have to distinguish between two cases:

$$(1) : \max\{\rho\|u_1^1 - u_{h_1}^1\|_1; \|u_2^0 - u_{h_2}^0\|_2\} = \rho\|u_1^1 - u_{h_1}^1\|_1 \quad (39)$$

or

$$(2) : \max\{\rho\|u_1^1 - u_{h_1}^1\|_1; \|u_2^0 - u_{h_2}^0\|_2\} = \|u_2^0 - u_{h_2}^0\|_2 \quad (40)$$

Case 1 implies

$$\|u_1^1 - u_{h_1}^1\|_1 \leq \|u_1^1 - w_{h_1}^1\|_1 + \rho\|u_1^1 - u_{h_1}^1\|_1 \quad (41)$$

$$\|u_2^0 - u_{h_2}^0\|_2 \leq \rho\|u_1^1 - u_{h_1}^1\|_1 \quad (42)$$

By adding  $\|u_1^1 - w_{h_1}^1\|_1$  in (42) we get

$$\|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1 \leq \|u_1^1 - w_{h_1}^1\|_1 + \rho\|u_1^1 - u_{h_1}^1\|_1 \quad (43)$$

So  $\|u_1^1 - w_{h_1}^1\|_1 + \rho\|u_1^1 - u_{h_1}^1\|_1$  is bounded below by  $\|u_1^1 - u_{h_1}^1\|_1$  and  $\|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1$ . This implies that

$$(a) : \|u_1^1 - u_{h_1}^1\|_1 \leq \|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1 \quad (44)$$

or

$$(b) : \|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1 \leq \|u_1^1 - u_{h_1}^1\|_1 \quad (45)$$

Which implies

$$(a) : \|u_1^1 - u_{h_1}^1\|_1 \leq \|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1 \leq \|u_1^1 - w_{h_1}^1\|_1 + \rho\|u_1^1 - u_{h_1}^1\|_1$$

or

$$(b) : \|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1 \leq \|u_1^1 - u_{h_1}^1\|_1 \leq \|u_1^1 - w_{h_1}^1\|_1 + \rho\|u_1^1 - u_{h_1}^1\|_1$$

Hence (a) and (b) are true because they both coincide with (41). So, there is either a contradiction and thus case 1 is impossible or case 1 is possible only if

$$\|u_1^1 - u_{h_1}^1\|_1 = \|u_2^0 - u_{h_2}^0\|_2 + \|u_1^1 - w_{h_1}^1\|_1 \quad (46)$$

Case 2 implies

$$\|u_1^1 - u_{h_1}^1\|_1 \leq \|u_1^1 - w_{h_1}^1\|_1 + \|u_2^0 - u_{h_2}^0\|_2 \quad (47)$$

$$\rho \|u_1^1 - u_{h_1}^1\|_1 \leq \|u_2^0 - u_{h_2}^0\|_2 \quad (48)$$

That is

$$\begin{aligned} \|u_1^1 - u_{h_1}^1\|_1 &\leq \left(\frac{1}{\rho}\right) \|u_2^0 - u_{h_2}^0\|_2 < \|u_2^0 - u_{h_2}^0\|_2 \\ &\leq \|u_1^1 - w_{h_1}^1\|_1 + \|u_2^0 - u_{h_2}^0\|_2 \end{aligned}$$

Which coincides with (47). So in both cases 1 and 2 we can write

$$\|u_1^1 - u_{h_1}^1\|_1 \leq \|u_1^1 - w_{h_1}^1\|_1 + \|u_2^0 - u_{h_2}^0\|_2 \quad (49)$$

which is equivalent to

$$\|u_1^1 - u_{h_1}^1\|_1 \leq \sum_{i=1}^1 \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^0 \|u_2^i - u_{h_2}^i\|_2 \quad (50)$$

Similarly, we obtain in domain 2

$$\begin{aligned} \|u_2^1 - u_{h_2}^1\|_2 &\leq \|u_2^1 - w_{h_2}^1\|_2 \\ &+ \max\{\rho \|u_2^1 - u_{h_2}^1\|_2; \|u_1^1 - u_{h_1}^1\|_1\} \end{aligned}$$

We then have to distinguish between two cases:

$$(1) : \max\{\rho \|u_2^1 - u_{h_2}^1\|_2; \|u_1^1 - u_{h_1}^1\|_1\} = \rho \|u_2^1 - u_{h_2}^1\|_2 \quad (51)$$

or

$$(2) : \max\{\rho \|u_2^1 - u_{h_2}^1\|_2; \|u_1^1 - u_{h_1}^1\|_1\} = \|u_1^1 - u_{h_1}^1\|_1 \quad (52)$$

Case 1 implies

$$\|u_2^1 - u_{h_2}^1\|_2 \leq \|u_2^1 - w_{h_2}^1\|_2 + \rho \|u_2^1 - u_{h_2}^1\|_2 \quad (53)$$

$$\|u_1^1 - u_{h_1}^1\|_1 \leq \rho \|u_2^1 - u_{h_2}^1\|_2 \quad (54)$$

By adding  $\|u_2^1 - w_{h_2}^1\|_2$  in (54) we get

$$\|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \leq \|u_2^1 - w_{h_2}^1\|_2 + \rho \|u_2^1 - u_{h_2}^1\|_2 \quad (55)$$

Hence  $\|u_2^1 - w_{h_2}^1\|_2 + \rho \|u_2^1 - u_{h_2}^1\|_2$  is bounded below by both  $\|u_2^1 - u_{h_2}^1\|_2$  and  $\|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1$  then

$$(a) : \|u_2^1 - u_{h_2}^1\|_2 \leq \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \quad (56)$$

or

$$(b) : \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \leq \|u_2^1 - u_{h_2}^1\|_2 \quad (57)$$

which implies

$$\begin{aligned} (a) : \|u_2^1 - u_{h_2}^1\|_2 &\leq \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \\ &\leq \|u_2^1 - w_{h_2}^1\|_2 + \rho \|u_2^1 - u_{h_2}^1\|_2 \end{aligned}$$

or

$$\begin{aligned} (b) : \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 &\leq \|u_2^1 - u_{h_2}^1\|_2 \\ &\leq \|u_2^1 - w_{h_2}^1\|_2 + \rho \|u_2^1 - u_{h_2}^1\|_2 \end{aligned}$$

Hence, (a) and (b) are true because they both coincide with (53). So, there is either a contradiction and thus case 1 is impossible or case 1 is possible only if

$$\|u_2^1 - u_{h_2}^1\|_2 = \|u_2^1 - w_{h_2}^1\|_2 + \rho \|u_2^1 - u_{h_2}^1\|_2 \quad (58)$$

Case 2 implies

$$\|u_2^1 - u_{h_2}^1\|_2 \leq \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1$$

$$\rho \|u_2^1 - u_{h_2}^1\|_2 \leq \|u_1^1 - u_{h_1}^1\|_1$$

So

$$\|u_2^1 - u_{h_2}^1\|_2 \leq \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \quad (59)$$

and

$$\begin{aligned} \|u_2^1 - u_{h_2}^1\|_2 &\leq \left(\frac{1}{\rho}\right) \|u_1^1 - u_{h_1}^1\|_1 < \|u_1^1 - u_{h_1}^1\|_1 \\ &\leq \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \end{aligned}$$

which coincides with (59). Then both cases 1 and 2 imply

$$\|u_2^1 - u_{h_2}^1\|_2 \leq \|u_2^1 - w_{h_2}^1\|_2 + \|u_1^1 - u_{h_1}^1\|_1 \quad (60)$$

Using (50) we get

$$\begin{aligned} \|u_2^1 - u_{h_2}^1\|_2 &\leq \|u_2^1 - w_{h_2}^1\|_2 \\ &+ \left(\sum_{i=1}^1 \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^0 \|u_2^i - u_{h_2}^i\|_2\right) \end{aligned}$$

or equivalently

$$\|u_2^1 - u_{h_2}^1\|_2 \leq \left(\sum_{i=0}^1 \|u_2^i - u_{h_2}^i\|_2 + \sum_{i=1}^1 \|u_1^i - w_{h_1}^i\|_1\right) \quad (61)$$

Now, let us assume that (62) and (63) are true

$$\|u_1^n - u_{h_1}^n\|_1 \leq \sum_{i=1}^n \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^{n-1} \|u_2^i - u_{h_2}^i\|_2 \quad (62)$$

$$\|u_2^n - u_{h_2}^n\|_2 \leq \sum_{i=0}^n \|u_2^i - u_{h_2}^i\|_2 + \sum_{i=1}^n \|u_1^i - w_{h_1}^i\|_1 \quad (63)$$

and prove that

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \sum_{i=1}^{n+1} \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^n \|u_2^i - u_{h_2}^i\|_2 \quad (64)$$

$$\|u_2^{n+1} - u_{h_2}^{n+1}\|_2 \leq \sum_{i=0}^{n+1} \|u_2^i - u_{h_2}^i\|_2 + \sum_{i=1}^{n+1} \|u_1^i - w_{h_1}^i\|_1 \quad (65)$$

Indeed, we have in domain 1

$$\begin{aligned} \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 &\leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \\ &\leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \\ &+ \max\left\{\left(\frac{1}{\beta}\right) \|F(u_1^{n+1}) - F(u_{h_1}^{n+1})\|_1; |u_2^n - u_{h_2}^n|_2\right\} \end{aligned}$$

then

$$\begin{aligned} \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 &\leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \\ &+ \max\left\{\left(\frac{1}{\beta}\right) \|F(u_1^{n+1}) - F(u_{h_1}^{n+1})\|_1; \|u_2^n - u_{h_2}^n\|_2\right\} \end{aligned}$$

thus

$$\begin{aligned} \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 &\leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \\ &+ \max\{\rho \|u_1^{n+1} - u_{h_1}^{n+1}\|_1; \|u_2^n - u_{h_2}^n\|_2\} \end{aligned}$$

We then have to distinguish between two cases:

$$(1) : \max\{\rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1; \|u_2^n - u_{h_2}^n\|_2\} = \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \quad (66)$$

or

$$(2) : \max\{\rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1; \|u_2^n - u_{h_2}^n\|_2\} = \|u_2^n - u_{h_2}^n\|_2 \quad (67)$$

Case 1 implies

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \quad (68)$$

$$\|u_2^n - u_{h_2}^n\|_2 \leq \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \quad (69)$$

By adding  $\|u_1^{n+1} - w_{h_1}^{n+1}\|_1$  in (73) we get

$$\|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1$$

So  $\|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1$  is bounded below by  $\|u_1^{n+1} - u_{h_1}^{n+1}\|_1$  and  $\|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1$ . This implies that

$$(a) : \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \quad (70)$$

or

$$(b) : \|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \quad (71)$$

Which implies

$$(a) : \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1$$

or

$$(b) : \|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1$$

Hence (a) and (b) are true because they both coincide with (68). So, there is either a contradiction and thus case 1 is impossible or case 1 is possible only if

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 = \|u_2^n - u_{h_2}^n\|_2 + \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 \quad (72)$$

Case 2 implies

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \|u_2^n - u_{h_2}^n\|_2 \quad (73)$$

$$\rho\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_2^n - u_{h_2}^n\|_2 \quad (74)$$

That is

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \left(\frac{1}{\rho}\right)\|u_2^n - u_{h_2}^n\|_2 < \|u_2^n - u_{h_2}^n\|_2 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \|u_2^n - u_{h_2}^n\|_2$$

Which coincides with (73). So in both cases 1 and 2 we can write

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \|u_2^n - u_{h_2}^n\|_2 \quad (75)$$

using (63) we get

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \|u_1^{n+1} - w_{h_1}^{n+1}\|_1 + \left(\sum_{i=0}^n \|u_2^i - u_{h_2}^i\|_2 + \sum_{i=1}^n \|u_1^i - w_{h_1}^i\|_1\right)$$

Hence

$$\|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \leq \sum_{i=1}^{n+1} \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^n \|u_2^i - u_{h_2}^i\|_2 \quad (76)$$

Which is the desired result (64). Estimate in domain 2 (65) can be proved similarly using estimate (64).

**Theorem 3.1** Let  $h = \max\{h_1, h_2\}$ . Then, for  $n$  large enough, there exists a constant  $C$  independent of both  $h$  and  $n$  such that

$$\|u_i - u_{h_i}^{n+1}\|_i \leq Ch^2 |\log h|^2; i = 1, 2. \quad (77)$$

**Proof** Let us give the proof for  $i = 1$ . The one for  $i = 2$  is similar and so will be omitted. Indeed, Let  $\kappa = \max\{k_1, k_2\}$ , then making use of Theorem 2.2, Lemma 3.1 and (34) respectively, we get

$$\|u_1 - u_{h_1}^{n+1}\|_1 \leq \|u_1 - u_1^{n+1}\|_1 + \|u_1^{n+1} - u_{h_1}^{n+1}\|_1 \quad (78)$$

So

$$\|u_1 - u_{h_1}^{n+1}\|_1 \leq \kappa^2 \|u_1 - u_1^0\|_1 + \left(\sum_{i=1}^{n+1} \|u_1^i - w_{h_1}^i\|_1 + \sum_{i=0}^n \|u_2^i - w_{h_2}^i\|_2\right)$$

Thus

$$\|u_1 - u_{h_1}^{n+1}\|_1 \leq \kappa^2 \|u_1 - u_1^0\|_1 + 2(n+1)Ch^2 |\log h| \quad (79)$$

So, for  $n$  large enough, we have

$$\kappa^{2n} = h^2 \quad (80)$$

then

$$n = \frac{|\log h|}{|\log \kappa|} = \bar{C} |\log h| \quad (81)$$

and thus

$$\|u_1 - u_{h_1}^{n+1}\|_1 \leq \kappa^2 \|u_1 - u_1^0\|_1 + 2(\bar{C} |\log h| + 1)Ch^2 |\log h| \quad (82)$$

So

$$\|u_1 - u_{h_1}^{n+1}\|_1 \leq Ch^2 |\log h|^2 \quad (83)$$

which is the desired result.

## REFERENCES

- [1] M. Boulbrachene, Ph. Cortey-Dumont, and J.-C. Miellou, "Mixing finite elements and finite differences in a subdomain method," in *Domain Decomposition for partial Differential Equations*, pp. 198–216, SIAM, Philadelphia, Pa, USA, 1988.
- [2] M. Boulbrachene and S. Saadi, "Maximum norm analysis of an overlapping nonmatching grids method for the obstacle problem" *Advances in Difference Equations*, pp. 1–10, 2006.
- [3] H. Brezis and M. Sibony, "Mthodes d'approximation et d'iteration pour les operateurs monotones" *Archive for Rational Mechanics and Analysis*, vol. 28, pp. 59–82, 1968.
- [4] X.-C. Cai, T.P. Mathew, and M. V. Sarkis, "Maximum norm analysis of overlapping nonmatching grid discretizations of elliptic equations," *SIAM Journal on Numerical Analysis*, vol. 37, no. 5, pp. 1709–1728, 2000.
- [5] P. G. Ciarlet and P.-A. Raviart, "Maximum principle and uniform convergence for the finite element method," *Computer Methods in Applied Mechanics and Engineering*, vol. 2, pp. 17–31, 1973.
- [6] A. Harbi and M. Boulbrachene, "Maximum Norm Analysis of a Non-matching Grids Method for Nonlinear Elliptic PDES", *Journal of Applied Mathematic*. Volume 2011.

- [7] J. Karatson and S. Korotov, "Discrete maximum principles for finite element solutions of nonlinear elliptic problems with mixed boundary conditions", *Numerische Mathematik*, vol. 99, no. 4, pp. 669–698, 2005.
- [8] P.-L. Lions, "On the Schwarz alternating method. I", in *Proceedings of the 1st International Symposium on Domain Decomposition Methods for Partial Differential Equations*, pp. 1–42, SIAM, Philadelphia, Pa, USA, 1988.
- [9] P.-L. Lions, "On the Schwarz alternating method. II. Stochastic interpretation and order properties", in *Proceedings of the 2nd International Symposium on Domain Decomposition Methods for Partial Differential Equations*, pp. 47–70, SIAM, Philadelphia, Pa, USA, 1989.
- [10] S.-H. Lui, "On monotone and Schwarz alternating methods for nonlinear elliptic PDEs", *Mathematical Modelling and Numerical Analysis*, vol. 35, no. 1, pp. 1–15, 2001.
- [11] S. H. Lui, "On linear monotone iteration and Schwarz methods for nonlinear elliptic PDEs", *Numerische Mathematik*, vol. 93, no. 1, pp. 109–129, 2002.
- [12] T. P. Mathew and G. Russo, "Maximum norm stability of difference schemes for parabolic equations on overset nonmatching space-time grids", *Mathematics of Computation*, vol. 72, no. 242, pp. 619–656, 2003.
- [13] J. Nitsche, " $L^\infty$ -convergence of finite element approximations", in *Proceedings of the Symposium on Mathematical Aspects of Finite Element Methods*, vol. 606 of *Lecture Notes in Mathematics*, pp. 261–274, 1977.