

# Extend three-wave method for the (3+1)-Dimensional Soliton Equation

Somayeh Arbabi Mohammad-Abadi and Maliheh Najafi

**Abstract**—In this paper, we study (3+1)-dimensional Soliton equation. We employ the Hirota's bilinear method to obtain the bilinear form of (3+1)-dimensional Soliton equation. Then by the idea of extended three-wave method, some exact soliton solutions including breather type solutions are presented.

**Keywords**—three-wave method; (3+1)-dimensional Soliton equation; Hirota's bilinear form.

## I. INTRODUCTION

The study of exact solutions of nonlinear partial differential equations plays an important role in soliton theory and explicit formulas of nonlinear partial differential equations play an essential role in the nonlinear science. Also, the explicit formulas may provide physical information and help us to understand the mechanism of related physical models. Recently, many kinds of powerful methods have been proposed to find exact solutions of nonlinear partial differential equations, e.g., the tanh-method [1], the homogeneous balance method [2], homotopy analysis method [3], [4], [5], [6], [7], [8], the  $F$ -expansion method [9], three-wave method [10], [11], [12], extended homoclinic test approach [13], [14], [15], the  $(\frac{G'}{G})$ -expansion method [16] and the exp-function method [17], [18], [19], [20], [21].

In this paper, by means of the Three-wave method, we will obtain some exact and new solutions for the (3+1)-dimensional soliton equation. In the following section we have a brief review on the Three-wave method and then we apply the method to find explicit formulas of solutions of the (3+1)-dimensional soliton equation in Section 3. The paper is concluded in Section 4.

## II. THREE-WAVE METHOD

Dai et al. [22], suggested the three-wave method for nonlinear evolution equations. The basic idea of this method applies the Painlevé analysis to make a transformation as

$$u = T(f) \quad (1)$$

for some new and unknown function  $f$ . Then we use this transformation in a high dimensional nonlinear equation of the general form

$$F(u, u_t, u_x, u_y, u_z, u_{xx}, u_{yy}, u_{zz}, \dots) = 0, \quad (2)$$

**S. Arbabi Mohammad-Abadi:** Department of Mathematics, Anar Branch, Islamic Azad University, Anar, Iran.  
 e-mail: (arbabi\_math87@yahoo.com).

**M. Najafi:** Department of Mathematics, Anar Branch, Islamic Azad University, Anar, Iran.

where  $u = u(x, y, z, t)$  and  $F$  is a polynomial of  $u$  and its derivatives. By substituting (1) in (2), the first one converts into the Hirota's bilinear form, which it will solve by taking a special form for  $f$  and assuming that the obtained Hirota's bilinear form has three-wave solutions, we can specify the unknown function  $f$ . For more details see [22], [23].

## III. (3+1)-DIMENSIONAL SOLITON EQUATION

In this paper, we investigate explicit formula of solutions of the following (3+1)-dimensional Soliton equation given in [24]

$$3u_{xz} - (2u_t + u_{xxx} - 2uu_x)_y + 2(u_x \partial_x^{-1} u_y)_x = 0, \quad (3)$$

or equivalently

$$3u_{xxz} - (2u_x t + u_{xxxx} - 2u_x u_{xx})_y + 2(u_{xx} u_y)_x = 0. \quad (4)$$

To solve eq. (3) we introduce a new dependent variable  $w$  by

$$w = -3(\ln f)_x \quad (5)$$

where  $f(x, y, z, t)$  is an unknown real function which will be determined. Substituting eq. (5) into eq. (4), we obtain the following Hirota's bilinear form

$$(3D_x D_z - D_y D_x^3 - 2D_y D_t) f \cdot f = 0, \quad (6)$$

where the D-operator is defined by

$$D_x^m D_y^k D_t^n f(x, y, t) \cdot g(x, y, t) = \left(\frac{\partial}{\partial x_1} - \frac{\partial}{\partial x_2}\right)^m \left(\frac{\partial}{\partial y_1} - \frac{\partial}{\partial y_2}\right)^k \left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2}\right)^n [f(x_1, y_1, t_1)g(x_2, y_2, t_2)], \quad (7)$$

and the right hand side is computed in

$$x_1 = x_2 = x, \quad y_1 = y_2 = y, \quad t_1 = t_2 = t.$$

Now we suppose the solution of eq. (6) as

$$f(x, y, z, t) = e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1} \quad (8)$$

where

$$\xi_1 = a_1 x + d_1 t \quad (9)$$

and

$$\xi_i = a_i x + b_i y + c_i z + d_i t, \quad i = 2, 3 \quad (10)$$

and  $a_1, d_1, \delta_1, a_i, b_i, c_i, d_i$  and  $\delta_i$  are some constants to be determined later. Substituting eq. (8) into eq. (6), and equating all coefficients of  $\exp(\xi_1)$ ,  $\exp(-\xi_1)$ ,  $\sin(\xi_2)$ ,  $\cos(\xi_2)$ ,  $\sinh(\xi_3)$  and  $\cosh(\xi_3)$  to zero, we get the following set of algebraic equations for  $a_1, d_1, \delta_1, a_i, b_i, c_i, d_i, \delta_i$ , ( $i = 2, 3$ )

$$\begin{aligned}
 2d_3b_3 + a_3^3b_3 + 3a_1^2a_3b_3 - 3a_3c_3 &= 0, \\
 -3a_3^2b_3a_1 + 3a_1c_3 - a_1^3b_3 - 2d_1b_3 &= 0, \\
 -3a_1^2a_2b_2 + a_2^3b_2 + 3a_2c_2 - 2d_2b_2 &= 0, \\
 a_1^3b_2 + 2d_1b_2 - 3a_2^2b_2a_1 - 3a_1c_2 &= 0, \\
 -3a_2^2a_3b_3 - 3a_3c_3 - 3a_3^2a_2b_2 + a_2^3b_2 & \\
 + a_3^3b_3 - 2d_2b_2 + 2d_3b_3 + 3a_2c_2 &= 0, \\
 -3a_2^2b_2a_3 + 2d_2b_3 + 2d_3b_2 + 3a_3^2b_3a_2 & \\
 -3a_2c_3 - 3a_3c_2 - a_2^3b_3 + a_3^3b_2 &= 0, \\
 -2\delta_1^2d_2b_2 + 4\delta_1^2a_2^3b_2 - 3\delta_2^2a_3c_3 & \\
 + 3\delta_1^2a_2c_2 + 4\delta_2^2a_3^3b_3 + 2\delta_2^2d_3b_3 &= 0.
 \end{aligned}$$

Solving the system of equations (11) with the aid of Maple, yields the following cases:

**A. Case I:**

$$\begin{aligned}
 a_2 = a_3 = d_2 = d_3 = 0, c_2 &= \frac{b_2c_3}{b_3}, \\
 d_1 &= -\frac{a_1(a_1^2b_3 - 3c_3)}{2b_3}
 \end{aligned}$$

for some arbitrary complex constants  $a_1, b_2, b_3, c_3$  and  $\delta_i, i = 1, 2, 3$ . Substitute eq. (12) into eq. (5) with eq. (8), we obtain the solution as

$$f(x, y, z, t) = e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1} \quad (13)$$

and

$$u(x, y, z, t) = -3 \frac{-a_1 e^{-\xi_1} + \delta_3 a_1 e^{\xi_1}}{e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1}} \quad (14)$$

for

$$\begin{aligned}
 \xi_1 &= a_1 x - \frac{a_1(a_1^2b_3 - 3c_3)t}{2b_3}, \\
 \xi_2 &= b_2 y + \frac{b_2c_3z}{b_3}, \quad \xi_3 = b_3 y + c_3 z
 \end{aligned}$$

If  $\delta_3 > 0$ , then we obtain the exact breather cross-kink solution

$$\begin{aligned}
 u(x, y, z, t) &= \\
 \frac{6a_1\sqrt{\delta_3} \sinh(\xi_1 - \theta)}{2\sqrt{\delta_3} \cosh(\xi_1 - \theta) + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3)}
 \end{aligned}$$

for

$$\theta = \frac{1}{2} \ln(\delta_3).$$

If  $\delta_3 < 0$ , then we obtain the exact breather cross-kink solution

$$\begin{aligned}
 u(x, y, z, t) &= \\
 \frac{6a_1\sqrt{-\delta_3} \cosh(\xi_1 - \theta)}{2\sqrt{-\delta_3} \sinh(\xi_1 - \theta) + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3)}
 \end{aligned}$$

for

$$\theta = \frac{1}{2} \ln(-\delta_3).$$

**B. Case II:**

$$\begin{aligned}
 a_2 = b_3 = c_3 = d_2 = 0, c_2 &= \frac{b_2(2d_1 + a_1^3)}{3a_1}, \\
 d_3 &= \frac{a_3(2d_1 + a_1^3 - a_3^2a_1)}{2a_1}
 \end{aligned}$$

for some arbitrary complex constants  $a_1, a_3, b_2, d_1, \delta_i, i = 1, 2, 3$ . Substitute eq. (16) into eq. (5) with eq. (8), we obtain the solution as follows

$$f(x, y, z, t) = e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1} \quad (17)$$

and

$$u(x, y, z, t) = \frac{-3(-a_1 e^{-\xi_1} + \delta_2 \sinh(\xi_3) a_3 + \delta_3 a_1 e^{\xi_1})}{e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1}} \quad (18)$$

for

$$\begin{aligned}
 \xi_1 &= a_1 x + d_1 t, \\
 \xi_2 &= b_2 y + \frac{b_2(2d_1 + a_1^3)z}{3a_1}, \\
 \xi_3 &= a_3 x + \frac{a_3(2d_1 + a_1^3 - a_3^2a_1)t}{2a_1},
 \end{aligned}$$

If  $\delta_3 > 0$ , then we obtain the exact breather cross-kink solution

$$\begin{aligned}
 u(x, y, z, t) &= \\
 \frac{6a_1\sqrt{\delta_3} \sinh(\xi_1 - \theta) - 3\delta_2 \sinh(\xi_3) a_3}{2\sqrt{\delta_3} \cosh(\xi_1 - \theta) + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3)}
 \end{aligned}$$

for

$$\theta = \frac{1}{2} \ln(\delta_3).$$

If  $\delta_3 < 0$ , then we obtain the exact breather cross-kink solution

$$\begin{aligned}
 u(x, y, z, t) &= \\
 \frac{6a_1\sqrt{-\delta_3} \cosh(\xi_1 - \theta) - 3\delta_2 \sinh(\xi_3) a_3}{2\sqrt{-\delta_3} \sinh(\xi_1 - \theta) + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3)}
 \end{aligned}$$

for

$$\theta = \frac{1}{2} \ln(-\delta_3).$$

C. CaseIII:

$$a_1 = d_1 = 0, c_3 = \frac{a_2^3 b_3 - 2 a_3 c_2 + a_3^2 b_3 a_2}{2 a_2},$$

$$b_2 = -\frac{a_2 b_3}{a_3}, d_3 = \frac{a_3 (3 a_2^3 b_3 - 6 a_3 c_2 + a_3^2 b_3 a_2)}{4 a_2 b_3},$$

$$d_2 = \frac{a_2^3 b_3 - 3 a_3 c_2}{2 b_3}, \delta_1 = \frac{\delta_2 a_3^2}{a_2^2}, \quad (20)$$

for some arbitrary complex constants  $a_2, d_2, \delta_1, \delta_2$ . Substitute eq. (20) into eq. (5) with eq. (8), we obtain the solution as follows

$$f(x, y, z, t) = 1 + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 \quad (21)$$

and

$$u(x, y, z, t) = \frac{3 \delta_1 \sin(\xi_2) a_2 - 3 \delta_2 \sinh(\xi_3) a_3}{1 + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3}, \quad (22)$$

for

$$\xi_2 = a_2 x - \frac{a_2 b_3 y}{a_3} + c_2 z + \frac{(a_2^3 b_3 - 3 a_3 c_2) t}{2 b_3},$$

$$\xi_3 = a_3 x + b_3 y + \frac{(a_2^3 b_3 - 2 a_3 c_2 + a_3^2 b_3 a_2) z}{2 a_2} + \frac{a_3 (3 a_2^3 b_3 - 6 a_3 c_2 + a_3^2 b_3 a_2) t}{4 a_2 b_3},$$

and

$$\delta_1 = \frac{\delta_2 a_3^2}{a_2^2}$$

D. CaseIV:

$$a_1 = -a_3, a_2 = i a_3, c_2 = \frac{b_2 c_3}{b_3}, d_1 = \frac{a_3 (4 a_3^2 b_3 - 3 c_3)}{2 b_3},$$

$$d_2 = -\frac{i (4 a_3^2 b_3 - 3 c_3) a_3}{2 b_3}, d_3 = -\frac{a_3 (4 a_3^2 b_3 - 3 c_3)}{2 b_3}, \quad (23)$$

for some arbitrary complex constants  $a_3, b_2, b_3, c_3, \delta_i, i = 1, 2, 3$ . Substitute eq. (23) into eq. (5) with eq. (8), we obtain the solution as follows

$$f(x, y, z, t) = e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1} \quad (24)$$

and

$$u(x, y, z, t) = -3 \frac{a_3 e^{-\xi_1} + i \delta_1 \sin(\xi_2) a_3 - \delta_2 \sinh(\xi_3) a_3 - \delta_3 a_3 e^{\xi_1}}{e^{-\xi_1} + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3) + \delta_3 e^{\xi_1}} \quad (25)$$

for

$$\xi_1 = -a_3 x + \frac{a_3 (4 a_3^2 b_3 - 3 c_3) t}{2 b_3}$$

$$\xi_2 = -i a_3 x - b_2 y - \frac{b_2 c_3 z}{b_3} + \frac{i (4 a_3^2 b_3 - 3 c_3) a_3 t}{2 b_3},$$

$$\xi_3 = -a_3 x - b_3 y - c_3 z + \frac{a_3 (4 a_3^2 b_3 - 3 c_3) t}{2 b_3}.$$

If  $\delta_3 > 0$ , then we obtain the exact breather cross-kink solution

$$u(x, y, z, t) = \frac{-6 a_3 \sqrt{\delta_3} \sinh(\xi_1 - \theta) - 3 i \delta_1 \sin(\xi_2) a_3 + 3 \delta_2 \sinh(\xi_3) a_3}{2 \sqrt{\delta_3} \cosh(\xi_1 - \theta) + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3)}$$

for

$$\theta = \frac{1}{2} \ln(\delta_3).$$

If  $\delta_3 < 0$ , then we obtain the exact breather cross-kink solution

$$u(x, y, z, t) = \frac{-6 a_3 \sqrt{-\delta_3} \cosh(\xi_1 - \theta) - 3 i \delta_1 \sin(\xi_2) a_3 + 3 \delta_2 \sinh(\xi_3) a_3}{2 \sqrt{-\delta_3} \sinh(\xi_1 - \theta) + \delta_1 \cos(\xi_2) + \delta_2 \cosh(\xi_3)}$$

for

$$\theta = \frac{1}{2} \ln(-\delta_3).$$

IV. CONCLUSION

In this paper, using the three-wave solution method we obtained some explicit formulas of solutions for the (3+1)-dimensional Soliton equation. Three-wave solution method with the aid of a symbolic computation software like Maple or Mathematica is an easy and straightforward method which can be apply to other nonlinear partial differential equations. It must be noted that, all obtained solutions have checked in the (3+1)-dimensional Soliton equation. All solutions satisfy in the equations.

REFERENCES

- [1] A.M. Wazwaz, *The tanh method: solitons and periodic solutions for the Dodd-Bullough-Tzikhailov and the Tzitzeica-Dodd-Bullough equations*, Chaos, Solitons and Fractals 25 (2005)55-63.
- [2] Z. Xiqiang, W. Limin, S. Weijun, *The repeated homogeneous balance method and its applications to nonlinear partial differential equations*, Chaos, Solitons and Fractals, 28(2) (2006) 448-453.
- [3] S.J. Liao, *Beyond Perturbation: Introduction to the homotopy analysis method*, Chapman & Hall/CRC Press, Boca Raton, 2003.
- [4] S.J. Liao, *On the homotopy analysis method for nonlinear problems*, Appl. Math. Comput., 147 (2006) 499-513.
- [5] S.J. Liao, *A general approach to get series solution of non-similarity boundary-layer flows*, Commun. Nonlinear Sci. Numer. Simul., 14(5)(2009) 2144-2159.
- [6] M.T. Darvishi, F. Khani, *A series solution of the foam drainage equation*, Comput. Math. Appl., 58 (2009) 360-368.
- [7] A. Aziz, F. Khani, M.T. Darvishi, *Homotopy analysis method for variable thermal conductivity heat flux gage with edge contact resistance*, Zeitschrift fuer Naturforschung A, 65a(10)(2010) 771-776.
- [8] F. Khani, M.T. Darvishi, R.S.R. Gorla, *Analytical investigation for cooling turbine disks with a non-Newtonian viscoelastic fluid*, Comput. Math. Appl., 61(7)(2011) 1728-1738.

- [9] E. Fan, Z. Jian, *Applications of the Jacobi elliptic function method to special-type nonlinear equations*, Phys. Lett. A, 305(6) (2002) 383-392.
- [10] M.T. Darvishi, Maliheh Najafi, Mohammad Najafi, *Exact three-wave solutions for high nonlinear form of Benjamin-Bona-Mahony-Burgers equations*, International Journal of Mathematical and Computer Sciences, 6(3)(2010) 127-131.
- [11] M.T. Darvishi, M. Najafi, *Some exact solutions of the (2+1)-dimensional breaking soliton equation using the three-wave method*, International Journal of Computational and Mathematical Sciences, 6(1)(2012)13-16.
- [12] M.T. Darvishi, Maliheh Najafi, Mohammad Najafi, *New exact solutions for the (3+1)-dimensional breaking soliton equation*, International Journal of Information and Mathematical Sciences, 6(2)(2010) 134-137.
- [13] M.T. Darvishi, Maliheh Najafi, Mohammad Najafi, *New application of EHTA for the generalized (2+1)-dimensional nonlinear evolution equations*, International Journal of Mathematical and Computer Sciences, 6(3)(2010) 132-138.
- [14] M.T. Darvishi, M. Najafi, *A modification of extended homoclinic test approach to solve the (3+1)-dimensional potential-YTSE equation*, Chin. Phys. Lett., 28(4)(2011) 040202.
- [15] M.T. Darvishi, M. Najafi, *Some complexiton type solutions of the (3+1)-dimensional Jimbo-Miwa equation*, International Journal of Computational and Mathematical Sciences, 6(1)(2012) 25-27.
- [16] M.T. Darvishi, Maliheh Najafi, Mohammad Najafi, *Traveling wave solutions for the (3+1)-dimensional breaking soliton equation by  $(\frac{G'}{G})$ -expansion method and modified F-expansion method*, International Journal of Computational and Mathematical Sciences, 6(2)(2012) 64-69.
- [17] J.H. He, M.A. Abdou, *New periodic solutions for nonlinear evolution equations using Exp-function method*, Chaos, Solitons and Fractals, 34 (2007) 1421-1429.
- [18] F. Khani, S. Hamed-Nezhad, M.T. Darvishi, S.-W. Ryu, *New solitary wave and periodic solutions of the foam drainage equation using the Exp-function method*, Nonlin. Anal.: Real World Appl., 10 (2009) 1904-1911.
- [19] B.-C. Shin, M.T. Darvishi, A. Barati, *Some exact and new solutions of the Nizhnik-Novikov-Vesselov equation using the Exp-function method*, Comput. Math. Appl., 58(11/12) (2009) 2147-2151.
- [20] F. Khani, M.T. Darvishi, A. Farmani, L. Kavitha, *New exact solutions of coupled (2+1)-dimensional nonlinear system of Schrödinger equations*, ANZIAM Journal, 52 (2010) 110-121.
- [21] X.H. Wu, J.H. He, *Exp-function method and its application to nonlinear equations*, Chaos, Solitons and Fractals, 38(3)(2008) 903-910.
- [22] Z.-D. Dai, S.-Q. Lin, D.-L. Li, G. Mu, *The three-wave method for nonlinear evolution equations*, Nonl. Sci. Lett. A 1(1) (2010)77-82.
- [23] C.-J. Wang, Z.-D. Dai, L. Liang, *Exact three-wave solution for higher dimensional KDV-type equation*, Appl. Math. Comput. 216 (2010.) 501-505.
- [24] J.-O. WU, X.-G. GENG, *Grammian Determinant Solution and Pfaffianization for a (3+1)-Dimensional Soliton Equation*, Commun. Theor. Phys. (Beijing, China) 52 (2009) 791-794.