

# Weak Convergence of Mann Iteration for a Hybrid Pair of Mappings in a Banach Space

Alemayehu Geremew Negash

**Abstract**—We prove the weak convergence of Mann iteration for a hybrid pair of maps to a common fixed point of a selfmap  $f$  and a multivalued  $f$  nonexpansive mapping  $T$  in Banach space  $E$ .

**Keywords**—Common fixed point, Mann iteration, Multivalued mapping, weak convergence.

## I. INTRODUCTION

LET  $E$  be a Banach space and  $K$ , a nonempty subset of  $E$ . We denote by  $2^E$ , the family of all subsets of  $E$ ;  $CB(E)$ , the family of nonempty closed and bounded subsets of  $E$  and  $C(E)$ , the family of nonempty compact subsets of  $E$ . Let  $f : K \rightarrow K$  be a selfmap. Let  $H$  be a Hausdorff metric on  $CB(E)$ . That is, for  $A, B \in CB(E)$ ,

$$H(A, B) = \max\{\sup_{x \in A} d(x, B), \sup_{x \in B} d(x, A)\},$$

where

$$d(x, B) = \inf\{\|x - y\| : y \in B\}.$$

A multivalued mapping  $T : K \rightarrow 2^K$  is called  $f$  nonexpansive if

$$H(Tx, Ty) \leq \|fx - fy\|,$$

for all  $x, y \in K$ .

If  $f = I_K$ , the identity mapping on  $K$ , then we call  $T$  is a multivalued nonexpansive mapping.

A point  $x$  is a fixed point of  $T$  if  $x \in Tx$ . A point  $x$  is called a common fixed point of  $f$  and  $T$  if  $fx = x \in Tx$ .  $F(T) = \{x \in K : x \in Tx\}$  stands for the fixed point set of a mapping  $T$  and  $F = F(T) \cap F(f) = \{x \in K : fx = x \in Tx\}$  stands for the common fixed point set of maps  $f$  and  $T$ .

Recently, Song and Wang [5] introduced the following Mann iterates of a Multivalued mapping  $T$ :

Let  $K$  be a nonempty convex subset of  $E$ ,  $\alpha_n \in [0, 1]$  and  $\gamma_n \in (0, \infty)$  such that  $\lim_{n \rightarrow \infty} \gamma_n = 0$ . Let  $T : K \rightarrow CB(K)$  be a multivalued mapping. Let  $x_0 \in K$ , and

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n y_n, \quad (1)$$

where  $y_n \in Tx_n$  such that

$$\|y_{n+1} - y_n\| \leq H(Tx_{n+1}, Tx_n) + \gamma_n, n = 0, 1, 2, \dots$$

Song and Wang [5] established the following theorem on the convergence of Mann iteration.

**Theorem 1.** [5] Let  $E$  be a Banach space satisfying Opial's condition and  $K$  be a nonempty, weakly compact and convex

Alemayehu Geremew Negash is with the Department of Mathematics, College of Natural Sciences, Jimma University, Jimma, P.O. Box 378, Ethiopia; (e-mail: alemg1972@gmail.com).

subset of  $E$ . Suppose that  $T : K \rightarrow CB(K)$  is a multivalued nonexpansive mappings for which  $F(T) \neq \emptyset$  and for which  $T(y) = \{y\}$  for each  $y \in F(T)$ . For  $x_0 \in K$ , let  $\{x_n\}$  be the Mann iteration defined by (1). Assume that

$$0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1.$$

Then the sequence  $\{x_n\}$  weakly converges to a fixed point of  $T$ .

The aim of this paper is to prove the weak convergence of Mann iteration for a hybrid pair of maps to a common fixed point of a selfmap  $f$  and a multivalued  $f$  nonexpansive mapping  $T$  in Banach space  $E$ . Our results extend the results of Song and Wang [5] to a hybrid pair of maps.

## II. PRELIMINARIES

Throughout this paper  $E$  denotes real Banach space. We denote the weak convergence of  $\{x_n\}$  to  $x$  in  $E$  by  $x_n \rightharpoonup x$ , and that of strong convergence by  $x_n \rightarrow x$ .

A Banach space  $E$  is said to satisfy Opial's condition [3] if for any sequence  $\{x_n\}$  in  $E$ ,  $x_n \rightharpoonup x$  ( $n \rightarrow \infty$ ) implies

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|,$$

for each  $y \in E$  with  $x \neq y$ .

Every Hilbert space and  $l^p$  ( $1 < p < \infty$ ) space satisfy Opial's condition [3].

A multivalued mapping  $T : K \rightarrow CB(K)$  is said to satisfy Condition I [4] if there is a nondecreasing function  $\varphi : [0, \infty) \rightarrow [0, \infty)$  with  $\varphi(0) = 0$ ,  $\varphi(r) > 0$  for  $r \in (0, \infty)$  such that

$$d(x, Tx) \geq \varphi(d(x, F(T))) \text{ for all } x \in K.$$

A selfmap  $f : E \rightarrow E$  is said to be weakly continuous if  $fx_n \rightharpoonup fx$  whenever  $x_n \rightharpoonup x$ . A map  $f : K \rightarrow E$  is said to be demiclosed at 0 if for each sequence  $\{x_n\}$  in  $K$  converging weakly to  $x$  and  $\{fx_n\}$  converging strongly to 0, we have  $fx = 0$ .

**Lemma 1.** [2] Let  $(E, d)$  be a complete metric space, and  $A, B \in CB(E)$  and  $a \in A$ . Then for each positive number  $\varepsilon$ , there exists  $b \in B$  such that

$$d(a, b) \leq H(A, B) + \varepsilon.$$

**Lemma 2.** [6] Let  $\{x_n\}$  and  $\{y_n\}$  be two bounded sequences in a Banach space  $E$  and  $\beta_n \in [0, 1]$  with

$$0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1.$$

Suppose  $x_{n+1} = \beta_n x_n + (1 - \beta_n)y_n$  for all integers  $n \geq 1$  and

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Then  $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$ .

We will construct the following iteration.

Let  $K$  be a nonempty subset of a metric space  $X$ . Let  $f : K \rightarrow K$ ,  $T : K \rightarrow CB(K)$  with  $f(K)$  is convex and  $Tx \subseteq f(K)$  for all  $x \in K$ . Let  $\alpha_n \in [0, 1]$ , and  $\gamma_n \in (0, \infty)$  such that  $\lim_{n \rightarrow \infty} \gamma_n = 0$ . Choose  $x_0 \in K$  and  $y_0 \in Tx_0$ . Let  $z_0 = fx_0$  and

$$\begin{aligned} z_1 &= fx_1 = (1 - \alpha_0)fx_0 + \alpha_0 y_0 \\ &= (1 - \alpha_0)z_0 + \alpha_0 y_0. \end{aligned}$$

From Lemma 1, there exists  $y_1 \in Tx_1$  such that

$$\|y_1 - y_0\| \leq H(Tx_1, Tx_0) + \gamma_0.$$

Let

$$z_2 = fx_2 = (1 - \alpha_1)z_1 + \alpha_1 y_1.$$

Inductively, we have

$$z_{n+1} = fx_{n+1} = (1 - \alpha_n)z_n + \alpha_n y_n, \quad (2)$$

where  $y_n \in Tx_n$  such that

$$\|y_{n+1} - y_n\| \leq H(Tx_{n+1}, Tx_n) + \gamma_n, \quad n = 0, 1, 2, \dots$$

### III. MAIN RESULTS

**Proposition 1.** [1] Let  $K$  be a nonempty subset of a Banach space  $E$ . Let  $f : K \rightarrow K$  be a selfmap with  $f(K)$  is convex. Suppose  $T : K \rightarrow CB(K)$  is a multivalued  $f$  nonexpansive mapping and  $Tx \subseteq f(K)$  for all  $x \in K$ . For  $x_0 \in K$ , let  $\{z_n\}$  be the Mann iteration associated with the maps  $T$  and  $f$ , defined by (2) and assume also that

$$0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1.$$

Then  $\lim_{n \rightarrow \infty} \|z_n - y_n\| = 0$  and  $\lim_{n \rightarrow \infty} d(z_n, Tx_n) = 0$ .

**Proof.** From the definition of the Mann iteration  $\{z_n\}$  given by (2), it follows that  $z_{n+1} = (1 - \alpha_n)z_n + \alpha_n y_n$ , where  $y_n \in Tx_n$  such that

$$\begin{aligned} \|y_{n+1} - y_n\| &\leq H(Tx_{n+1}, Tx_n) + \gamma_n \\ &\leq \|z_{n+1} - z_n\| + \gamma_n, \quad n = 0, 1, 2, \dots \end{aligned}$$

Therefore,

$$\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|z_{n+1} - z_n\|) \leq \limsup_{n \rightarrow \infty} \gamma_n = 0.$$

Hence, all conditions of Lemma 2 are satisfied. Hence, by Lemma 2, we obtain  $\lim_{n \rightarrow \infty} \|z_n - y_n\| = 0$ .

Since  $y_n \in Tx_n$  for all  $n = 0, 1, 2, \dots$ , we have  $d(z_n, Tx_n) \leq \|z_n - y_n\|$ .

Hence,  $\lim_{n \rightarrow \infty} d(z_n, Tx_n) = 0$ .

**Definition 1.** A multivalued mapping  $T : K \rightarrow CB(K)$  is said to satisfy *Condition I* associated with a selfmap  $f : K \rightarrow K$  if there is a nondecreasing function

$\varphi : [0, \infty) \rightarrow [0, \infty)$  with  $\varphi(0) = 0$ ,  $\varphi(r) > 0$  for  $r \in (0, \infty)$  such that

$$d(fx, Tx) \geq \varphi(d(fx, F)), \quad \text{for all } x \in K.$$

where  $F = \{x \in K : fx = x \in Tx\}$ .

If  $f = I_K$ , in Definition 1, then  $T$  is said to satisfy 'Condition I' (Senter and Dotson [4]).

**Theorem 2.** Let  $K$  be a nonempty closed subset of a Banach space  $E$ . Let  $f : K \rightarrow K$  be a continuous selfmap with  $f(K)$  is convex. Suppose  $T : K \rightarrow CB(K)$  is a multivalued  $f$  nonexpansive mapping for which  $Tx \subseteq f(K)$  for all  $x \in K$ ;  $F = F(T) \cap F(f) \neq \emptyset$ , and satisfies condition *I* associated with  $f$ . For  $x_0 \in K$ , let  $\{z_n\}$  be the Mann iteration associated with the maps  $T$  and  $f$ , defined by (2) and assume also that

$$0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1.$$

If  $T(y) = \{y\}$  for each  $y \in F(T)$ , then the Mann iteration  $\{z_n\}$  strongly converges to a common fixed point of  $f$  and  $T$ .

**Proof.** It follows from Proposition 1 that  $\lim_{n \rightarrow \infty} d(z_n, Tx_n) = 0$ .

Now let  $p \in F = F(T) \cap F(f)$ . Then,

$$\begin{aligned} \|z_{n+1} - p\| &\leq (1 - \alpha_n)\|z_n - p\| + \alpha_n\|y_n - p\| \\ &\leq (1 - \alpha_n)\|z_n - p\| + \alpha_n H(Tx_n, Tp) \\ &\leq (1 - \alpha_n)\|z_n - p\| + \alpha_n\|z_n - p\| \\ &= \|z_n - p\|, \quad n = 0, 1, 2, \dots \end{aligned}$$

which gives

$$d(z_{n+1}, F) \leq d(z_n, F), \quad n = 0, 1, 2, \dots$$

Then the sequence  $\{d(z_n, F)\}$  is a non increasing sequence of nonnegative reals and hence  $\lim_{n \rightarrow \infty} d(z_n, F)$  exists.

Since  $T$  satisfies Condition *I* associated with  $f$ , we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \varphi(d(z_n, F)) &= \lim_{n \rightarrow \infty} \varphi(d(fx_n, F)) \\ &\leq \lim_{n \rightarrow \infty} d(fx_n, fx_n) \\ &= \lim_{n \rightarrow \infty} d(z_n, fx_n) = 0. \end{aligned}$$

Hence,  $\lim_{n \rightarrow \infty} \varphi(d(z_n, F)) = 0$ .

Since  $\varphi$  is non decreasing function, we get  $\lim_{n \rightarrow \infty} d(z_n, F) = 0$ .

Hence, for  $\varepsilon > 0$ , there exist natural number  $n_0$  such that if  $n \geq n_0$ ,  $d(z_n, F) < \frac{\varepsilon}{4}$ . In particular,  $\inf\{\|z_{n_0} - y\| : y \in F\} < \frac{\varepsilon}{4}$ . So there must exist a  $z \in F$  such that  $\|z_{n_0} - z\| < \frac{\varepsilon}{2}$ .

Now for  $m, n \geq n_0$ , we have,

$$\begin{aligned} \|z_{n+m} - z_n\| &\leq \|z_{n+m} - z\| + \|z_n - z\| \\ &\leq 2\|z_n - z\| \\ &< 2\left(\frac{\varepsilon}{2}\right) = \varepsilon. \end{aligned}$$

Hence,  $\{z_n\}$  is a Cauchy sequence in a closed subset  $F$  of a Banach space  $E$ , therefore it converges to a point, say  $p \in K$ . Hence,  $d(p, F) = 0$ . Since  $F$  is closed,  $p \in F$ .

Hence, the conclusion of the theorem follows.

**Corollary 1.** [5] Let  $K$  be a nonempty, closed and convex subset of a Banach space  $E$ . Suppose that  $T : K \rightarrow CB(K)$  is a multi-valued nonexpansive mapping for which  $F(T) \neq \emptyset$  and for which  $T(y) = \{y\}$  for each  $y \in F(T)$  and satisfies condition I. For  $x_0 \in K$ , let  $\{x_n\}$  be the Mann iteration defined by (1). Assume that

$$0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1.$$

Then the sequence  $\{x_n\}$  strongly converges to a fixed point of  $T$ .

**Proof.** Follows from Theorem 2 by taking  $f = I_K$ .

**Theorem 3.** Let  $E$  be a Banach space satisfying Opial's condition and  $K$  be a nonempty weakly compact subset of  $E$ . Let  $f : K \rightarrow K$  be a weakly continuous selfmap with  $f(K)$  is convex. Suppose  $T : K \rightarrow CB(K)$  is a multivalued  $f$  nonexpansive mapping for which  $Tx \subseteq f(K)$  for all  $x \in K$ ;  $F(T) \cap F(f) \neq \emptyset$ , and  $d(x, Tx) \leq d(fx, Tx)$  for all  $x, y \in K$ . For  $x_0 \in K$ , let  $\{z_n\}$  be the Mann iteration associated with the maps  $T$  and  $f$ , defined by (2) and assume also that

$$0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1.$$

If  $T(y) = \{y\}$  for each  $y \in F(T)$  and  $I - f$  is demiclosed at 0, then the Mann iteration  $\{z_n\}$  weakly converges to a common fixed point of  $f$  and  $T$ .

**Proof.** It follows from Proposition 1 that  $\lim_{n \rightarrow \infty} d(z_n, Tx_n) = 0$ . Further, since  $d(x_n, Tx_n) \leq d(z_n, Tx_n)$  we get  $\lim_{n \rightarrow \infty} d(x_n, Tx_n) = 0$ . Now let  $p \in F(T) \cap F(f)$ . Then,

$$\begin{aligned} \|z_{n+1} - p\| &\leq (1 - \alpha_n)\|z_n - p\| + \alpha_n\|y_n - p\| \\ &\leq (1 - \alpha_n)\|z_n - p\| + \alpha_n H(Tx_n, Tp) \\ &\leq (1 - \alpha_n)\|z_n - p\| + \alpha_n\|z_n - p\| \\ &= \|z_n - p\|, \quad n = 0, 1, 2, \dots \end{aligned}$$

Then the sequence  $\{\|z_n - p\|\}$  is a decreasing sequence of nonnegative reals and hence  $\lim_{n \rightarrow \infty} \|z_n - p\|$  exists for each  $p \in F(T) \cap F(f)$ .

From the weak compactness of  $K$ , there exists a subsequence  $\{x_{n_j}\}$  of  $\{x_n\}$  such that  $x_{n_j} \rightharpoonup q$  as  $j \rightarrow \infty$ , for some  $q \in K$ . By the weak continuity of  $f$ , we have  $z_{n_j} \rightharpoonup fq = y$  (say) as  $j \rightarrow \infty$ . Suppose  $q \notin Tq$ . By the compactness of  $Tq$ , for any given  $x_{n_j}$ , there exists  $w_j \in Tq$  such that  $\|x_{n_j} - w_j\| = d(x_{n_j}, Tq)$ . Since  $Tq$  is compact,  $w_k$  has a convergent subsequence. For simplicity sake, we write that subsequence as  $w_j$  itself. So,  $w_j \rightarrow w \in Tq$ . Then  $q \neq w$ . Now the Opial's property of  $E$  implies that,

$$\begin{aligned} \limsup_{j \rightarrow \infty} \|z_{n_j} - w\| &\leq \limsup_{j \rightarrow \infty} [\|z_{n_j} - w_j\| + \|w_j - w\|] \\ &\leq \limsup_{j \rightarrow \infty} \|z_{n_j} - w_j\| \\ &\leq \limsup_{j \rightarrow \infty} [d(z_{n_j}, Tx_{n_j}) + H(Tx_{n_j}, Tq)] \end{aligned}$$

$$\begin{aligned} &= \limsup_{j \rightarrow \infty} \|z_{n_j} - y\| \\ &< \limsup_{j \rightarrow \infty} \|z_{n_j} - w\|, \end{aligned}$$

a contradiction. Hence,  $q = w$ , and so  $q \in Tq$ . Since,

$$\|fx_{n_j} - x_{n_j}\| \leq d(z_{n_j}, Tx_{n_j}) + d(Tx_{n_j}, x_{n_j}),$$

we have  $(I - f)(x_{n_j}) \rightarrow 0$  strongly as  $k \rightarrow \infty$ . By the demiclosedness of  $I - f$ , we have  $(I - f)(q) = 0$  and hence  $q = fq = y$ .

Hence,  $fy = y \in Ty$ .

Hence,  $y$  is a common fixed point of  $f$  and  $T$ .

Next we show that  $z_n \rightarrow y$  as  $n \rightarrow \infty$ . Suppose not. There exists another subsequence  $\{z_{n_k}\}$  of  $\{z_n\}$  such that  $z_{n_k} \rightharpoonup z \neq y$ . Then, by similar argument as above, we have  $fz = z \in Tz$ .

From Opial's property, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|z_n - z\| &= \limsup_{k \rightarrow \infty} \|z_{n_k} - z\| \\ &< \limsup_{k \rightarrow \infty} \|z_{n_k} - y\| \\ &= \limsup_{j \rightarrow \infty} \|z_{n_j} - y\| \\ &< \limsup_{j \rightarrow \infty} \|z_{n_j} - z\| \\ &= \lim_{n \rightarrow \infty} \|z_n - z\|, \end{aligned}$$

a contradiction. Hence,  $z_n \rightarrow y$  as  $n \rightarrow \infty$ .

Thus, the conclusion of the theorem follows.

**Corollary 2.** If  $f = I_K$ , then we get Theorem 1. Hence, Theorem 3 extends Theorem 1 to a pair of maps.

## REFERENCES

- [1] G. V. R. Babu, G.N. Alemayehu, Strong Convergence of Mann Iteration for a Hybrid Pair of Mappings in a Banach Space, *Mathematica Aeterna*, 3 (2013)(7), 501-507.
- [2] S. B. Nadler Jr., Multivalued contraction mappings, *Pacific J. Math.* 30 (1969) 475-487.
- [3] Z. Opial, Weak Convergence of the sequence of successive approximations for nonexpansive mappings, *Bull. Amer. Math. Soc.* 73 (1967) 591-597.
- [4] H. F. Senter, W. G. Dotson, Approximating fixed points of nonexpansive mappings, *Proc. Amer. Math. Soc.* 44 (1974) 375-380.
- [5] Y. Song, H. Wang, Convergence of iterative algorithms for multivalued mappings in Banach spaces, *Nonlinear Analysis* 70 (2009) 1547-1556.
- [6] T. Suzuki, Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces, *Fixed Point Theory Appl.* 2005 (1) (2005) 103-123, doi:10.1155 / FPTA.2005.103.