

Early Onset Neonatal Sepsis Pathogens in Malaysian Hospitals: Determining Empiric Antibiotic

Nazedah Ain Ibrahim, Mohamed Mansor Manan

Abstract—Information regarding early onset neonatal sepsis (EONS) pathogens may vary between regions. Global perspectives showed Group B *Streptococcal* (GBS) as the most common causative pathogens, but the widespread use of intrapartum antibiotics has changed the pathogens pattern towards gram negative microorganisms, especially *E. coli*. Objective of this study is to describe the pathogens isolated, to assess current treatment and risk of EONS. Records of 899 neonates born in three General Hospitals between 2009 until 2012 were retrospectively reviewed. Proven was found in 22 (3%) neonates. The majority was isolated with gram positive organisms, 17 (2.3%). All grams positive and most gram negative organisms showed sensitivity to the tested antibiotics. Only two rare gram negative organisms showed total resistant. Male was possible risk of proven EONS. Although proven EONS remains uncommon in Malaysia, nonetheless, the effect of intrapartum antibiotics still required continuous surveillance.

Keywords—Early onset neonatal sepsis, neonates, pathogens, gram positive, gram negative.

I. INTRODUCTION

NEONATAL sepsis is the major cause of hospitalization in neonatal intensive care unit (NICU). It is a significant cause of neonatal morbidity and mortality especially in the first 3 days of life which also defined as early onset neonatal sepsis (EONS) [1], [2].

There are a few antibiotics approaches being used worldwide in order to reduce the burden. First approach is intrapartum antibiotic prophylaxis for mother with risk factors [3]-[8] and second approach is empiric antibiotics for suspected EONS (neonates with high risk or clinical signs) [9]-[11].

EONS is normally caused by microorganisms that colonize the mother's genitourinary tract and occur within 72 hours of life [1]. The most common microorganisms isolated include group B *Streptococcus* (GBS), *Escherichia coli*, coagulase-negative *Staphylococcus* (CoNS), *Haemophilus influenzae* and *Listeria monocytogenes* [1].

Historically, Group B *Streptococcal* (GBS) has become the primary cause of EONS but the incidence has declined through the widespread use of intrapartum antibiotic prophylaxis [12], [13]. Most of the studies from year 2000 onwards have reported the shift of predominantly gram

positive microorganism causing EONS to predominantly gram negative especially *Escherichia coli* over time [12], [13].

However, information regarding EONS pathogens may vary between regions. It is crucial to know the common pathogens involved in EONS in our setting to facilitate the choice of antibiotic and improve the EONS management. We conducted multi-centre retrospective review over 899 neonates with suspected EONS to describe the common pathogens isolated in this specific population, to assess current treatment and risk of EONS.

II. METHODS

A. Sample Population and Study Design

Records of 899 neonates born in three General Hospitals (Hospital Raja Permaisuri Bainun Ipoh, Hospital Pulau Pinang and Hospital Sultanah Aminah Johor Bahru) in year 2009 until 2012 were retrospectively reviewed. Only neonates with blood cultures taken and documented prior empiric antibiotics administration and within 72 hours of life were included. Blood culture result was traced and documented in designated data collection form.

B. Study Approval

The study was approved by the Medical Research and Ethics Committee Ministry of Health Malaysia (NMRR-11-975-10283) and Research Ethics Committee Universiti Teknologi MARA (UiTM) (600-RMI (5/1/6/01)).

C. Statistical Analysis

All data were entered on a SPSS for windows version 16. The frequencies and percentages of each continuous variable studied was calculated and presented in the form of table. Categorical variables were assessed using Pearson Chi-Square test (χ^2 test) or Fisher's exact test. Logistic-regression models were used to assess associations between proven EONS and neonatal characteristics, maternal characteristics and risk. For all statistical analyses, the significance was set at 0.05.

III. RESULTS

Of 899 suspected EONS reviewed, there were 734 (82%) with documented blood culture taken prior empiric antibiotic within 72 hours of life. Out of it, 22 cases were proven EONS with positive blood culture. Demographics characteristics of suspected and proven EONS were shown in Table I. The male to female ratio was 2.14:1. Malay ethnicity shows the highest number of proven EONS as compared to other ethnics. Both suspected and proven EONS occurred more in premature and low birth weight neonates with mean of gestational age less

A.I. Nazedah is with the Pharmacy Department, Sungai Buloh Hospital, Jalan Hospital, 47000 Sungai Buloh, Selangor, Malaysia (phone: +6012-5858549; e-mail: naze_03@yahoo.com).

M.M. Mohamed is with the Faculty of Pharmacy, UiTM Puncak Alam Campus, Bandar Puncak Alam, 42300, Selangor, Malaysia (phone: +603-32584645; fax: +603-32584602; e-mail: mmmanan2002@yahoo.com).

than 37 weeks and birth weight less than 2.5 kg. Length of hospital stay was significantly longer in proven EONS group.

TABLE I
DEMOGRAPHICS CHARACTERISTICS

| Characteristics | Suspected EONS n = 712 | Proven EONS n = 22 | P value |
|---------------------------------------|---------------------------|-----------------------|---------------|
| Gender | | | 0.277 |
| Male, n (%) | 430 (60.40) | 15 (68.18) | |
| Female, n (%) | 282 (39.61) | 7 (31.82) | |
| Ethnic | | | 0.462 |
| Malay, n (%) | 449 (63.06) | 18 (81.82) | |
| Chinese, n (%) | 119 (16.71) | 1 (4.54) | |
| Indian, n (%) | 75 (10.53) | 1 (4.54) | |
| Others, n (%) | 69 (9.69) | 2 (9.09) | |
| Gestational age (week), mean (±SD) | 34.90 (3.975) | 33.19 (4.535) | 0.269 |
| Birth weight (kg), mean (±SD) | 2.27 (0.841) | 1.98 (0.848) | 0.113 |
| Length of stay (day), mean (±SD) | 17.89 (17.289) | 24.59 (19.271) | 0.002* |

* There was a significant association between proven EONS and prolongation of hospital stayed.

Table II shows the distribution of pathogens isolated from blood cultures. Gram positive pathogens contribute majority of the proven EONS case, 17 (77.27%) and the remaining were gram negative pathogens, 5 (22.73%). *Coagulase negative staphylococci* (CoNS), *Bacillus sp.* and *Streptococcus pneumoniae* were common isolated that account to more than 60% of total blood cultures isolated. Another 40% were isolated with other pathogens which contributed only one case each. Death was reported in one *Streptococcus pneumoniae* case due to invasive infection within 7 days of life.

TABLE II

DISTRIBUTION OF PATHOGENS AMONG 22 CASES OF PROVEN EONS OCCURRING IN 734 SUSPECTED EONS BORN IN THREE GOVERNMENT HOSPITALS BETWEEN 2009-2012

| Organism | n (%) |
|--|------------|
| Gram positive organisms | 17 (77.27) |
| Coagulase negative <i>staphylococci</i> (CoNS) | 7 (31.82) |
| <i>Bacillus sp.</i> | 5 (22.73) |
| <i>Streptococcus pneumoniae</i> * | 2 (9.09) |
| <i>Streptococcus spp.</i> | 1 (4.54) |
| <i>Enterococcus sp.</i> | 1 (4.54) |
| Group B <i>Streptococcal</i> (GBS) | 1 (4.54) |
| Gram negative organisms | 5 (22.73) |
| <i>Stenotrophomonas (xantho) maltophi</i> | 1 (4.54) |
| <i>Haemophilus influenza</i> | 1 (4.54) |
| <i>Spingomonas paucimobilis</i> | 1 (4.54) |
| <i>Enterobacter gergoviae</i> | 1 (4.54) |
| <i>E. coli</i> | 1 (4.54) |
| Total | 22 (100) |

* One case death within 7 days of life due to invasive infection.

Table III shows the antibiotic susceptibility pattern of isolated pathogens. Only recommended antibiotics for suspected EONS were reported. CoNS shows some cases were resistant to penicillin however, there were highly susceptible to gentamicin. Most of the *Bacillus sp.* was highly resistant to ampicillin and penicillin however, there were susceptible to

gentamicin and cefotaxime. In gram negative pathogens, *Stenotrophomonas (xantho) maltophi* shows resistant to gentamicin and *Enterobacter gergoviae* shows resistant to all tested antibiotics (ampicillin, gentamicin and cefotaxime).

TABLE III
ANTIBIOTIC SUSCEPTIBILITY PATTERN

| Total Case, n | Antibiotic tested (n) | | | | |
|--|-----------------------|---------------|---------------|----------------------|---------------|
| | Ampicillin | Penicillin | Gentamicin | Cefotaxime | |
| Gram positive | | | | | |
| Coagulase negative <i>staphylococci</i> (CoNS) | 7 | NT (7) | NT (5), R (2) | NT (1), S (6) | NT (7) |
| <i>Bacillus sp.</i> | 5 | R (4), NT (1) | R (3), NT (2) | I (1), S (3), NT (1) | NT (4), S (1) |
| <i>Streptococcus pneumoniae</i> | 2 | NT (2) | S (1), R (1) | NT (2) | NT (1), S (1) |
| <i>Streptococcus spp.</i> | 1 | S | S | S | NT |
| <i>Enterococcus sp.</i> | 1 | NT | S | NT | S |
| Group B <i>Streptococcal</i> (GBS) | 1 | NT | S | NT | S |
| Gram negative | | | | | |
| <i>Stenotrophomonas (xantho) maltophi</i> | 1 | NT | NT | R | NT |
| <i>Haemophilus influenza</i> | 1 | S | NT | NT | S |
| <i>Sphingomonas paucimobilis</i> | 1 | NT | NT | S | NT |
| <i>Enterobacter gergoviae</i> | 1 | R | NT | R | R |
| <i>E. coli</i> | 1 | S | NT | S | S |

NT = not tested, S = sensitive, R = resistant, I = intermediate.

Table IV shows the association between proven EONS and characteristics of neonates, mothers and the risk of EONS. There was no significant association found in this study. However, male, surfactant, caesarean delivery and prolonged rupture of membrane >18 hours were a possible risk of proven EONS.

IV. DISCUSSION

Blood culture is the most common investigations done prior empiric antibiotics administration in the diagnosis of serious infections [14]. The cultures will be incubated up to 5 days but, in modern culturing system the pathogens can be identified within 48 hours [15] and antibiotic use can be de-escalated within 3 days.

However, cultures with proven EONS remained scarce among neonates. The low number of positive cultures observed in the present study of 3% (n=22) was similarly reported by Clark et al. (2006) [10] and Metsvaht et al. (2010) [11]. This study also reported that male was more predominance to get proven EONS which also concordance with previous reports [16].

TABLE IV
CHARACTERISTICS AND THE RISK OF 22 NEONATES WITH PROVEN EONS

| Characteristic | Total, n | OR (95% CI) | P value |
|---|-------------|-------------------|---------|
| Neonate | | | |
| Gender | | | |
| Male* | 15 | | |
| Female | 7 | 1.34 (0.96-1.87) | 0.085 |
| Ethnic | | | |
| Malay* | 18 | | |
| Chinese | 1 | 0.21 (0.03-1.59) | 0.130 |
| Indian | 1 | 0.33 (0.04-2.53) | 0.287 |
| Others | 2 | 0.72 (0.16-3.19) | 0.668 |
| Premature (GA<37 weeks) | | | |
| No* | 6 | | |
| Yes | 16 | 1.19 (0.24-5.87) | 0.833 |
| Low birth weight (BW <2.5kg) | | | |
| No* | 6 | | |
| Yes | 16 | 1.86 (0.41-8.50) | 0.422 |
| Ventilator | | | |
| No* | 5 | | |
| Intubated | 11 | 0.98 (0.34-2.88) | 0.976 |
| CPAP | 6 | 1.11 (0.33-3.71) | 0.863 |
| Surfactant | | | |
| No* | 15 | | |
| Yes | 7 | 1.99 (0.79-4.99) | 0.145 |
| Maternal | | | |
| LSCS | | | |
| No* | 13 | | |
| Yes | 9 | 0.57 (0.24-1.37) | 0.208 |
| UTI / vaginal infection | | | |
| No* | 20 | | |
| Yes | 2 | 0.46 (0.05-4.51) | 0.504 |
| Maternal fever | | | |
| No* | 20 | | |
| Yes | 2 | 1.00 (0.15-6.63) | 0.997 |
| Chorioamnionitis | | | |
| No* | 20 | | |
| Yes | 2 | 1.23 (0.15-10.10) | 0.847 |
| Meconium stained amniotic fluid (MSAF) | | | |
| No* | 18 | | |
| Yes | 4 | 0.73 (0.09-6.24) | 0.774 |
| Prolonged rupture of membrane > 18 hour | | | |
| No* | 16 | | |
| Yes | 6 | 1.60 (0.62-4.17) | 0.334 |
| Perinatal asphyxia | | | |
| No* | 21 | | |
| Yes | 1 | 0.69 (0.09-5.54) | 0.724 |
| Intrapartum antibiotic | | | |
| Anytime during hospitalization | | | |
| No* | 17 | | |
| Yes | 5 | 1.24 (0.44-3.56) | |
| Prior delivery | | | |
| No* | 16 | | |
| Yes | 6 | 1.49 (0.55-4.04) | 0.432 |

*Neonates in this category served as the reference group.

As mentioned previously, microorganism isolated in EONS was closely related to maternal genital tract colonization [17]. There are many varieties of microorganisms causing EONS in developing countries. The most common microorganisms reported to be associated with EONS were group B streptococcus (GBS), *Escherichia coli* (*E. coli*), enterococcus and *Listeria monocytogenes*. Coagulase negative staphylococci (CoNS) normally can be found in both early and late onset neonatal sepsis [2]. Before the era of intrapartum antibiotic prophylaxis (IAP), GBS was the leading cause of mortality and morbidity among neonates [18].

In the three hospitals setting, antibiotics prophylaxis during pregnancy was commonly prescribed to mothers with risk factors for GBS transmission to neonates. For example, mothers with GBS colonization and preterm premature rupture of membrane will be covered for IAP and this may cause prolonged use of IAP [19].

In 1996, the American College of Obstetrician and Gynecologists (ACOG) and Centers for Disease Control and Prevention (CDC) recommended universal screening for all pregnancy women at 35-37 weeks of gestation before being administered with IAP to prevent GBS early onset sepsis [7]. Routine screening for GBS during pregnancy can prevent more cases of EONS compared to the risk based approach [18], [20]. The use of IAP caused marked reduction of GBS infection by 80-95% among neonates [18].

Many studies reported the use of IAP can affect the pattern of microorganism in EOS by increasing in the frequencies of non-GBS or antimicrobial-resistant [21]. The use of any IAP of more than 4 hours was associated with an increase in *E. coli* infection and ampicillin-resistant infection [22]-[24]. This phenomenon happens frequently in premature and LBW especially in very low birth weight (VLBW) neonates [12], [13], [21].

However in this study, only 22.73% of the proven infection cases were exposed to IAP and this could have affected the impact of IAP to these patients. The common antibiotics used in IAP for the present study were injection ampicillin, tablet erythromycin ethylsuccinate, injection cefoperazone, ampicillin/sulbactam and amoxicillin/clavulanate.

Our data findings suggested the risk of proven infection was higher in premature and LBW neonates (>70%) could be due to their immature immune system. More than 70% of microorganisms isolated in this study were gram positive organism. These findings are different from a previous report where the microorganism that caused EONS was predominantly gram negative [22]. It may be due to the lower number of IAP exposure in culture proven sepsis case as mentioned previously.

Overall analysis showed that the number of proven CoNS became the most common isolated gram positive organism. This organism was common organism isolated in birth canal. This finding was similar with the finding reported by Ghotaslou et al. 2007 in Iran [25]. Even though CoNS were commonly reported as blood culture contaminants, it has gained clinical importance as the etiology agents of catheter-associated bacteremia [14].

Minimum inhibitory concentration (MIC) is defined as the lowest concentration of antimicrobial required to inhibit the visible growth of a microorganism after overnight incubation [26]. It is the gold standard for determining the susceptibility of organisms to antimicrobial and being used by diagnostic laboratories to confirm unusual antimicrobial resistance [26].

Antimicrobial susceptibility surveillance plays an important role in determining empiric antibiotic therapy. Many studies show that by interpreting antimicrobial susceptibility data especially in specific infections [27]-[29] or patients [30]-[32] can provide the update on the most effective empirical treatment and subsequently can avoid the emergence of resistances.

In our study, penicillin (n=6) showed the highest reported resistance followed by ampicillin resistance (n=5), gentamicin resistance (n=2) and cefotaxime resistance last (n=1). Higher number of ampicillin resistance cases in this study was consistent with the findings reported by other authors [12], [22]-[24].

Bacillus sp. showed highly resistant to penicillin and ampicillin, however it was sensitive to either gentamicin or cefotaxime. *E. coli* proven infections occurred in one case and was sensitive to all antibiotic tested. This finding contradicts a report where increase in IAP exposure will increase resistant *E. coli* infection [12]. These findings showed the choices of antibiotics used for EONS in the present studied neonates are effective where penicillin or ampicillin plus gentamicin is the recommended regimen of choice for EONS and ampicillin plus cefotaxime was the alternative regimen [33], [34].

V. CONCLUSION

Proven EONS remains uncommon in Malaysia, only 3% reported in this study and majority isolated pathogens was gram positive organisms. The effect of intrapartum antibiotics still required continuous surveillance. However, this review show that current treatment used for managing suspected EONS was still effective.

ACKNOWLEDGMENT

The authors would like to thank the Director General of Health Malaysia for permission to publish this paper. The authors also acknowledge the help and corporation from the Hospital Director, staffs in NICU, and staff of the Medical Record Unit of Hospital Raja Permaisuri Bainun Ipoh, Hospital Pulau Pinang and Hospital Sultanah Aminah Johor Bahru, Malaysia.

REFERENCES

- [1] L. A-B Ann. et al., "Neonatal Sepsis," Retrieved September 3, 2011 from <http://emedicine.medscape.com/article/978352-overview>, 2011.
- [2] S. Vergnano, M. Sharland, P. Kazembe, and C. P. T. Mwansambo, "Health Neonatal sepsis: an international perspective," *Archives of Disease in Childhood. Fetal Neonatal, Edition*, 2005, 90, F220-F224.
- [3] W.E. Benitz, J.B. Gould, and M.L. Druzin, "Antimicrobial prevention of early-onset Group B Streptococcal Sepsis: Estimates of risk reduction based on a critical literature review" *Journal of the American Academy of Pediatrics*, 1999, 103, e78.

- [4] S. Kenyon, M. Boulvain and J. Neilson, "Antibiotics for preterm rupture of membranes (Review)," *Cochrane Database of Systematic Review*, 2007, 4, CD001058.
- [5] T. Siriwachirachai, U.S. Sangkomkamhang, P. Lumbiganon, and M. Laopaiboon, "Antibiotics for meconium-stained amniotic fluid in labour for preventing maternal and neonatal infections (Review)," *Cochrane Database System Review*, 2010, 12, CD007772.
- [6] C. H. Raynes-Greenow, C. L Roberts, J. C. Bell, B. Peat, G. L. Gilbert and S. Parker, "Antibiotics for ureaplasma in the vagina in pregnancy (Review)," *Cochrane Database of Systematic Review*, 2011, 9, CD003767.
- [7] J.R. Verani, L. McGee and S.J. Schrag, "Prevention of perinatal group B streptococcal disease-revised guidelines from CDC, 2010. Division of Bacterial Diseases, national Center for Immunization and Respiratory Diseases, Center for Disease Control and Prevention (CDC)," *MMWR Recomm Rep* 2010, 59 (RR-10):1-36.
- [8] ACOG, Committee opinion, "Prevention of early-onset group B streptococcal disease in newborns," *The American College of Obstetricians and Gynecologists*, 2011, 485.
- [9] E.I. Mtitimila and R.W. Cooke, "Antibiotic regimens for suspected early neonatal sepsis," *Cochrane Database of Systematic Review*, 2004, CD004495.
- [10] R.H. Clark, B.T. Bloom, A.R. Spitzer and D.R. Gerstmann, "Empiric use of ampicillin and cefotaxime, compared with ampicillin and gentamicin, for neonates at risk for sepsis is associated with an increased risk of neonatal death," *Journal of the American Academy of Pediatrics*, 2006, 117, 67-74.
- [11] T. Metsvaht, M-L. Ilmoja, Ü. Parm, L. Maipuu, M. Merila and I. Lutsar, "Comparison of ampicillin plus gentamicin vs. penicillin plus gentamicin in empiric treatment of neonates at risk of early onset sepsis," *Acta Paediatrica*, 2010, 99, 665-672.
- [12] J. S. Schrag, J. L. Hadler, K.E. Arnold, P. Martell-Cleary, A. Reingold, and A. Schuchat, "Risk factors for invasive, early onset *Escherichia coli* infections in the era of widespread intrapartum antibiotic use," *Journal of the American Academy of Pediatrics*, 2006, 118, 570-576.
- [13] M.J. Bizzarro, L-M. Dembry, R.S. Baltimore and P.G. Gallagher, "Changing patterns in neonatal *Escherichia coli* sepsis and the ampicillin resistance in the era of intrapartum antibiotic prophylaxis," *Journal of the American Academy of Pediatrics*, 2008, 121, 689-696.
- [14] M.P. Weinstein, "Blood culture contamination: Persisting problems and partial progress," *Journal of Clinical Microbiology*, 2003, 41, 6, 2275-2278.
- [15] J.S. Gerdes, "Diagnosis and management of bacterial infections in the neonate," *Journal of Pediatric Clinical of North America*, 2004, 51, 939-959.
- [16] R.Y. Bhat, and L.P. Baby, "Early onset of neonatal sepsis: analysis the risk factors and the bacterial isolated by using the BacT alert system," *Journal of Clinical and Diagnostic Research*, 2011, 5, 7, 1385-1388.
- [17] WHO, 17th Expert Committee on the Selection and Use of Essential Medicines, "Review of the efficacy and safety of procaine benzylpenicillin in neonates," 2009.
- [18] B.J. Stoll, N.I. Hansen, R.D. Higgins, A.A. Fanaroff, S. Duara, R. Goldberg, A. Laptook, M. Walsh, W. Oh and E. Hale, "Very low birth weight preterm infants with early onset neonatal sepsis: The predominance of gram negative infections continues in the National Institute of Child Health and Human Development Neonatal Research Network, 2002-2003," *The Pediatric Infection Disease Journal*, 2005, 24, 635-639.
- [19] N.E. Glass, J. Sculkin, S. Chamany, L.E. Riley, A. Schuchat and S. Schrag, "Opportunities to reduce overuse of antibiotics for perinatal group B streptococcal disease prevention and management of preterm premature rupture of membranes," *Infectious Diseases in Obstetrics and Gynecology*, 2005, 13, 1, 5-10.
- [20] S. Schrag, R. Gorwitz, K. Fultz-Butts, and A. Scuchat, "Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC," *MMWR Recommendations and Reports*, 2002, 51(RR-11):1-22.
- [21] M.R. Moore, S.J. Schrag and A. Schuchat, "Effects of intrapartum antimicrobial prophylaxis for prevention of group B streptococcal disease on the incidence and ecology of early onset neonatal sepsis," *The Lancet Infectious Diseases*, 2003, 3, 201-213.
- [22] B.J. Stoll, N. Hansen, A. Fanaoff, L.L. Wright, W.A. Carlo, R.A. Ehrenkranz, J.A. Lemons, E.F. Donovan, A.R. Stark, J.E. Tayson, W. Oh, C. R. Bauer, S.B. Korones, S. Shankaran, A.R. Laptook, D.K. Stevenson, L-A. Papile and K. Poole, "Changes in pathogens causing

- Early-Onset Sepsis in very-low-birth-weight infants,” *New England Journal of Medicine*, 2002, 347, 240-247.
- [23] P. Kuhn, C. Dheu, B. Colender, D. Chognot, L. Keller, H. Demil, L. Donato, B. Langer, J. Messer and D. Astruc, “Incidence and distribution of pathogens in early onset neonatal sepsis in the era of antenatal antibiotics,” *Paediatric and Perinatal Epidemiology*, 2010, 24,479-487.
- [24] C.H. Tsai, Y.Y. Chen, K.G. Wang, C.Y. Chen and C.P. Chen, “Characteristics of early onset sepsis caused by *Escherichia coli*,” *Taiwanese Journal of Obstetrics & Gynecology*, 2011 51,26-30.
- [25] R. Ghotaslou, Z. Ghorashi and M-R. Nahaei, “Klebsiella pneumonia in neonatal sepsis: A 3 year study in the Pediatric Hospital of Tabriz, Iran, 2007, 60,126-128.
- [26] J.M. Andrews, “Determination of minimum inhibitory concentrations,” *Journal of Antimicrobial Chemotherapy*, 2001, 48,S1,5-16.
- [27] C.M. Muvunyi, F. Masaisa, C. Bayingana, L. Mutesa, A. Musemakweri, G. Muhirwa and G. (W.) Claeys, “Decreased susceptibility to commonly used antimicrobial agents in bacterial pathogens isolated from urinary tract infections in Rwanda: Need for new antimicrobial guidelines,” *American Journal of Tropical Medicine and Hygiene*, 2011, 84,6,923-928.
- [28] H. Gazi, S. Kurutepe, S. Surucuoglu, A. Teker, and B. Ozbakkaloglu, “Antimicrobial susceptibility of bacterial pathogens in the oropharynx of healthy school children in Turkey,” *Indian Journal of Medical Research*, 2004, 120,489-494.
- [29] A. Nerurkar, P. Solanky and S.S. Naik, “Bacterial pathogens in urinary tract infection and antibiotic susceptibility pattern,” *Journal of Pharmaceutical and Biomedical Sciences*, 2012, 21,21,1-3.
- [30] M. Radji, Siti Fauziah and N. Aribinuko, “Antibiotic sensitivity pattern of bacterial pathogens in the intensive care unit of Fatmawati Hospital, Indonesia,” *Asian Pacific Journal of Tropical Biomedicine*, 2011, 1,1,39-42.
- [31] I. Shaifali, U. Gupta, S.E. Mahmood and J. Ahmed, “Antibiotic susceptibility patterns of urinary pathogens in female outpatients,” *North American Journal of Medical Sciences*, 2012, 4,4,163-169.
- [32] E. Saez-Lopez, E. Guiral and S.M. Soto, “Neonatal sepsis by bacteria: A big problem for children,” *Clinical Microbiology*, 2013, 2,6,1000125.
- [33] *National Antibiotic Guideline*. Malaysia: Ministry of Health, 2008. Page 181.
- [34] A.I. Nazedah, M.M. Mohamed, and A.A. Noorizan, “The Effectiveness of empiric antibiotic therapy in the prevention of early onset sepsis,” *International Journal of Current Research*, 2014, 4,6258-6265.