

# Spline Collocation for Solving System of Fredholm and Volterra Integral Equations

N. Ebrahimi, J. Rashidinia

**Abstract**—In this paper, numerical solution of system of Fredholm and Volterra integral equations by means of the Spline collocation method is considered. This approximation reduces the system of integral equations to an explicit system of algebraic equations. The solution is collocated by cubic B-spline and the integrand is approximated by the Newton-Cotes formula. The error analysis of proposed numerical method is studied theoretically. The results are compared with the results obtained by other methods to illustrate the accuracy and the implementation of our method.

**Keywords**—Convergence analysis, Cubic B-spline, Newton-Cotes formula, System of Fredholm and Volterra integral equations.

## I. INTRODUCTION

CONSIDER the system of integral equations of the form:

$$F(t) = G(t) + \int_a^b K(t, x, F(x))dx, \quad t \in [a, b] \quad (1)$$

where

$$\begin{aligned} F(t) &= [f_1(t), f_2(t), \dots, f_n(t)]^T, \\ G(t) &= [g_1(t), g_2(t), \dots, g_n(t)]^T, \\ K(t, x, F(x)) &= \sum_{j=1}^n k_{ij}(t, x, f_j(x)), \quad i = 1, \dots, n. \end{aligned}$$

In (1) the known kernel  $K$  is continuous, the function  $G(x)$  is given, existence and uniqueness of the solution of (1) is given by [22] and  $F(x)$  is the solution to be determined [1], [2]. We consider the  $i$ -th equation of (1):

$$f_i(t) = g_i(t) + \sum_{j=1}^n \int_a^b k_{ij}(t, x, f_j(x)) dx, \quad t \in [a, b], \quad i = 1, \dots, n \quad (2)$$

There has been considerable interest in solving integral equation (1). E. Babolian et al. [13] applied an Adomian decomposition method for solving system of linear Fredholm integral equations of the second kind. Numerical solution of the system of linear Fredholm integral equations (1), has been proposed by Maleknejad et al. in [14], [15]. Rationalized Haar functions have been used for direct numerical solutions in [14] and also Block-Pulse functions to propose solutions for the system of Fredholm integral equations have been developed in [15]. This type of equations have been solved in many papers with different methods such as Taylor's expansion [16], [17], operational matrices method [18] homotopy perturbation method [19]-[22], Sinc collocation method [23], [24] and

Adomian's decomposition method [25]-[28]. Using a global approximation to the solution of Fredholm and Volterra integral equation of the second kind is constructed by means of the cubic spline quadrature in [9]-[12].

In this paper, we use cubic B-spline collocation for approximation unknown function and use of the Newton-Cotes rules for approximating integrand.

## II. CUBIC B-SPLINE

We introduce the cubic B-spline space and basis functions to construct an interpolant  $S_N = [s_1, s_2, \dots, s_n]^T$ , to be used in the formulation of the cubic B-spline collocation method.

Let  $\pi: \{a = t_0 < t_1 < \dots < t_N = b\}$ , be a uniform partition of the interval  $[a, b]$  with step size  $h = \frac{b-a}{N}$ . The cubic spline space is denoted by

$$S_3(\pi) = \{s \in C^2[a, b]; s_i|_{[t_k, t_{k+1}]} \in P_3, i = 1, \dots, n, k = 0, \dots, N\},$$

where  $P_3$  is the class of cubic polynomials. The construction of the cubic B-spline interpolate  $S_N$  to the analytical solution  $F(t) = [f_1(t), f_2(t), \dots, f_n(t)]^T$ , for (1) can be performed with the help of the four additional knots such that

$$t_{-2} < t_{-1} < t_0 \text{ and } t_N < t_{N+1} < t_{N+2}.$$

Following [9] we can define a cubic B-spline  $s_i(t)$  of the form

$$s_i(t) = \sum_{k=-1}^{N+1} c_{ki} B_k^3(t) \quad (3)$$

where

$$B_k^3(t) = \frac{1}{6h^3} \begin{cases} (t - t_{k-2})^3, & t \in [t_{k-2}, t_{k-1}] \\ h^3 + 3h^2(t - t_{k-1}) + 3h(t - t_{k-1})^2 - 3(t - t_{k-1})^3, & t \in [t_{k-1}, t_k] \\ h^3 + 3h^2(t_{k+1} - t) + 3h(t_{k+1} - t)^2 - 3(t_{k+1} - t)^3, & t \in [t_k, t_{k+1}] \\ (t_{k+2} - t)^3, & t \in [t_{k+1}, t_{k+2}] \\ 0, & \text{otherwise,} \end{cases}$$

satisfying the following interpolator conditions:

$$s_i(t_k) = f_i(t_k), \quad 1 \leq i \leq n, \quad 0 \leq k \leq N,$$

and the end conditions

$$\begin{aligned} \text{(i)} \quad s_i'(t_0) &= f_i'(t_0), \quad s_i'(t_N) = f_i'(t_N), \quad i = 1, \dots, n, \text{ or} \\ \text{(ii)} \quad D^j s_i(t_0) &= D^j s_i(t_N), \quad j = 1, 2, \text{ or} \\ \text{(iii)} \quad s_i''(t_0) &= 0, \quad s_i''(t_N) = 0. \end{aligned} \quad (4)$$

N. Ebrahimi is with the School of Mathematics, Islamic Azad University, Central Tehran Branch, Iran (e-mail: ebrahimi\_nehzat@yahoo.com).

J. Rashidinia is with the School of Mathematics, Iran University of Science & Technology, Narmak, Tehran 16844, Iran (e-mail: rashidinia@iust.ac.ir).

### III. THE COLLOCATION METHOD

#### A. Nonlinear Fredholm Integral Equation

In the given nonlinear Fredholm integral equation (2), we can approximate the unknown function by cubic B-spline (3),

$$s_i(t) = g_i(t) + \sum_{j=1}^n \int_a^b k_{ij}(t, x, s_j(x)) dx, t \in [a, b], i = 1, \dots, n. \quad (5)$$

We now collocate (5) at collocation points  $t_k = a + kh, h = \frac{b-a}{N}, k = 0, 1, \dots, N$ , and we obtain

$$s_i(t_k) = g_i(t_k) + \sum_{j=1}^n \int_a^b k_{ij}(t_k, x, s_j(x)) dx, i = 1, \dots, n, k = 0, \dots, N. \quad (6)$$

To approximate the integral equation (6), we can use the Newton-Cotes formula [2], when  $n$  is even then the Simpson rule can be used and when  $n$  is multiple of 3, we have to use the three-eighth rule, then we get the following nonlinear system:

$$s_i(t_k) = g_i(t_k) + h \sum_{j=1}^n \sum_{r=0}^N w_{k,r} k_{ij}(t_k, x_r, s_j(x_r)), i = 1, \dots, n, k = 0, \dots, N, \quad (7)$$

where  $x_r = a + rh, r = 0, \dots, N$ , we need more equations to obtain the unique solution for (7). Hence by associating (7) with (4) for the  $i$ -th equation, we have the following nonlinear system  $(N+3) \times (N+3)$ :

$$\begin{cases} s_i(t_k) = g_i(t_k) + h \sum_{j=1}^n \sum_{r=0}^N w_{k,r} k_{ij}(t_k, x_r, s_j(x_r)), k = 0, \dots, N, \\ D^j s_i(t_0) = D^j s_i(t_N), \quad j = 1, 2, \end{cases} \quad (8)$$

where  $w_{k,r}$  represents the weights for a quadrature rule of Newton-Cotes type. By solving the above nonlinear system for each  $i$ , we can determine the coefficients in (3), by setting coefficients in (3), we obtain the approximate solution for (1).

#### B. Nonlinear Volterra Integral Equation

Now we consider the system of nonlinear Volterra integral equations

$$F(t) = G(t) + \int_a^t K(t, x, F(x)) dx, t \in [a, b] \quad (9)$$

the solutions of (9) can be replaced with cubic B-spline and so we collocate (9) at collocation points  $t_k = a + kh, h = \frac{t-a}{N}, k = 0, 1, \dots, N$ , and we obtain

$$\begin{cases} s_i(t_k) = g_i(t_k) + \sum_{j=1}^n \int_a^{t_k} k_{ij}(t_k, x, s_j(x)) dx, i = 1, \dots, n, k = 1, \dots, N, \\ s_i(t_0) = g_i(t_0). \end{cases} \quad (10)$$

To approximate the integral equation (10), we can use the Newton-Cotes formula, when  $n$  is even then the Simpson rule can be used and when  $n$  is multiple of 3, we have to use the three-eighth rule, then we get the following nonlinear system:

$$\begin{cases} s_i(t_k) = g_i(t_k) + h \sum_{j=1}^n \sum_{r=0}^k w_{k,r} k_{ij}(t_k, x_r, s_j(x_r)), k = 1, \dots, N, \\ s_i(t_0) = g_i(t_0). \end{cases} \quad (11)$$

We need more equations to obtain the unique solution for (11). Hence by associating (11) with (4) for the  $i$ -th equation, we have the following nonlinear system  $(N+3) \times (N+3)$ :

$$\begin{cases} s_i(t_k) = g_i(t_k) + h \sum_{j=1}^n \sum_{r=0}^k w_{k,r} k_{ij}(t_k, x_r, s_j(x_r)), k = 1, \dots, N, \\ D^j s_i(t_0) = D^j s_i(t_N), \quad j = 1, 2, \\ s_i(t_0) = g_i(t_0). \end{cases} \quad (12)$$

By solving the above nonlinear system for each  $i$ , we can determine the coefficients in (3), by setting coefficients in (3), we obtain the approximate solution for (9).

### IV. ANALYSIS: CONVERGENCE OF THE APPROXIMATE SOLUTION

In this section, we consider the error analysis of the system of nonlinear Volterra integral equations of the second kind.

We assume that  $S_N = [s_1, s_2, \dots, s_n]^T$ , is approximated solution for  $F(t) = [f_1(t), f_2(t), \dots, f_n(t)]^T$ . We need to recall the following definition in [9].

Definition: The most immediate error analysis for spline approximates  $s$  to a given function  $f$  defined on an interval  $[a, b]$  follows from the second integral relations.

If  $f \in C^4[a, b]$ , then  $\|D^j(f - S)\| \leq \gamma h^{4-j}, j = 0, 1, 2, 3, 4$ , where  $\|f\|_\infty = \max_{0 \leq i \leq N} \sup_{t_{i-1} \leq t \leq t_i} |f(t)|$ ,

and  $D^j$  the  $j$ -th derivative.

Theorem: The approximate method

$$S_N(t_k) = G(t_k) + h \sum_{r=0}^k w_{k,r} K(t_k, x_r, S_N(x_r)), k = 1, \dots, N, \quad (13)$$

for solution of the system of nonlinear Volterra integral (9) is converge and the error bounded is

$$|E_{Nk}| \leq \frac{hWL}{1-hWL} \sum_{r=0}^{k-1} |E_{Nr}| + \frac{1}{1-hWL} |E(h, t_k)|.$$

Proof: We know that at  $t_k = a + kh, h = \frac{t-a}{N}, k = 0, 1, \dots, N$ , the corresponding approximation method for the system of nonlinear Volterra integral equation (9) is

$$S_N(t_k) = G(t_k) + h \sum_{r=0}^k w_{k,r} K(t_k, x_r, S_N(x_r)), k = 1, \dots, N. \quad (14)$$

By discretizing (9) and approximating the integrand by the Newton-Cotes formula, we obtain

$$F(t_k) = G(t_k) + h \sum_{r=0}^k w_{k,r} K(t_k, x_r, F(x_r)) + E(h, t_k), k = 1, \dots, N, \quad (15)$$

where

$$E(h, t_k) = \int_a^{t_k} K(t_k, x, F(x)) dx - h \sum_{r=0}^k w_{k,r} K(t_k, x_r, F(x_r)).$$

By subtracting (15) from (14) and using interpolatory conditions of cubic B-spline, we get

$$S_N(t_k) - F(t_k) = h \sum_{r=0}^k w_{k,r} [K(t_k, x_r, S_N(x_r)) - K(t_k, x_r, F(x_r))] - E(h, t_k).$$

We suppose that  $W = \max_{r,k} |w_{k,r}|$  and  $S_N(t_k) = S_{Nk}$ ,  $F(t_k) = F_k$ ,  $k = 1, \dots, N$ , and kernel  $K$  satisfies a Lipschitz condition in its third argument of the form

$$|K(t, x, S_N) - K(t, x, F)| \leq L|S_N - F|,$$

where  $L$  is independent of  $t, x, S_N$  and  $F$ . We get

$$|S_{Nk} - F_k| \leq hWL \sum_{r=0}^k (S_{Nr} - F_r) + |E(h, t_k)|.$$

Then, for sufficiently small  $h$  (see [2]), we have

$$|E_{Nk}| \leq \frac{hWL}{1 - hWL} \sum_{r=0}^{k-1} |E_{Nr}| + \frac{1}{1 - hWL} |E(h, t_k)|,$$

where  $E_{Nk} = S_{Nk} - F_k$ ,  $k = 1, \dots, N$ .

Since by assumption both the quadrature error and the function approximate error are zero in the limit, it follows when  $h \rightarrow 0$ ,  $\lim \max |E(h, t_k)| = 0$ , then the above term is zero. We get for a fixed  $k$ ,  $|E_{Nk}| \rightarrow 0$  as  $h \rightarrow 0$ .

#### V. NUMERICAL EXAMPLES

In order to test the applicability of the presented method, we consider four examples of the system of linear and nonlinear Volterra and Fredholm integral equations. The absolute errors in the solution for various values of  $N$  are tabulated in tables. Our numerical solutions are compared with [3]-[8], [19].

Example 1. Consider the system of following linear Fredholm integral equations with exact solution

$$\begin{aligned} (f_1(t), f_2(t))^T &= (e^t, e^{-t})^T, \\ f_1(t) &= 2e^t + \frac{e^{t+1} - 1}{t + 1} - \int_0^1 e^{t-x} f_1(x) dx - \int_0^1 e^{(t+2)x} f_2(x) dx, \\ f_2(t) &= e^t + e^{-t} + \frac{e^{t+1} - 1}{t + 1} - \int_0^1 e^{tx} f_1(x) dx - \int_0^1 e^{t+x} f_2(x) dx. \end{aligned}$$

This system has been solved by our method with  $N = 10, 30, 60$ , the absolute errors at the particular grid points are tabulated in Table I, and compared with the absolute errors obtained by [4], [6]. This table verified that our results are more accurate in comparison with these methods.

Example 2. Consider the system of following linear Volterra integral equations with exact solution

$$\begin{aligned} (f_1(t), f_2(t))^T &= (\cos t, \sin t)^T, \\ f_1(t) &= -1 + \cos t - t \cos t^2 - \frac{1}{2}(-2 + t) \sin t + \cos t (1 + \sin t) \\ &\quad + \int_0^t (\sin(t-x) - 1) f_1(x) dx + \int_0^t (1 - x \cos t) f_2(x) dx, \\ f_2(t) &= -t + \sin t + \int_0^t f_1(x) dx + \int_0^t (t-x) f_2(x) dx. \end{aligned}$$

This system has been solved by our method with  $N = 10, 20, 30$ , the absolute errors at the particular grid points are tabulated in Table II, and compared with the absolute errors obtained by [7]. This table verified that our results are more accurate in comparison with [7].

Example 3. Consider the system of following nonlinear Fredholm integral equations with exact solution, [8],

$$\begin{aligned} (f_1(t), f_2(t))^T &= (t - 1, t^2)^T, \\ f_1(t) &= \frac{-10}{9} + \frac{11}{12} + \int_0^1 (txf_1^2(x) + x^2f_2^3(x)) dx, \\ f_2(t) &= \frac{1}{20} + \frac{t}{5} + t^2 + \int_0^1 (xf_1^3(x) - tf_2^2(x)) dx. \end{aligned}$$

The absolute errors at the particular grid points are tabulated in Table III. This table verified that our results are more accurate.

Example 4. Consider the system of following nonlinear Volterra integral equation with exact solution

$$\begin{aligned} (f_1(t), f_2(t))^T &= (t, t)^T, \\ f_1(t) &= t - t^2 + \int_0^t (f_1(x) + f_2(x)) dx, \\ f_2(t) &= t - \frac{t^2}{2} - \frac{t^3}{3} + \int_0^t (f_1^2(x) + f_2(x)) dx. \end{aligned}$$

The absolute errors at the particular grid points are tabulated in Table IV and compared with the absolute errors obtained by with [3], [5], [19]. This table verified that our results are more accurate in comparison.

TABLE I  
 THE ERROR  $\|E\|$  IN SOLUTION OF EXAMPLE 1 AT PARTICULAR POINTS

t	Our method N = 10	Our method N = 30	Our method N = 60	method in [6]	method in [4]
Results f1(x)					
0	1.71E-05	2.11E-07	1.32E-08	1.13E-03	1.2E-03
0.1	1.78E-05	2.19E-07	1.37E-08	1.15E-03	1.15E-03
0.2	1.83E-05	2.26E-07	1.41E-08	1.19E-03	1.19E-03
0.3	1.87E-05	2.31E-07	1.44E-08	1.29E-03	1.28E-03
0.4	1.87E-05	2.32E-07	1.45E-08	1.44E-03	1.44E-03
0.5	1.84E-05	2.29E-07	1.43E-08	1.67E-03	1.69E-03
0.6	1.77E-05	2.19E-07	1.37E-08	1.72E-03	1.70E-03
0.7	1.63E-05	2.02E-07	1.26E-08	1.77E-03	1.77E-03
0.8	1.41E-05	1.75E-07	1.09E-08	1.93E-03	1.93E-03
0.9	1.09E-05	1.35E-07	8.46E-09	2.19E-03	2.20E-03
1	6.51E-06	8.01E-08	5.01E-09	2.61E-02	4.21E-02
Results f2(x)					
0	3.73E-06	4.61E-08	2.88E-09	6.70E-04	6.67E-04
0.1	3.60E-06	4.45E-08	2.78E-09	7.31E-04	7.31E-04
0.2	3.59E-06	4.44E-08	2.77E-09	7.43E-04	7.43E-04
0.3	3.75E-06	4.63E-08	2.89E-09	7.13E-04	7.13E-04
0.4	4.15E-06	5.13E-08	3.21E-09	6.60E-04	6.49E-04
0.5	4.87E-06	6.03E-08	3.77E-09	5.55E-04	5.55E-04
0.6	6.04E-06	7.47E-08	4.67E-09	5.43E-04	5.44E-04
0.7	7.77E-06	9.62E-08	6.01E-09	4.92E-04	4.92E-04
0.8	1.02E-05	1.26E-07	7.92E-09	4.02E-04	4.02E-04
0.9	1.36E-05	1.68E-07	1.05E-08	2.78E-04	2.30E-04
1	1.82E-05	2.22E-07	1.41E-08	1.23E-04	1.14E-03

TABLE II  
 THE ERROR  $\|E\|$  IN SOLUTION OF EXAMPLE 2 AT PARTICULAR POINTS

t	Our method N = 10	Our method N = 20	Our method N = 30	method in [7]
Results f1(x)				
0	1.25E-06	7.78E-08	1.53E-08	0
0.1	1.03E-06	6.39E-08	1.26E-08	1.37E-04
0.2	7.78E-07	4.83E-08	9.53E-09	9.27E-04
0.3	5.01E-07	3.10E-08	6.12E-09	2.67E-03
0.4	1.99E-07	1.22E-08	2.41E-09	5.45E-03
0.5	1.22E-07	7.80E-09	1.54E-09	9.23E-03
0.6	4.61E-07	2.89E-08	5.72E-09	1.38E-02
0.7	8.15E-07	5.10E-08	1.01E-08	1.93E-02
0.8	1.17E-06	7.37E-08	1.45E-08	2.56E-02
0.9	1.55E-06	9.68E-08	1.91E-08	3.32E-02
1	1.92E-06	1.20E-07	2.37E-08	4.19E-02
Results f2(x)				
0	1.01E-06	6.32E-08	1.25E-08	0
0.1	1.26E-06	7.92E-08	1.56E-08	1.52E-04
0.2	1.52E-06	9.51E-08	1.88E-08	1.14E-03
0.3	1.77E-06	1.11E-07	2.19E-08	3.71E-03
0.4	2.03E-06	1.26E-07	2.51E-08	8.57E-03
0.5	2.28E-06	1.43E-07	2.82E-08	1.64E-02
0.6	2.53E-06	1.58E-07	3.13E-08	2.78E-02
0.7	2.79E-06	1.74E-07	3.44E-08	4.25E-02
0.8	3.05E-06	1.90E-07	3.76E-08	5.91E-02
0.9	3.30E-06	2.06E-07	4.07E-08	7.48E-02
1	3.55E-06	2.22E-07	4.38E-08	8.71E-02

TABLE III  
 THE ERROR  $\|E\|$  IN SOLUTION OF EXAMPLE 3 AT PARTICULAR POINTS

t	N = 10	N = 30	N = 60
Results f1(x)			
0	1.94E-04	2.44E-06	1.52E-07
0.1	1.88E-04	2.37E-06	1.48E-07
0.2	1.82E-04	2.30E-06	1.44E-07
0.3	1.77E-04	2.23E-06	1.39E-07
0.4	1.71E-04	2.16E-06	1.35E-07
0.5	1.65E-04	2.09E-06	1.31E-07
0.6	1.60E-04	2.02E-06	1.26E-07
0.7	1.55E-04	1.95E-06	1.22E-07
0.8	1.49E-04	1.88E-06	1.17E-07
0.9	1.43E-04	1.81E-06	1.13E-07
1	1.38E-04	1.74E-06	1.09E-07
Results f2(x)			
0	5.62E-05	7.05E-07	4.41E-08
0.1	5.28E-05	6.63E-07	4.15E-08
0.2	4.94E-05	6.21E-07	3.88E-08
0.3	4.60E-05	5.78E-07	3.62E-08
0.4	4.26E-05	5.36E-07	3.35E-08
0.5	3.92E-05	4.93E-07	3.09E-08
0.6	3.58E-05	4.51E-07	2.82E-08
0.7	3.25E-05	4.09E-07	2.56E-08
0.8	2.91E-05	3.66E-07	2.29E-08
0.9	2.57E-05	3.24E-07	2.03E-08
1	2.23E-05	2.82E-07	1.76E-08

TABLE IV  
 THE ERROR  $\|E\|$  IN SOLUTION OF EXAMPLE 4 AT PARTICULAR POINTS

t	Our method N = 10	RungeKutta method [5]	Adomian method [5]	Modified method [3]	HPM method [19]
Results f1(x)					
0	9.36E-17	0	0	0	
0.1	5.55E-17	1.82E-08	0	8.33E-04	9.67E-07
0.2	5.55E-16	1.17E-02	4.0E-06	1.57	3.0E-05
0.3	1.11E-16	2.45E-02	3.80E-05	3.13E-03	2.21E-04
0.4	1.11E-16	2.98E-02	2.23E-04	7.97E-02	9.05E-04
0.5	2.22E-16	2.65E-02	8.90E-04	8.90E-02	2.69E-03
0.6	2.22E-05	1.07E-02	2.77E-03	8.74E-02	1.37E-03
0.7	0	2.53E-02	7.20E-03	7.12E-02	2.61E-02
0.8	5.55E-16	9.72E-02	1.65E-02	7.79E-02	2.93E-02
0.9	3.33E-16	0.23	3.40E-02	8.90E-03	3.60E-02
1	0	0.52	0.13		6.87E-02
Results f2(x)					
0	1.35E-16	0	0	0	
0.1	8.33E-17	8.71E-08	2.10E-06	3.58E-04	4.71E-05
0.2	0	6.22E-03	6.70E-05	1.33E-02	7.11E-04
0.3	1.66E-16	1.84E-02	5.19E-04	1.82E-02	3.40E-03
0.4	1.11E-16	2.83E-02	2.24E-03	1.37E-02	1.02E-02
0.5	2.22E-16	3.21E-02	7.03E-03	1.73E-02	2.35E-02
0.6	3.33E-16	2.16E-02	1.78E-02	9.70E-03	4.61E-02
0.7	2.22E-16	1.62E-02	3.98E-02	1.15E-02	8.08E-02
0.8	1.11E-16	1.07E-01	7.94E-02	3.09E-02	0.13
0.9	4.44E-16	0.31	0.15	2.35E-02	0.19
1	0	1.75	0.25	9.37E-02	0.29

REFERENCES

- [1] Atkinson, K. E: The Numerical Solution of Integral Equations of the Second Kind, Cambridge University Press, New York, (1997).
- [2] Delves, L. M , Mohammed, J. L: Computational Methods for Integral Equations , Cambridge University Press, Cambridge, (1985).
- [3] Saeed, R. K , Ahmed, C. S: Approximate Solution for the System of Non-linear Volterra Integral Equations of the Second Kind by using Block-by-block Method, Australian Journal of Basic and Applied Sciences, 2(1), 114-124(2008).
- [4] Babolian, E , Masouri, Z, Hatamzadeh-Varmazyar, S: A Direct Method for Numerically Solving Integral Equations System Using Orthogonal Triangular Functions, Int. J. Industrial Mathematics, 1(2), 135-145(2009).
- [5] Maleknejad, K, Shahrezaee, M: Using RungeKutta method for numerical solution of the system of Volterra integral equation, Applied Mathematics and Computation, 149, 399-410(2004).
- [6] Babolian, E , Mordad, M: A numerical method for solving systems of linear and nonlinear integral equations of the second kind by hat basis functions, Computers and Mathematics with Applications, 62, 187-198(2011).
- [7] Rabbani, M, Maleknejad, K , Aghazadeh, N: Numerical computational solution of the Volterra integral equations system of the second kind by using an expansion method, Applied Mathematics and Computation, 187, 1143-1146(2007).
- [8] Bakodah, H. O: Some Modifications of Adomian Decomposition Method Applied to Nonlinear System of Fredholm Integral Equations of the Second Kind, Int. J. Contemp. Math. Sciences, 7(19), 929 - 942(2012)
- [9] Prenter, P.M: Spline and Variational Methods, Wiley & Sons, New-York, (1975).
- [10] Mahmoodi, Z, Rashidinia, J, Babolian, E: B-Spline collocation method for linear and nonlinear Fredholm and Volterra integro-differential equations, Applicable Analysis, 1-16(2012).
- [11] Rashidinia, J, Babolian, E, Mahmoodi, Z : Spline Collocation for Fredholm Integral Equations, Mathematical Sciences, 5(2), 147-158(2011).
- [12] Rashidinia, J, Babolian, E, Mahmoodi, Z: Spline Collocation for nonlinear Fredholm Integral Equations, International Journal of Mathematical Modelling & Computations, 1(1), 69-75(2011).

- [13] Babolian,E, Biazar,J, Vahidi,A.R: The decomposition method applied to systems of Fredholm integral equations of the second kind, *Appl. Math. Comput.*, 148 ,443-452(2004).
- [14] Maleknejad,K , Mirzaee,F: Numerical solution of linear Fredholm integral equations system by rationalized Haar functions method, *Int. J. Comput. Math.*, 80(11),1397-1405(2003).
- [15] Maleknejad,K, Shahrezaee,M,Khatami,H: Numerical solution of integral equations system of the second kind by Block-Pulse functions, *Appl. Math. Comput.* ,166,15-24(2005).
- [16] Maleknejad,K,Aghazadeh,N, Rabbani,M: Numerical solution of second kind Fredholm integral equations system by using a Taylor-series expansion method, *Appl. Math. Comput.*,175,1229-1234(2006).
- [17] Maleknejad,K,Mahmoudi,Y: Numerical solution of linear Fredholm integral equation by using hybrid Taylor and block-pulse functions, *Appl. Math. Comput.*,149,799-806(2004).
- [18] Babolian,E, Fattahzadeh,F: Numerical computation method in solving integral equations by using Chebyshev wavelet operational matrix of integration, *Appl. Math. Comput.*,188,1016-1022(2007).
- [19] E. Yusufu Glu,E: A homotopy perturbation algorithm to solve a system of Fredholm-Volterra type integral equations, *Mathematical and Computer Modelling*,47, 1099-1107(2008).
- [20] Matinfar,M,Saeidy,M,Vahidi,J: Application of Homotopy Analysis Method for Solving Systems of Volterra Integral Equations , *Adv. Appl. Math. Mech.*, 4 ,36-45(2012).
- [21] Javidi,M,Golbabai,A: A numerical solution for solving system of Fredholm integral equations by using homotopy perturbation method, *Appl. Math. Comput.*,189,1921-1928(2007).
- [22] Biazar,J, Eslami,M: Modified HPM for solving systems of Volterra integral equations of the second kind ,*Journal of King Saud University* , 23,35-39(2011).
- [23] Muhammad,M,Nurmuhammad,A,Mori,M, M. Sugihara,M: Numerical solution of integral equations by means of the Sinc collocation method based on the double exponential transformation, *J. Comput. Appl. Math.*,177 ,269-286(2005).
- [24] Rashidinia,J,Zarebnia,M: Convergence of approximate solution of system of Fredholm integral equations, *J. Math. Anal. Appl.*,333,1216-1227(2007).
- [25] Wazwaz,A. M: *A First Course in Integral Equations*, WSPC, New Jersey, (1997).
- [26] Biazar,J: Solution of systems of integral-differential equations by Adomian decomposition method, *Appl. Math. Comput.*,168,1232-1238(2005).
- [27] Jafari,H, Daftardar-Gejji,V: Solving a system of nonlinear fractional differential equations using Adomian decomposition, *J. Comput. Appl. Math.*,196 ,644-651(2006).
- [28] Vahidi,A.R,Mokhtar,M: On the decomposition method for system of linear Fredholm integral equations of the second kind, *Appl. Math. Sci.*, 2,57-62(2008).