

## Termination of the Brachial Artery in the Arm and Its Clinical Significance

**Authors :** Ramya Rathan, Miral N. F. Salama

**Abstract :** The variations in the arteries have been drawing attention of anatomists for a long time because of their clinical significance. The brachial artery is the principal artery of the arm which is the continuation of the axillary artery from the lower border of the Teres Major. It terminates into the radial and ulnar arteries below the elbow joint at the neck radius. The present study aims at exploring the clinical significance of the high termination of the brachial artery. During the routine cadaveric dissection of the arm, for the undergraduate students of medicine at our university, we observed a high bifurcation of the radial and the ulnar artery at the midshaft of the humerus. The median nerve was seen passing between these two junctions. Further, the course and the relations of this artery were studied. The accurate knowledge regarding these kinds of variation in the blood vessels is mandatory for planning of designing. General physicians, surgeons and radiologists should keep in mind the variations in the branching pattern of the arteries in their daily medical, diagnostic and therapeutic procedures to avoid complications in diagnostic and surgical procedures.

**Keywords :** brachial artery, high termination, radial artery, ulnar artery

**Conference Title :** ICMMA 2019 : International Conference on Microscopic and Macroscopic Anatomy

**Conference Location :** Barcelona, Spain

**Conference Dates :** February 11-12, 2019