

Social Ties and the Prevalence of Single Chronic Morbidity and Multimorbidity among the Elderly Population in Selected States of India

Authors : Sree Sanyal

Abstract : Research in ageing often highlights the age-related health dimension more than the psycho-social characteristics of the elderly, which also influences and challenges the health outcomes. Multimorbidity is defined as the person having more than one chronic non-communicable diseases and their prevalence increases with ageing. The study aims to evaluate the influence of social ties on self-reported prevalence of multimorbidity (selected chronic non-communicable diseases) among the selected states of elderly population in India. The data is accessed from Building Knowledge Base on Population Ageing in India (BKPAI), collected in 2011 covering the self-reported chronic non-communicable diseases like arthritis, heart disease, diabetes, lung disease with asthma, hypertension, cataract, depression, dementia, Alzheimer's disease, and cancer. The data of the above diseases were taken together and categorized as: 'no disease', 'one disease' and 'multimorbidity'. The predicted variables were demographic, socio-economic, residential types, and the variable of social ties includes social support, social engagement, perceived support, connectedness, and importance of the elderly. Predicted probability for multiple logistic regression was used to determine the background characteristics of the old in association with chronic morbidities showing multimorbidity. The finding suggests that 24.35% of the elderly are suffering from multimorbidity. Research shows that with reference to 'no disease', according to the socio-economic characteristics of the old, the female oldest old (80+) from others in caste and religion, widowed, never had any formal education, ever worked in their life, coming from the second wealth quintile standard, from rural Maharashtra are more prone with 'one disease'. From the social ties background, the elderly who perceives they are important to the family, after getting older their decision-making status has been changed, prefer to stay with son and spouse only, satisfied with the communication from their children are more likely to have less single morbidity and the results are significant. Again, with respect to 'no disease', the female oldest old (80+), who are others in caste, Christian in religion, widowed, having less than 5 years of education completed, ever worked, from highest wealth quintile, residing in urban Kerala are more associated with multimorbidity. The elderly population who are more socially connected through family visits, public gatherings, gets support in decision making, who prefers to spend their later years with son and spouse only but stays alone shows lesser prevalence of multimorbidity. In conclusion, received and perceived social integration and support from associated neighborhood in the older days, knowing about their own needs in life facilitates better health and wellbeing of the elderly population in selected states of India.

Keywords : morbidity, multi-morbidity, prevalence, social ties

Conference Title : ICHAA 2019 : International Conference on Healthy and Active Aging

Conference Location : Toronto, Canada

Conference Dates : June 17-18, 2019