

Depressive Symptoms in Children with Epilepsy Attending a Tertiary Care Hospital in Oman

Authors : Hamood Al Kiyumi, Salim Al Huseini, Khalid Al Risi, Hassan Mirza, Amira Al Hosni, Sanjay Jaju, Asaad Al Habsi

Abstract : Objectives: The aim of this study was to assess the proportion of depressive symptoms along with demographic data in children diagnosed with epilepsy in a tertiary care institution in Oman. Methods: This cross-sectional study was conducted between June 2016 and August 2018. We have included 75 children with age group from five to 12 years old, attending epilepsy clinic at Sultan Qaboos University Hospital who were diagnosed with epilepsy and already on treatment. Patients were excluded if they have mental retardation. Validated Depression Scale for Children (CES-DC) questionnaire was utilized to assess the level of depressive symptoms among children. In addition, we have looked at associated factors including seizure status in the last three months, compliance with antiepileptic medications, type of epilepsy, and number of antiepileptic medications. Results: In this study, we found that depressive symptoms were present in 39 (52%) of patients. We also found that 96% of the patients were compliant to medications. In addition, seizure was present in the last three months in 48% of the sample studies. There was no statistically significant association between any of the studied variables and depression. Conclusions: Although depression is highly prevalent in children with epilepsy, this study did not find any significant association between the CES-DC scores and the studied factors.

Keywords : depression, children, epilepsy, Oman

Conference Title : ICPMH 2019 : International Conference on Positive Mental Health

Conference Location : Tokyo, Japan

Conference Dates : March 25-26, 2019