

## Assessment of Predictive Confounders for the Prevalence of Breast Cancer among Iraqi Population: A Retrospective Study from Baghdad, Iraq

**Authors :** Nadia H. Mohammed, Anmar Al-Taie, Fadia H. Al-Sultany

**Abstract :** Although breast cancer prevalence continues to increase, mortality has been decreasing as a result of early detection and improvement in adjuvant systemic therapy. Nevertheless, this disease required further efforts to understand and identify the associated potential risk factors that could play a role in the prevalence of this malignancy among Iraqi women. The objective of this study was to assess the perception of certain predictive risk factors on the prevalence of breast cancer types among a sample of Iraqi women diagnosed with breast cancer. This was a retrospective observational study carried out at National Cancer Research Center in College of Medicine, Baghdad University from November 2017 to January 2018. Data of 100 patients with breast cancer whose biopsies examined in the National Cancer Research Center were included in this study. Data were collected to structure a detailed assessment regarding the patients' demographic, medical and cancer records. The majority of study participants (94%) suffered from ductal breast cancer with mean age 49.57 years. Among those women, 48.9% were obese with body mass index (BMI)  $35 \text{ kg/m}^2$ . 68.1% of them had positive family history of breast cancer and 66% had low parity. 40.4% had stage II ductal breast cancer followed by 25.5% with stage III. It was found that 59.6% and 68.1% had positive oestrogen receptor sensitivity and positive human epidermal growth factor (HER2/neu) receptor sensitivity respectively. In regard to the impact of prediction of certain variables on the incidence of ductal breast cancer, positive family history of breast cancer ( $P < 0.0001$ ), low parity ( $P < 0.0001$ ), stage I and II breast cancer ( $P = 0.02$ ) and positive HER2/neu status ( $P < 0.0001$ ) were significant predictive factors among the study participants. The results from this study provide relevant evidence for a significant positive and potential association between certain risk factors and the prevalence of breast cancer among Iraqi women.

**Keywords :** Ductal Breast Cancer, Hormone Sensitivity, Iraq, Risk Factors

**Conference Title :** ICBMP 2019 : International Conference on Biomedicine and Medical Pharmacology

**Conference Location :** Dublin, Ireland

**Conference Dates :** March 21-22, 2019