

Postpartum Female Sexual Dysfunctions in Hungary: A Cross-Sectional Study

Authors : Katalin Szöllősi, László Szabó

Abstract : Introduction and purpose: Even though female sexual dysfunctions are common among women in the postpartum period, the profile of these disturbances has not been well investigated in Hungary yet. The aim of the study was to evaluate the postpartum female sexual functions in Hungary. This research sought to investigate the possible predictor factors which can influence postpartum female sexual functions. Method and sample: This was a cross-sectional study, including patients from two maternity clinics in Budapest. 113 women were recruited into our study 3 months after their childbirth. 53 had vaginal birth, 60 had a caesarian section. Data were collected from medical reports in addition by using self-developed questions and validated questionnaires in order to measure important predictors which may be responsible for postpartum sexual dysfunctions such as mode of delivery, parity, urinary incontinence and body image. Sexual functions were evaluated by the Hungarian version of the Female Sexual Function Index (FSFI). The Hungarian version of Body Image Questionnaire-Short Form14 (BSQ-SF14) was applied for assessing body image. Results: 82,3% of the participants began to have sexual intercourse within three months postpartum. 53,98% of the participants reported sexual dysfunctions (cut-off FSFI score 26,55). According to our results mode of delivery, parity, hemorrhoids, time of intercourse, resumption was not associated with female sexual dysfunctions. We found correlation at a tendential level between urinary incontinence and sexual dysfunctions ($p=0,003$, $R=0,26$). We found a negative correlation at a tendential level between the total score of BSQ-SF14 and FSFI ($p=0,03$, $R=-0,269$). Only 32,74% of women reported discussing sexual life with health care professionals. However, 67,25% of them would have had the need to be asked about their postpartum health issues. Conclusions and recommendations: The prevalence of female sexual dysfunctions were relatively high after childbirth. We found that incontinence and body image was associated with sexual dysfunctions; other risk factors remained unknown. Despite regular contact with health care professionals, women rarely get any information about postpartum sexual health issues. The high prevalence of dysfunctions indicates the need for further investigation to address other risk factors and proper counselling of women after childbirth.

Keywords : body image, postpartum, sexual dysfunction, urinary incontinence

Conference Title : ICPS 2019 : International Conference on Psychiatry and Sexology

Conference Location : Tokyo, Japan

Conference Dates : March 23-24, 2020