Protective Effect of Hesperidin against Cyclophosphamide Hepatotoxicity in Rats

Authors : Amr A. Fouad, Waleed H. Albuali, Iyad Jresat

Abstract : The protective effect of hesperidin was investigated in rats exposed to liver injury induced by a single intraperitoneal injection of cyclophosphamide (CYP) at a dose of 150 mg kg-1. Hesperidin treatment (100 mg kg-1/day, orally) was applied for seven days, starting five days before CYP administration. Hesperidin significantly decreased the CYP-induced elevations of serum alanine aminotransferase, and hepatic malondialdehyde and myeloperoxidase activity, significantly prevented the depletion of hepatic glutathione peroxidase activity resulted from CYP administration. Also, hesperidin ameliorated the CYP-induced liver tissue injury observed by histopathological examination. In addition, hesperidin decreased the CYP-induced expression of inducible nitric oxide synthase, tumor necrosis factor- α , cyclooxygenase-2, Fas ligand, and caspase-9 in liver tissue. It was concluded that hesperidin may represent a potential candidate to protect against CYP-induced hepatotoxicity.

Keywords : hesperidin, cyclophosphamide, liver, rats

Conference Title : ICABB 2014 : International Conference on Applied Biology and Biotechnology

Conference Location : Stockholm, Sweden

Conference Dates : July 14-15, 2014

1