

Identification and Quantification of Sesquiterpene Lactones of Sagebrush (*Artemisia tridentate*) and Its Chemical Modification

Authors : Rosemary Anibogwu, Kavita Sharma, Karl De Jesus

Abstract : Sagebrush is an abundant and naturally occurring plant in the Intermountain West region of the United States. The plant contains an array of bioactive compounds such as flavonoids, terpenoids, sterols, and phenolic acids. It is important to identify and characterize these compounds because Native Americans use sagebrush as herbal medicine. These compounds are also utilized for preventing infection in wounds, treating headaches and colds, and possess antitumor properties. This research is an exploratory study on the sesquiterpene present in the leaves of sagebrush. The leaf foliage was extracted with 100 % chloroform and 100 % methanol. The percentage yield for the crude was considerably higher in chloroform. The Thin Layer Chromatography (TLC) analysis of the crude extracted unveiled a brown band at $R_f = 0.25$ and a dark brown band at $R_f = 0.74$, along with three unknown faint bands the 254 nm UV lamp. Furthermore, the two distinct brown (Achillin) and dark brown band (Hydroxyachillin) in TLC were further utilized in the isolation of pure compounds with column chromatography. The structures of Achillin and Hydroxyachillin were elucidated based on extensive spectroscopic analysis, including TLC, High-Performance Liquid Chromatography (HPLC), 1D- and 2D-Nuclear Magnetic Resonance (NMR), and Mass Spectroscopy (MS). The antioxidant activities of crude extract and three pure compounds were evaluated in terms of their peroxy radical scavenging by Ferric Reducing Ability of Plasma (FRAP) and 1,1-Diphenyl-2-picryl-hydrazyl (DPPH) methods. The crude extract showed the antioxidant activity of $18.99 \pm 0.51 \mu\text{mol TEg}^{-1} \text{FW}$ for FRAP and $11.59 \pm 0.38 \mu\text{mol TEg}^{-1} \text{FW}$ for DPPH. The activities of Achillin, Hydroxyachillin, and Quercetagenin trimethyl ether were 13.03, 15.90 and 14.02 $\mu\text{mol TEg}^{-1} \text{FW}$ respectively for the FRAP assay. The three purified compounds have been submitted to the National Cancer Institute 60 cancer cell line for further study.

Keywords : HPLC, nuclear magnetic resonance spectroscopy, sagebrush, sesquiterpene lactones

Conference Title : ICSRD 2020 : International Conference on Scientific Research and Development

Conference Location : Chicago, United States

Conference Dates : December 12-13, 2020