

Determinants of Post-Psychotic Depression in Schizophrenia Patients in ACSH and Mekelle Hospital Tigray, Ethiopia, 2019

Authors : Ashenafi Ayele, Shewit Haftu, Tesfalem Araya

Abstract : Background: "Post-psychotic depression", "post schizophrenic depression", and "secondary depression" have been used to describe the occurrence of depressive symptoms during the chronic phase of schizophrenia. Post-psychotic depression is the most common cause of death due to suicide in schizophrenia patients. Overall lifetime risk for patients with schizophrenia is 50% for suicide attempts and 9-13% lifetime risk for completed suicide and also it is associated with poor prognosis and poor quality of life. Objective: To assess determinant of post psychotic depression in schizophrenia patients ACSH and Mekelle General Hospital, Tigray Ethiopia 2019. Methods: An institutional based unmatched case control study was conducted among 69 cases and 138 controls with the ratio of case to control 1 ratio 2. The sample is calculated using epi-info 3.1 to assess the determinant factors of post-psychotic depression in schizophrenia patients. The cases were schizophrenia patients who have been diagnosed at least for more than one-year stable for two months, and the controls are any patients who are diagnosed as schizophrenia patients. Study subjects were selected using a consecutive sampling technique. The Calgary depression scale for schizophrenia self-administered questionnaire was used. Before the interview, it was assessed the client's capacity to give intended information using a scale called the University of California, San Diego Brief Assessment of Capacity to Consent (UBACC). Bivariant and multiple Logistic regression analysis was performed to determine between the independent and dependent variables. The significant independent predictor was declared at 95% confidence interval and P-value of less than 0.05. Result: Females were affected by post psychotic depression with the (AOR=2.01, 95%CI: 1.003- 4.012, P=0.49).Patients who have mild form of positive symptom of schizophrenia affected by post psychotic depression with (AOR =4.05, 95%CI: 1.888- 8.7.8, P=0001).Patients who have minimal form of negative symptom of schizophrenia are affected by post psychotic depression with (AOR =4.23, 95%CI: 1.081-17.092, P=.038). Conclusion: In this study, sex (female) and presence of positive and negative symptoms of schizophrenia were significantly associated. It is recommended that the post psychotic depression should be assessed in every schizophrenia patient to decrease the severity of illness, and to improve patient's quality of life.

Keywords : determinants, post-psychotic depression, Mekelle city

Conference Title : ICNDPD 2020 : International Conference on Neuropsychiatric Disorders and Psychiatric Disorder

Conference Location : Toronto, Canada

Conference Dates : June 18-19, 2020