A Case Report: The Role of Gut Directed Hypnotherapy in Resolution of Irritable Bowel Syndrome in a Medication Refractory Pediatric Male Patient

Authors: Alok Bapatla, Pamela Lutting, Mariastella Serrano

Abstract: Background: Irritable Bowel Syndrome (IBS) is a functional gastrointestinal disorder characterized by abdominal pain associated with altered bowel habits in the absence of an underlying organic cause. Although the exact etiology of IBS is not fully understood, one of the leading theories postulates a pathology within the Brain-Gut Axis that leads to an overall increase in gastrointestinal sensitivity and pejorative changes in gastrointestinal motility. Research and clinical practice have shown that Gut Directed Hypnotherapy (GDH) has a beneficial clinical role in improving Mind-Gut control and thereby comorbid conditions such as anxiety, abdominal pain, constipation, and diarrhea. Aims: This study presents a 17-year old male with underlying anxiety and a one-year history of IBS-Constipation Predominant Subtype (IBS-C), who has demonstrated impressive improvement of symptoms following GDH treatment following refractory trials with medications including bisacodyl, senna, docusate, magnesium citrate, lubiprostone, linaclotide. Method: The patient was referred to a licensed clinical psychologist specializing in clinical hypnosis and cognitive-behavioral therapy (CBT), who implemented "The Standardized Hypnosis Protocol for IBS" developed by Dr. Olafur S. Palsson, Psy.D at the University of North Carolina at Chapel Hill. The hypnotherapy protocol consisted of a total of seven weekly 45-minute sessions supplemented with a 20-minute audio recording to be listened to once daily. Outcome variables included the GAD-7, PHQ-9 and DCI-2, as well as self-ratings (ranging 0-10) for pain (intensity and frequency), emotional distress about IBS symptoms, and overall emotional distress. All variables were measured at intake prior to administration of the hypnosis protocol and at the conclusion of the hypnosis treatment. A retrospective IBS Questionnaire (IBS Severity Scoring System) was also completed at the conclusion of the GDH treatment for pre-and post-test ratings of clinical symptoms. Results: The patient showed improvement in all outcome variables and self-ratings, including abdominal pain intensity, frequency of abdominal pain episodes, emotional distress relating to gut issues, depression, and anxiety. The IBS Questionnaire showed a significant improvement from a severity score of 400 (defined as severe) prior to GDH intervention compared to 55 (defined as complete resolution) at four months after the last session. IBS Questionnaire subset questions that showed a significant score improvement included abdominal pain intensity, days of pain experienced per 10 days, satisfaction with bowel habits, and overall interference of life affected by IBS symptoms. Conclusion: This case supports the existing research literature that GDH has a significantly beneficial role in improving symptoms in patients with IBS. Emphasis is placed on the numerical results of the IBS Questionnaire scoring, which reflects a patient who initially suffered from severe IBS with failed response to multiple medications, who subsequently showed full and sustained

Keywords: pediatrics, constipation, irritable bowel syndrome, hypnotherapy, gut-directed hypnosis **Conference Title:** ICFGD 2022: International Conference on Functional Gastrointestinal Disorders

Conference Location : Tokyo, Japan **Conference Dates :** January 07-08, 2022