

Phenotype of Cutaneous Squamous Cell Carcinoma in a Brazilian City with a Tropical Climate

Authors : Julia V. F. Cortes, Maria E. V. Amarante, Carolina L. Cerdeira, Roberta B. V. Silva

Abstract : Nonmelanoma skin cancer is more commonly diagnosed than all other malignancies combined. In that group, cutaneous squamous cell carcinoma stands out for having the highest probability of metastasis and recurrence after treatment, in addition to being the second most prevalent form of skin cancer. Its main risk factors include exposure to carcinogens, such as ultraviolet radiation related to sunlight exposure, smoking, alcohol consumption, and human papillomavirus (HPV) infection. Considering the increased risk of skin cancer in the Brazilian population, caused by the high incidence of solar radiation, and the importance of identifying risk phenotypes for the accomplishment of public health actions, an epidemiological study was conducted in a city with a tropical climate located in southeastern Brazil, aiming to identify the target population and assist in primary and secondary prevention. This study describes the profile of patients with cutaneous squamous cell cancer, correlating the variables, sex, age, and differentiation. The study used as primary data source the results of anatomopathological exams delivered from January 2015 to December 2019 for patients registered at one pathology service, which analyzes the results of biopsies. Thus, 66 patients with cutaneous squamous cell carcinoma were analyzed. The most affected age group was 60 years or older (78.79%), emphasizing that moderately differentiated (79.49%) and well-differentiated forms (66.67%) are prevalent in this age group, resulting in a difference of 12.82 percentage points between them. In addition, the predominant sex was male (58%), and it was found that half of the women and 65.79% of men had a moderately differentiated type, whereas the well-differentiated type was slightly more frequent in women. It is worth noting that the moderately differentiated subtype has a 59.20% prevalence among all cases. Thus, it was concluded that the most affected age group was 60 years or older and that men were more affected. As for the subtype, the moderately differentiated one, which is recognized for presenting the second-highest risk for metastasis, was prevalent in this study, affecting 6.6% more men and predominating in the elderly.

Keywords : cutaneous squamous cell carcinoma, epidemiology, skin cancer, spinal cell cancer

Conference Title : ICCDDD 2021 : International Conference on Clinical Dermatology and Dermatological Diseases

Conference Location : New York, United States

Conference Dates : December 09-10, 2021