

A Randomized, Controlled Trial To Test Behavior Change Techniques (BCTs) To Improve Low Intensity Physical Activity In Older Adults

Authors : Ciaran Friel, Jerry Suls, Patrick Robles, Frank Vicari, Joan Duer-Hefele, Karina W. Davidson

Abstract : Physical activity guidelines focus on increasing moderate intensity activity for older adults, but adherence to recommendations remains low. This is despite the fact that scientific evidence supports that any increase in physical activity is positively correlated with health benefits. Behavior change techniques (BCTs) have demonstrated effectiveness in reducing sedentary behavior and promoting physical activity. This pilot study uses a Personalized Trials (N-of-1) design to evaluate the efficacy of using four BCTs to promote an increase in low-intensity physical activity (2,000 steps of walking per day) in adults aged 45-75 years old. The 4 BCTs tested were goal setting, action planning, feedback, and self-monitoring. BCTs were tested in random order and delivered by text message prompts requiring participant response. The study recruited health system employees in the target age range, without mobility restrictions and demonstrating interest in increasing their daily activity by a minimum of 2,000 steps per day for a minimum of five days per week. Participants were sent a Fitbit Charge 4 fitness tracker with an established study account and password. Participants were recommended to wear the Fitbit device 24/7, but were required to wear it for a minimum of ten hours per day. Baseline physical activity was measured by the Fitbit for two weeks. Participants then engaged with a clinical research coordinator to review comprehension of the text message content and required actions for each of the BCTs to be tested. Participants then selected a consistent daily time in which they would receive their text message prompt. In the 8 week intervention phase of the study, participants received each of the four BCTs, in random order, for a two week period. Text message prompts were delivered daily at a time selected by the participant. All prompts required an interactive response from participants and may have included recording their detailed plan for walking or daily step goal (action planning, goal setting). Additionally, participants may have been directed to a study dashboard to view their step counts or compare themselves with peers (self-monitoring, feedback). At the end of each two week testing interval, participants were asked to complete the Self-Efficacy for Walking Scale (SEW_Dur), a validated measure that assesses the participant's confidence in walking incremental distances and a survey measuring their satisfaction with the individual BCT that they tested. At the end of their trial, participants received a personalized summary of their step data in response to each individual BCT. Analysis will examine the novel individual-level heterogeneity of treatment effect made possible by N-of-1 design, and pool results across participants to efficiently estimate the overall efficacy of the selected behavioral change techniques in increasing low-intensity walking by 2,000 steps, 5 days per week. Self-efficacy will be explored as the likely mechanism of action prompting behavior change. This study will inform the providers and demonstrate the feasibility of N-of-1 study design to effectively promote physical activity as a component of healthy aging.

Keywords : aging, exercise, habit, walking

Conference Title : ICEPAA 2021 : International Conference on Exercise and Physical Activity in Aging

Conference Location : New York, United States

Conference Dates : December 09-10, 2021