

Clinical Profile, Evaluation, Management and Visual Outcome of Idiopathic Intracranial Hypertension in a Neuro-Ophthalmology Clinic in Jeddah, Saudi Arabia

Authors : Rahaf Mandura

Abstract : Background: Idiopathic intracranial hypertension (IIH) is a disorder with elevated intracranial pressure (ICP) more than 250 mm H₂O, without evidence of meningeal inflammation, space-occupying lesion, or venous thrombosis. The aim of this research is to study the clinical profile, evaluation, management, and visual outcome in a hospital-based population of IIH cases in Jeddah. Methodology: This is a retrospective observational study that included the medical records of all patients referred to neuro-ophthalmology service for evaluation of papilledema. The medical records have been reviewed from October 2018 to February 2020 at Jeddah Eye Hospital (JEH), Saudi Arabia. A total of fifty-one patients presented with papilledema in the studied period. Forty-seven patients met our inclusion criteria and were included in the study. Results: Most of the patients were females (43, 91.5%) with a mean age of presentation of 30.83±11.40 years. The most common presenting symptom was headache (40 patients, 85.1%), followed by transient visual obscuration (20 patients, 42.6%), and reduced visual acuity (15 patients, 31.9%). All 47 patients were started on medical treatment with oral acetazolamide with four patients (8.5%) shifted to topiramate because of the lack of response or intolerance to acetazolamide while four patients (8.5%) underwent lumbar-peritoneal shunt because of inadequate control of the disease despite the treatment with medical therapy. For both eyes, the change in visual acuity across all assessment points was statistically significant. Nevertheless, there were no significant changes in the visual field findings among all of the compared assessment points. Conclusion: The present study has shown that IIH-related papilledema is common in young female patients with headaches, transient visual obscurations and reduced visual acuity. Those are the commonest symptoms in our IIH population. Medical treatment of IIH is significantly efficacious and should be considered in order to enhance the prognosis of IIH-related complications. Therefore, the visual status should be frequently monitored for these patients.

Keywords : idiopathic intracranial hypertension, intracranial hypertension, papilledema, headache

Conference Title : ICO 2021 : International Conference on Ophthalmology

Conference Location : Cairo, Egypt

Conference Dates : December 13-14, 2021