

Fear of Covid-19 a Major Contributing Factor to Insomnia in General Iranian Population

Authors : Amin Nakhostin-Ansari, Samaneh Akbarour, Khosro Sadeghniaat Haghighi, Zahra Banafsheh Alemohammad, Farnaz Etesam, Arezu Najafi, Mahnaz Khalafehnlsaz

Abstract : Introduction: The outbreak of coronavirus disease has considerably burdened the healthcare system in Iran. This study aimed to evaluate the characteristics of insomnia experienced by the general Iranian population during the COVID-19 pandemic. Method: A scale(FCV-19) was used for Fear of COVID-19, Insomnia Severity Index (ISI), Patient Health Questionnaire-2 (PHQ-2), and Generalized Anxiety Disorder Scale-2 (GAD-2) for detailed characterization of insomnia and its patterns Results: In total, 675 people with insomnia with the mean age of 40.28 years (SD=11.15) participated in this study. Prevalence of difficulty initiating sleep (DIS), difficulty maintaining sleep (DMS), and early morning awakening (EMA) were 91.4%, 86.7%, and 77%, respectively. DIS, DMS, and EMA were more common in people with depression and anxiety. FCV-19 score was higher in those with more severe types of DIS, DMS, and EMA ($P<0.001$). FCV-19 was a risk factor for all patterns of insomnia (OR=1.19, 1.12, 1.02 for DIS, DMS, and EMA, respectively). Conclusion: fear of COVID-19 is a major factor to insomnia patterns. Investigation of COVID-19 fear in people with insomnia and the addition of attributed relieving or management strategies to conventional management of insomnia are reasonable approaches to improve the sleep condition of people in the pandemic.

Keywords : insomnia, difficulty maintaining sleep, COVID-19, Coronavirus

Conference Title : ICONCE 2021 : International Conference on Outbreak of a Novel Coronavirus and Epidemiology

Conference Location : Copenhagen, Denmark

Conference Dates : June 10-11, 2021