Interdialytic Acupuncture Is an Add-on Option for Preserving Residual Renal Function: A Case Series Report

Authors: Lai Tzu-Hsuan, Lai Jung-Nien, Lin Jaung-Geng, Kao Shung-Te, Hsuan-Kuang Jung

Abstract : Background: Whether acupuncture therapy contributes to preserving residual renal function (RRF) remains largely unknown. This case series evidenced the potential beneficial effects of acupuncture for preserving RRF in five patients with the end-stage renal disease under hemodialysis (HD) treatment. Participants: Five patients on HD receiving eight sessions of weekly 30-min interdialytic acupuncture (Inter-A) with residual urine volume (rUV) and residual glomerular filtration rate (rGFR) recorded once every two weeks were included for analysis. Outcomes: Changes in rUV and rGFR calculated using 24-hour urine collection data were analyzed to assess RRF. Variations in hemoglobin, urea Kt/V and serum albumin levels measured monthly were analyzed to evaluate HD adequacy. Results: After eight Inter-A sessions, mean (standard deviation (SD)) rUV and rGFR increased from 612 (184) ml/day and 1.48 (.94) ml/min/1.73 m2 at baseline to 803(289) ml/day and 2.04(1.17) ml/min/1.73m2 at 2- and 4-week follow-up, respectively. The mean percentage difference increased by 31% in rUV and 38% in rGFR. Routine measurements on HD adequacy also showed improvement. Conclusions: Acupuncture might be an optional add-on treatment for HD population with poor control of water; however, further well-designed controlled trials are warranted.

Keywords: end-stage renal disease, hemodialysis, acupuncture, residual renal function, residual urine volume

Conference Title: ICUN 2021: International Conference on Urology and Nephrology

Conference Location: London, United Kingdom

Conference Dates: May 24-25, 2021