

Endocrine Therapy-Induced Alopecia in Patients with Breast Cancer in Tunisia

Authors : Aref Zribi, Sonia Ben Nasr, Sana Fendri, Mahdi Balti, Abderazzek Haddaoui

Abstract : Background: Despite their benefit, Endocrine therapies (ET) are known to have substantial adverse events (AEs) such as hot flashes, mood disorders and osteoarticular pain. ET induced alopecia(EIA) is less frequently noted by patients and is less reported in the literature. The aim of our study was to report ET alopecia characteristics and their influence on patient and treatment observance. Method: We conducted a retrospective study including luminal BC patients treated in the oncology department of the military hospital of Tunis between January 2015 and December 2020. Patients treated with previous chemotherapy-inducing alopecia were excluded. Results: 145 female patients were included. The median age was 59 years. EIA was reported in 44% of cases. Alopecia was attributed to aromatase inhibitors in 53% and tamoxifen in 21%. Severity was grade 1 in 80% and grade 2 in the remaining cases. ET discontinuation because of alopecia was noted in 6.5 % of patients. Moderate improvement of alopecia was observed with topical minoxidil and Thallium metallicum 9CH homeopathy during ET in 60% of patients. Conclusions: EIA is frequent in BC patients and should be considered to improve treatment observance and patients' quality of life.

Keywords : endocrine therapy, alopecia, breast cancer, Tunisia

Conference Title : ICG 2021 : International Conference on Gynecology

Conference Location : Sydney, Australia

Conference Dates : August 30-31, 2021