

Outcome of Emergency Response Team System in In-Hospital Cardiac Arrest

Authors : Jirapat Suriyachaisawat, Ekkit Surakarn

Abstract : Introduction: To improve early detection and mortality rate of In- Hospital Cardiac arrest, Emergency Response Team (ERT) system was planned and implemented since June 2009 to detect pre-arrest conditions and for any concerns. The ERT consisted of on duty physicians and nurses from emergency department. ERT calling criteria consisted of acute change of HR < 40 or > 130 beats per minute, systolic blood pressure < 90mmHg, respiratory rate <8 or > 28 breaths per minute, O2 saturation < 90%, acute change in conscious state, acute chest pain or worried about the patients. From the data on ERT system implementation in our hospital in early phase (during June 2009-2011), there was no statistic significance in difference in In-Hospital cardiac arrest incidence and overall hospital mortality rate. Since the introduction of the ERT service in our hospital, we have conducted continuous educational campaign to improve awareness in an attempt to increase use of the service. Methods: To investigate outcome of ERT system in In-Hospital cardiac arrest and overall hospital mortality rate. We conducted a prospective, controlled before-and after examination of the long term effect of a ERT system on the incidence of cardiac arrest. We performed Chi -square analysis to find statistic significance. Results: Of a total 623 ERT cases from June 2009 until December 2012, there were 72 calls in 2009, 196 calls in 2010 ,139 calls in 2011 and 245 calls in 2012. The number of ERT calls per 1000 admissions in year 2009-10 was 7.69, 5.61 in 2011 and 9.38 in 2013. The number of Code blue calls per 1000 admissions decreased significantly from 2.28 to 0.99 per 1000 admissions (P value < 0.001). The incidence of cardiac arrest decreased progressively from 1.19 to 0.34 per 1000 admissions and significant in difference in year 2012 (P value < 0.001). The overall hospital mortality rate decreased by 8 % from 15.43 to 14.43 per 1000 admissions (P value 0.095). Conclusions: ERT system implementation was associated with progressive reduction in cardiac arrests over three year period, especially statistic significant in difference in 4th year after implementation. We also found an inverse association between number of ERT use and the risk of occurrence of cardiac arrests, But we have not found difference in overall hospital mortality rate.

Keywords : emergency response team, ERT, cardiac arrest, emergency medicine

Conference Title : ICEM 2015 : International Conference on Emergency Medicine

Conference Location : Montreal, Canada

Conference Dates : May 11-12, 2015