

## Comparative Acaricidal Efficacy of Fluralaner vs Oral Ivermectin Against Tick Infestation in Dogs

**Authors :** Tayyaba Zahra, Shehla Gul Bokhari, Asim Khalid Mahmood, Raheela Akhtar, Khizar Matloob

**Abstract :** In Pakistan, dogs are commonly infested with ticks, especially in summers, causing not only dermatological issues but also systemic problems. Persistence of tick infestation often leads to heavy losses. Different acaricides are locally available with variable efficacy; however, recurrence of infestation is commonly reported. The present study was thus designed to compare the efficacy of a novel drug Fluralaner and conventionally used Ivermectin against tick infestation. Dogs positive for tick infestation were randomly divided into 2 groups viz, Groups A and B having 8 dogs each. Ticks were enumerated manually from the whole body of dogs at day 0 before the administration of drugs. Dogs in Group A were treated with Fluralaner at day 0, and dogs in Group B were treated with Ivermectin. Post-treatment, ticks were counted again at days 7, 14, 21, 28, and 35. At day 07 of the study, no tick was found on the dogs treated with Fluralaner, while many ticks were present on the dogs treated with Ivermectin showing an efficacy up to 50%. On the consecutive follow-up evaluations, similar results were found for Fluralaner while the efficacy of Ivermectin was further reduced to less than 50%. Furthermore, Fluralaner treated dogs had better RBC counts, PCV, Hgb concentration, LFTs, RFTs post-treatment than the dogs treated with Ivermectin. Statistically, oral Fluralaner proved a more effective drug ( $P \leq 0.05$ ) than oral Ivermectin against tick infestation in dogs.

**Keywords :** fluralaner, ivermectin, dogs, tick infestations

**Conference Title :** ICABVM 2021 : International Conference on Animal Breeding and Veterinary Medicine

**Conference Location :** Istanbul, Türkiye

**Conference Dates :** December 20-21, 2021