Bioassay Guided Isolation of Antimicrobial and Cytotoxic Components from Ethyl Acetate Extracts of Cassia Sieberiana D.C. (Fabaceae)

Authors: Abubakar Sani

Abstract : The leaves extracts of Cassia sieberiana D.C. were screened for cytotoxicity using Brine Shrimp Test (BST) and antimicrobial bioassay against Staphylococcus aureus, Salmonella typhi and Escherichia coli using crude ethanol extract, Chloroform soluble fraction, aqueous soluble fraction, ethyl acetate soluble fraction, methanol soluble fraction and n-hexane soluble fraction. The Ethyl acetate fraction obtained proved to be most active in inducing complete lethality at minimum doses in BST and also active on Salmonella typhi. The Bioactivity result was used to guide the column chromatography which led to the isolation of pure compound CSB-8 which was found active in the BST with LC₅₀ value of 34(722-182)μg/ml and showed remarkable activity on Salmonella typhi (zone of inhibition 25mm) at 10,000μg/ml. The ¹H-NMR, ¹³C NMR, FTIR and GC-MS spectra of compound suggested the proposed structure to be 2-pentadecanone.

Keywords: brine shrimp, Cassia sieberiana D. C, Column chromatography, antimicrobial bioassay

Conference Title: ICP 2021: International Conference on Phytochemistry

Conference Location: Istanbul, Türkiye Conference Dates: December 20-21, 2021