Cytotoxic Activity Of Major Iridoids From Barleria Trispinosa (Forssk.) Vahl. Growing In Saudi Arabia

Authors : Hamza Assiry, Gamal A. Mohamed, Sabrin R. M. Ibrahim, Hossam M. Abdallah

Abstract : Chemical investigation of the aerial parts of Barleria trispinosa(Forssk.) Vahl. resulted in isolation of four major iridoids that were identified as 6,8-O,O-diacetylshanhiside methyl ester (acetyl barlerin) (1), 8-O-acetylshanzhiside methyl ester (barlerin) (2), shanzhiside methyl ester (3), and $6 - \Box$ -L-rhamnopyranosyl-8-O-acetylshanzhiside methyl ester (4). The isolated compounds were confirmed by detailed one and two-dimensional NMR. Isolated compounds were tested for their cytotoxic activity on breast cancer (MCF-7, MDA-MB-231) and colon cancer (LS174T) cell linesusing sulphorhodamine B (SRB) assay. It is noteworthy that compound 1 demonstrated a significant cytotoxic potential towards MDA-MB-231 cell line with IC5016.7 \pm 2.7µg / mL compared to doxorubicin whereas compounds 2, showed moderate cytotoxic potential with IC5021.2 \pm 1.9µg / mL on MCF-7. The other compounds showed moderate activity on the tested cell lines.

Keywords : acanthaceae, cytotoxicity, metabolites, barleria trispinosa

Conference Title : ICMPPPNP 2022 : International Conference on Medicinal Plants, Pharmacognosy, Phytochemistry and Natural Products

1

Conference Location : Rome, Italy

Conference Dates : March 03-04, 2022