

## Gestational Diabetes Mellitus (GDM) Increasing Postpartum Screening to Prevent T2D

**Authors :** Boma Nellie S, Nambiar Ritu, K. Kanchanmala, T. Rashida, Israell Imelda, Moul Khusnud, Michael Marina

**Abstract :** Gestational diabetes (GDM) imparts an increased life long risk of developing Type 2 Diabetes Mellitus (T2DM) and cardiovascular disease in women. Once diagnosed with GDM women have up to 74% increased cumulative risk developing T2DM in 10-15 years. Identifying women at increased risk of developing T2DM and offering them pharmacological and lifestyle management interventions will delay or eliminate the development of diabetes in this population. While ADA recommends that all gestational diabetics be offered postnatal screening, worldwide the screening rates from 35-75% and Al Rahba Hospital with a robust universal antenatal screening program for GDM was at a dismal 9% in 2011. A multidisciplinary team was put together involving OB/Gyn Physicians, Midwives, Nurses (ward and OPD) Diabetic Educators, Dietitians, Medical Records, Laboratory & IT with the implementation of multiple strategies to increase the uptake of postpartum screening of the gestational diabetic.

**Keywords :** GDM, postnatal screening, preventing type 2 diabetes, lifestyle management

**Conference Title :** ICOG 2014 : International Conference on Obstetrics and Gynaecology

**Conference Location :** Bangkok, Thailand

**Conference Dates :** December 18-19, 2014