

Honey Dressing versus Silver Sulfadiazine Dressing for Wound Healing in Second Degree Thermal Burn Patients

Authors : Syed Faizan Hassan Shah

Abstract : Introduction: Burn injuries are among the most devastating of all injuries. Burns is the fourth most common type of trauma worldwide. Approximately 90 percent of burns occur in low to middle-income countries. Nearly half a million Americans each year, with approximately 40,000 hospitalizations and 3,400 deaths annually, suffer burns. The survival rate for admitted burn patients has improved consistently over the past four decades, largely attributed to national decreases in burn size, improvements in burn critical care, and advancements in burn wound care. Objectives: The present study was conducted to compare the efficacy of Honey dressing versus Silver Sulfadiazine dressing for complete wound healing in the 2nd-degree thermal burn. Study Design: A Randomized controlled trial was carried out in the Department of General Surgery/burn unit of Ayub Teaching Hospital Abbottabad from July to December 2018. The study population included thermal burn patients presenting with ASA-I, ASA-II, and body surface area less than 50% of the age group above 12 to 60 years of either gender. All the patients were randomly divided into two equal groups of patients by blocked randomization using permuted block of 6. In group 'A,' patients underwent dressing by honey method, and patients in group 'B' had silver sulfadiazine dressing. The dressing was changed every 48 hours by a senior surgeon, and the condition of the wound was observed. Time duration till complete wound healing was noted in the Proforma. Results: A total of 100 patients were selected and divided into two groups of 50 patients in each two groups. The mean age of the patients was 27.66 ± 13.388 ranging from 12 to 60 years of age, and the mean duration of complete healing of wound in days was 20.20 ± 6.251 , ranging from 2 to 30 days. Mean comparison of age with both groups, age of the patients was 21.24 ± 3.761 (n=50) in group 'A,' i.e., honey dressing, and 19.16 ± 7.911 (n=50) was in group 'B,' i.e., silver sulfadiazine dressing. Efficacy in the honey dressing group was found effective in 48(75.0%) and ineffective in 2(5.6%) out of 50 patients. Efficacy in silver sulfadiazine dressing group 16(25.0%) was three found effective and in 34(94.4%) was ineffective out of 50 patients. There was a statistically significant difference between both groups. (P=0.000) . Conclusion: honey dressing is more effective as compared to silver sulfadiazine dressing in terms of complete wound healing in second-degree thermal burn patients; our study also concluded the same.

Keywords : efficacy, honey dressing, silver sulfadiazine dressing, wound healing

Conference Title : ICSAT 2022 : International Conference on Surgery, Anesthesiology and Trauma

Conference Location : London, United Kingdom

Conference Dates : September 22-23, 2022