

Gold Nanoprobes Assay for the Identification of Foodborn Pathogens Such as *Staphylococcus aureus*, *Listeria monocytogenes* and *Salmonella enteritis*

Authors : D. P. Houhoula, J. Papaparaskevas, S. Konteles, A. Dargenta, A. Farka, C. Spyrou, M. Ziaka, S. Koussisis, E. Charvalos

Abstract : Objectives: Nanotechnology is providing revolutionary opportunities for the rapid and simple diagnosis of many infectious diseases. *Staphylococcus aureus*, *Listeria monocytogenes* and *Salmonella enteritis* are important human pathogens. Diagnostic assays for bacterial culture and identification are time consuming and laborious. There is an urgent need to develop rapid, sensitive, and inexpensive diagnostic tests. In this study, a gold nanoprobe strategy developed and relies on the colorimetric differentiation of specific DNA sequences based approach on differential aggregation profiles in the presence or absence of specific target hybridization. Method: Gold nanoparticles (AuNPs) were purchased from Nanopartz. They were conjugated with thiolated oligonucleotides specific for the *femA* gene for the identification of members of *Staphylococcus aureus*, the *mecA* gene for the differentiation of *Staphylococcus aureus* and MRSA *Staphylococcus aureus*, *hly* gene encoding the pore-forming cytolysin listeriolysin for the identification of *Listeria monocytogenes* and the *invA* sequence for the identification of *Salmonella enteritis*. DNA isolation from *Staphylococcus aureus* *Listeria monocytogenes* and *Salmonella enteritis* cultures was performed using the commercial kit Nucleospin Tissue (Macherey Nagel). Specifically 20 μ l of DNA was diluted in 10mMPBS (pH5). After the denaturation of 10min, 20 μ l of AuNPs was added followed by the annealing step at 58oC. The presence of a complementary target prevents aggregation with the addition of acid and the solution remains pink, whereas in the opposite event it turns to purple. The color could be detected visually and it was confirmed with an absorption spectrum. Results: Specifically, 0.123 μ g/ μ l DNA of *St. aureus*, *L.monocytogenes* and *Salmonella enteritis* was serially diluted from 1:10 to 1:100. Blanks containing PBS buffer instead of DNA were used. The application of the proposed method on isolated bacteria produced positive results with all the species of *St. aureus* and *L. monocytogenes* and *Salmonella enteritis* using the *femA*, *mecA*, *hly* and *invA* genes respectively. The minimum detection limit of the assay was defined at 0.2 ng/ μ L of DNA. Below of 0.2 ng/ μ L of bacterial DNA the solution turned purple after addition of HCl, defining the minimum detection limit of the assay. None of the blank samples was positive. The specificity was 100%. The application of the proposed method produced exactly the same results every time (n = 4) the evaluation was repeated (100% repeatability) using the *femA*, *hly* and *invA* genes. Using the gene *mecA* for the differentiation of *Staphylococcus aureus* and MRSA *Staphylococcus aureus* the method had a repeatability 50%. Conclusion: The proposed method could be used as a highly specific and sensitive screening tool for the detection and differentiation of *Staphylococcus aureus* *Listeria monocytogenes* and *Salmonella enteritis*. The use AuNPs for the colorimetric detection of DNA targets represents an inexpensive and easy-to-perform alternative to common molecular assays. The technology described here, may develop into a platform that could accommodate detection of many bacterial species.

Keywords : gold nanoparticles, pathogens, nanotechnology, bacteria

Conference Title : ICSRD 2020 : International Conference on Scientific Research and Development

Conference Location : Chicago, United States

Conference Dates : December 12-13, 2020