## Effectiveness of Myofascial Release Technique in Treatment of Sacroiliac Joint Hypo-Mobility in Postnatal Women

Authors : Ahmed A. Abd El Rahim, Mohamed M. M. Essa, Magdy M. A. Shabana, Said A. Mohamed, Mohamed Ibrahim Mabrouk

**Abstract :** Background: Sacroiliac joint (SIJ) dysfunction is considered the main cause of pregnancy-related back pain, which may continue to persist postnatally. Myofascial release technique (MFR) is an application of low-intensity, prolonged stretch to myofascial structures to improve function by increasing the sliding properties of restricted myofascial tissues. Purpose: This study was designed to investigate the effect of MFR on postnatal SIJ hypo-mobility. Materials and Methods: Fifty postnatal women complaining of SIJ hypo-mobility participated in this study. Their ages ranged from 26 to 35 yrs., and their body mass index (BMI) didn't exceed 30 kg/m2. They were randomly assigned to two equal groups, group A (Gr. A) and group B (Gr. B). Both groups received three sessions per week for eight successive weeks. Gr. A received a traditional physical therapy program, while Gr. B received a traditional physical therapy program in addition to MFR. Doppler imaging of vibration was utilized to measure SIJ mobility pre- and post-intervention, and an electronic digital goniometer was used to measure back flexion and extension Range of motion. Results: Findings revealed a statistical improvement in post-intervention values of SIJ mobility in addition to trunk flexion and extension ROM in Gr. B compared to Gr. A (P<0.001). Conclusion: Adding MFR to traditional physical therapy programs is highly recommended in the treatment of SIJ hypo-mobility in postnatal women.

Keywords : sacroiliac hypo-mobility, sacroiliac dysfunction, myofascial release technique, traditional physical therapy, postnatal

**Conference Title :** ICPTT 2023 : International Conference on Physical Therapy Types **Conference Location :** New York, United States **Conference Dates :** August 10-11, 2023