

Case Report: Treatment Resistant Schizophrenia in an Immigrant Adolescent

Authors : Omaymah Al-Otoom, Rajesh Mehta

Abstract : Introduction: Migration is an established risk factor in the development of schizophrenia and other forms of psychosis. The exposure to different social adversities, including social isolation, discrimination, and economic stress, is thought to contribute to elevated rates of psychosis in immigrants and their children. We present a case of resistant schizophrenia in an immigrant adolescent. Case: The patient is a 15-year-old male immigrant. In October 2021, the patient was admitted for irritability, suicidal ideations, and hallucinations. He was treated with Fluoxetine 10 mg daily for irritability. In November 2021, he presented with similar manifestations. Fluoxetine was discontinued, and Risperidone 1 mg at bedtime was started for psychotic symptoms. In March 2022, he presented with commanding auditory hallucinations (voices telling him that people were going to kill his father). Risperidone was gradually increased to 2.5 mg twice daily for hallucinations. The outpatient provider discontinued Risperidone and started Olanzapine 7.5 mg and Lurasidone 40 mg daily. In August 2022, he presented with worsening paranoia due to medication non-adherence. The patient had limited improvement on medications. In October 2022, the patient presented to the ED for visual hallucinations and aggression towards the family. His medications were Olanzapine 10 mg daily, Lurasidone 60 mg daily, and Haloperidol 2.5 mg twice daily. In the ED, he received multiple as-needed medications and was placed in seclusion for his aggressive behavior. The patient showed a positive response to a higher dose of Olanzapine and decreased dose of Lurasidone. The patient was discharged home in stable condition. Two days after discharge, he was brought for bizarre behavior, visual hallucinations, and homicidal ideations at school. Due to concerns for potential antipsychotic side effects and poor response, Lurasidone and Olanzapine were discontinued, and he was discharged home on Haloperidol 5 mg in the morning and 15 mg in the evening. Clozapine treatment was recommended on an outpatient basis. He has no family history of psychotic disorders. He has no history of substance use. A medical workup was done, the electroencephalogram was normal, and the urine toxicology was negative. Discussion: Our patient was on three antipsychotics at some point with no improvement in his psychotic symptoms, which qualifies as treatment-resistant schizophrenia (TRP). It is well recognized that migrants are at higher risk of different psychiatric disorders, including posttraumatic stress disorder, affective disorders, schizophrenia, and psychosis. This is thought to be related to higher exposure to traumatic life events compared to the general population. In addition, migrants are more likely to experience poverty, separation from family members, and discrimination which could contribute to mental health issues. In one study, they found that people who migrated before the age of 18 had twice the risk of psychotic disorders compared to the native-born population. It is unclear whether migration increases the risk of treatment resistance. In a Canadian study, neither ethnicity nor migrant status was associated with treatment resistance; however, this study was limited by its small sample size. There is a need to implement psychiatric prevention strategies and outreach programs through research to mitigate the risk of mental health disorders among immigrants.

Keywords : psychosis, immigrant, adolescent, treatment resistant schizophrenia

Conference Title : ICPMHD 2023 : International Conference on Psychiatry and Mental Health Disorders

Conference Location : New York, United States

Conference Dates : January 30-31, 2023