Etiological Factors for Renal Cell Carcinoma: Five-Year Study at Mayo Hospital Lahore

Authors : Muhammad Umar Hassan

Abstract : Renal cell carcinoma is a subset of kidney cancer that arises in the lining of DCT and is present in parenchymal tissue. Diagnosis is based on lab reports, including urinalysis, renal function tests (RFTs), and electrolyte balance, along with imaging techniques. Organ failure and other complications have been commonly observed in these cases. Over the years, the presentation of patients has varied, so carcinoma was classified on the basis of site, shape, and consistency for detailed analysis. Lifestyle patterns and occupational history were inquired about and recorded. Methods: Data from 100 patients presenting to the oncology and nephrology department of Mayo Hospital in the year 2015-2020 were included in this retrospective study on a random basis. The study was specifically focused on three risk factors. Smoking, occupational exposures, and Hakim medicine are taken by the patient for any cause. After procurement of data, follow-up contacts of these patients were established, resulting in a detailed analysis of lifestyle. Conclusion: The inference drawn is a direct causal link between smoking, industrial workplace exposure, and Hakim medicine with the development of Renal Cell Carcinoma. It was shown in the majority of the patients and hence confirmed our hypothesis.

Keywords : renal cell carcinoma, kidney cancer, clear cell carcinoma

Conference Title : ICOMO 2023 : International Conference on Oncogenesis and Molecular Oncology

Conference Location : London, United Kingdom

Conference Dates : February 16-17, 2023

1

ISNI:000000091950263