## Inf-y and Il-2 Asses the Therapeutic Response in Anti-tuberculosis Patients at Jamot Hospital Yaounde, Cameroon

Authors : Alexandra Emmanuelle Membangbi, Jacky Njiki Bikoï, Esther Del-florence Moni Ndedi, Marie Joseph Nkodo Mindimi, Donatien Serge Mbaga, Elsa Nguiffo Makue, André Chris Mikangue Mbongue, Martha Mesembe, George Ikomey Mondinde, Eric Walter Perfura-yone, Sara Honorine Riwom Essama

Abstract : Background: Tuberculosis (TB) is one of the top lethal infectious diseases worldwide. In recent years, interferon-y (INF-y) release assays (IGRAs) have been established as routine tests for diagnosing TB infection. However, produced INF-y assessment failed to distinguish active TB (ATB) from latent TB infection (LTBI), especially in TB epidemic areas. In addition to IFN-y, interleukin-2 (IL-2), another cytokine secreted by activated T cells, is also involved in immune response against Mycobacterium tuberculosis. The aim of the study was to assess the capacity of IFN- $\gamma$  and IL2 to evaluate the therapeutic response of patients on anti-tuberculosis treatment. Material and Methods: We conducted a cross-sectional study in the Pneumonology Departments of the Jamot Hospital in Yaoundé between May and August 2021. After signed the informed consent, the sociodemographic data, as well as 5 mL of blood, were collected in the crook of the elbow of each participant. Sixty-one subjects were selected (n = 61) and divided into 4 groups as followed: group 1: resistant tuberculosis (n = 13), group 2: active tuberculosis (n=19), group 3 cured tuberculosis (n=16), and group 4: presumed healthy persons (n=13). The cytokines of interest were determined using an indirect Enzyme-linked Immuno-Sorbent Assay (ELISA) according to the manufacturer's recommendations. P-values < 0.05 were interpreted as statistically significant. All statistical calculations were performed using SPSS version 22.0 Results: The results showed that men were more 14/61 infected (31,8%) with a high presence in active and resistant TB groups. The mean age was 41.3±13.1 years with a 95% CI = [38.2-44.7], the age group with the highest infection rate was ranged between 31 and 40 years. The IL-2 and INF- $\gamma$  means were respectively 327.6±160.6 pg/mL and 26.6±13.0 pg/mL in active tuberculosis patients, 251.1±30.9 pg/mL and 21.4±9.2 pg/mL in patients with resistant tuberculosis, while it was 149.3±93.3 pg/mL and 17.9±9.4 pg/mL in cured patients, 15.1±8.4 pg/mL and 5.3±2.6 pg/mL in participants presumed healthy (p <0.0001). Significant differences in IFN- $\gamma$  and IL-2 rates were observed between the different groups. Conclusion: Monitoring the serum levels of INF-y and IL-2 would be useful to evaluate the therapeutic response of anti-tuberculosis patients, particularly in the both cytokines association case, that could improve the accuracy of routine examinations. **Keywords** : antibiotic therapy, interferon gamma, interleukin 2, tuberculosis

**Conference Title :** ICIADCI 2023 : International Conference on Immunological Aspects of Diseases and Clinical Immunology **Conference Location :** Prague, Czechia

1

Conference Dates : September 04-05, 2023