Osteitis in the Diabetic Foot in Algeria

Authors : Mohamed Amine Adaour, Mohamed Sadek Bachene, Mosaab Fortassi, Wafaa Siouda

Abstract : — Foot infections are responsible for a significant number of hospitalizations and amputations in diabetic patients. The objective of our study is to analyze and evaluate the management of diabetic foot in a surgical setting. A retrospective study was conducted based on a selected case of suspected diabetic foot infections of osteitis treated at the Mohamed Boudiaf hospital in Medea. The case was reiterated as a therapeutic charge, consisting of treating first the infection of the soft tissues, then the osteitis: biopsy after at least 15 days of cessation of antibiotic therapy. Successful treatment of osteitis was defined at the end of a follow-up period of complete wound healing, lack of bone resection/amputation surgery at the initial bone site during follow-up , Instead, biopsies are prescribed in the treatment of soft tissue infection. The mean duration of treatment for soft tissue infection was 2-3 weeks, the duration of the antibiotic-free window of therapy prior to bone biopsy was 2-4 weeks. This patient received medical management without surgical resection. The success rate for treating osteitis at one year was 73%, and healing at one year was 88%. It is often limited to a sausage of the foot at the cost of repeated amputations. The best management remains prevention, which necessarily involves setting up a specialized and adapted centre.

Keywords : diabetic foot, bone biopsy, osteitis, algeria

Conference Title : ICGDDM 2023 : International Conference on General Diabetes and Diabetes Management

Conference Location : London, United Kingdom **Conference Dates :** December 11-12, 2023

Open Science Index, Medical and Health Sciences Vol:17, No:12, 2023 publications.waset.org/abstracts/167195.pdf