

Study of the Prevalence, Associated Factors and Impact of Maternal Perinatal Depression in Women Alexandria 2022

Authors : Nermeen Saad Elbeltagy, Hoda Ghareeb, Hesham Adel Elsheshtawy, Nadim Hamed, Amany Ibrahim Mostafa, Sara Hazem Hassan

Abstract : Introduction: Depression is one of the most common mental health problems occurring in women during their child bearing years. Perinatal depression refers to major and minor depressive episodes that occur either during pregnancy or aer delivery. Although perinatal depression is common in developing countries, it is under-recognized in low and middle income countries making a substantial contribution to maternal and infant morbidity and mortality. About 12.5 - 42% of pregnant women and, 12 - 50% of post natal mothers in low and middle income countries such as Ethiopia had depression AIM OF THE WORK: To study prevalence, associated factors and impact of maternal perinatal depression in Alexandria. Patients and method: This study was conducted on 300 mothers at the postnatal ward in ElShatby Maternity Hospital from April 2022 unl October 2022. Females with past history of depression before pregnancy or females who receive medications inducing depression were excluded. The participants were asked to complete the questionnaire that includes the Edinburgh Postnatal Depression Scale (EPDS) as a screening test to obtain information concerning the current frame of mind at antepartum, partum and postpartum periods Results: The prevalence of perinatal depression was 22.3%. It was found that there is a significant negave moderate correlation between socioeconomic status and perinatal depression($r=-0.42$). The present study revealed that about two thirds (60.7%) of postpartum women had low socioeconomic level. Also, less than one fourth (20%) of parents had high education and only one fourth (25.3%) of postpartum women were working. There was a statically significance difference between the number of previous abortions and perinatal depression ($p=0.04$). There was a significant moderate correlation between the amount of blood lost during delivery and an increased risk of developing postpartum depression. The prevalence of perinatal depression was high in cases of female neonates more than male ones. Conclusion: the prevalence of perinatal depression among the studied women was 22.3% of studied group. The significant factors identified in this study can be targeted to reduce the occurrence of perinatal depression among pregnant women in Alexandria through appropriate health interventions which includes perinatal depression screening, counseling, and the provision of support for pregnant women during antenatal care as well as lifestyle modification.

Keywords : mental health, depression in pregnancy, mental disorders, psychology in pregnancy

Conference Title : ICHECM 2023 : International Conference on Healthcare Emergency and Crisis Management

Conference Location : Berlin, Germany

Conference Dates : July 17-18, 2023