

Effect of Different Weed Management Strategies in Chickpea Yield

Authors : Ijaz Ahmed Khan, Zaheen Ullah, Rahamdad, Gul Hassan

Abstract : A field experiment was conducted at Agricultural Research Station Ahmad Wala, Karak, Khyber Pakhtunkhwa Province during rabi season of 2010-011 to study the effect of different weed management practices on weed control in chickpea under field conditions. The results revealed that treatments showed significant influence on weed density, seed yield kg ha⁻¹ and other growth parameters. Significantly lower weed density (98 m⁻²) was recorded with the application of Isoproturon 500 EW as compared to control plots having 368.3 weeds m⁻². Moreover, significantly highest seed yield (1583.3 kg ha⁻¹) was produced in the plots assigned with Isoproturon 500 EW followed by Eucalyptus extract that produce seed yield of 1416.7 kg ha⁻¹. It was concluded from the study that Isoproturon 500 EW is the best option for controlling weeds and increase the seed yield kg ha⁻¹ of chickpea.

Keywords : chickpea, herbicides, weed control, weeds extracts

Conference Title : ICAB 2015 : International Conference on Agriculture and Biotechnology

Conference Location : Jeddah, Saudi Arabia

Conference Dates : January 26-27, 2015