

Characteristics of Clinical and Diagnostic Aspects of Benign Diseases of Cervix in Women

Authors : Gurbanova J., Majidova N., Ali-Zade S., Hasanova A., Mikailzade P.

Abstract : Currently, the problem of oncogynecological diseases is widespread and remains relevant in terms of quantitative growth. It is known that due to the increase in the number of benign diseases of the cervix, the development of precancerous conditions occurs. Benign diseases of the cervix represent the most common gynecological problem, which are often precursors of malignant neoplasms, especially cervical cancer. According to statistics, benign diseases of the cervix cover 25-45% of all gynecological diseases. Among women's oncogynecological diseases, cervical cancer ranks second in the world after breast cancer and ranks first in the mortality rate among oncological diseases in economically underdeveloped countries. We performed a comprehensive clinical and laboratory examination of 130 women aged 18 to 73 with benign cervical diseases. 59 (38.5%) women of reproductive age, as well as 39 (30%) premenopausal and 41 (31.5%) menopausal patients, participated in the study. Detailed anamnesis was collected from all patients, objective and gynecological examination was performed, laboratory and instrumental examinations (USM, IPV DNA, smear microscopy, and PCR bacteriological examination of sexually transmitted infections), simple and extended colposcopy, liquid-based PAP-smear smear and PAP-classic smear examinations were conducted. As a result of the research, the following nosological forms were found in women with benign diseases of the cervix: non-specific vaginitis in 10 (7.7%) cases; ectopia, endocervicitis - 60(46.2%); cervical ectropion - 7(5.4%); cervical polyp - 9(6.9%); cervical leukoplakia - 15(11.5%); atrophic vaginitis - 7(5.4%); condyloma - 12(9.2%); cervical stenosis - 2(1.5%); endometriosis of the cervix - was noted in 8 (6.2%) cases ($p < 0.001$), respectively. Characteristics of the menstrual cycle among the examined women: normal cycle in 97 (74.6%) cases; oligomenorrhea - 23 (17.7%); polymenorrhea - 4(3.1%); algomenorrhea - noted in 6 (4.6%) cases ($p < 0.001$). Cytological examination showed that: the specificity of liquid-based cytology was 76.2%, and the traditional PAP test was set at 70.6%. The overall diagnostic value was calculated to be 86% in liquid-based cytology and 78.5% in conventional PAP tests. Treatment of women with benign diseases of the cervix was carried out by diathermocoagulation method and "FOTEK EA 141M" device. It should be noted that 6 months after the treatment, after treatment with the "FOTEK EA 141M" device, there was no relapse in any patient. Recurrence was found in 23.7% of patients after diathermoelectrocoagulation. Thus, it is clear from the above that the study of cervical pathologies, the determination of optimal examinations, and effective treatment methods is one of the urgent problems facing obstetrics and gynecology.

Keywords : cervical cancer, cytological examination, PAP-smear, non-specific vaginitis

Conference Title : ICWGCRM 2023 : International Conference on Women Gynecology, Childbirth and Reproductive Medicine

Conference Location : Dubai, United Arab Emirates

Conference Dates : November 13-14, 2023