## Prevalence, Awareness, and Risk Factors of Diabetes in Ahvaz: South West of Iran

Authors: Leila Yazdanpanah, Hajieh Shahbazian, Seyed Mahmoud Latifi, Armaghan Moravej Aleali, Saeed Ghanbari

Abstract: Introduction: This study was designed to determine the prevalence of diabetes in people aged over 20 years in Ahvaz, Iran. Material and Methods: The study population selected by cluster sampling. Fasting blood sugar (FBS) assessed after minimum 8 hours night fasting. A questionnaire included: age, sex, weight, height, blood pressure, waist circumference and previous history of diabetes were completed for each patient. FBS≥126mg/dl and/or oral hypoglycemic treatment and/or insulin was defined as diabetes, FBS=100-125 mg/dl as impaired fasting glucose (IFG) and FBS<100mg/dl as normal. Results: Study population was 936 persons (47.2 % male and 52.8% female). The mean age of a population was 42.2±14 years. Diabetes was detected in 15.1 % of population. Only 57cases(6.1%) were aware of their disease and 9% had unknown diabetes. Diabetes was detected in 14.5% of male (11.3% unknown and 3.2 % known diabetes) and in 11.7% of female (7% unknown and 4.7% known diabetes). Prevalence of diabetes had no significant difference (P=0.21) in male and female but unknown diabetes was significantly higher in male (P=0.025). Prevalence of diabetes was increased with rising of age between 20-60 years old but decreasing after 60 years old. Diabetes was related to age, waist circumference and systolic and diastolic blood pressure, TG level and BMI in both sex (P=0.0001). Conclusion: More than half of female and three-fourth of male diabetic patients are unaware of their disease in South of Iran. Diabetes screening should be intensified in this population.

**Keywords:** diabetes, prevalence, risk factor, awareness

Conference Title: ICDM 2015: International Conference on Diabetes and Metabolism

**Conference Location :** London, United Kingdom **Conference Dates :** February 16-17, 2015