The Prevalence of Coronary Artery Disease and Its Risk Factors in Rural and Urban Areas of Pakistan

Authors : Muhammad Kamran Hanif Khan, Fahad Mushtaq

Abstract : Background: In both developed and underdeveloped countries, coronary artery disease (CAD) is a serious cause of death and disability. Cardiovascular disease (CVD) is becoming more prevalent in emerging countries like Pakistan due to the spread and acceptance of Western lifestyles. Material and Methods: An observational cross-sectional investigation was conducted, and data collection relied on a random cluster sampling method. The sample size for this cross-sectional study was calculated using the following factors: estimated true proportion of 17.5%, desired precision of 2%, and confidence interval of 95%. The data for this study was collected from a sample of 1387 adults. Results: The average age of those living in rural areas is 55.24 years, compared to 52.60 years for those living in urban areas. The mean fasting blood glucose of the urban participants is 105.28 mg/dL, which is higher than the mean fasting blood glucose of the rural participants, which is 191.97 mg/dL. CAD prevalence is greater in urban areas than in rural areas. ECG abnormalities prevalence is 16.1% in females compared to 12.5% in men. Conclusion: The prevalence of CAD is more common in urban areas than in rural ones for all of the measures of CAD used in the study.

Keywords : CVD prevalence, CVD risk factors, rural area, urban area

Conference Title : ICEMN 2024 : International Conference on Emergency Medicine and Public Health

Conference Location : Florence, Italy

Conference Dates : May 16-17, 2024

1