

Response of Subfossile Diatoms, Cladocera and Chironomidae in Sediments of Small Ponds to Changes in Wastewater Discharges from a Zn-Pb Mine

Authors : Ewa Szarek-Gwiazda, Agata Z. Wojtal, Agnieszka Pocięcha, Andrzej Kownacki, Dariusz Ciszewski

Abstract : Mining of metal ores is one of the largest sources of heavy metals, which deteriorate aquatic systems. The response of organisms to environmental changes can be well recorded in sediments of the affected water bodies and may be reconstructed basing on analyses of organisms remains. The present study aimed at response of diatoms, Cladocera and Chironomidae communities to impact of Zn-Pb mine water discharge recorded in sediment cores of small subsidence ponds on the Chechło River floodplain (Silesia-Krakow Region, southern Poland). We hypothesis various response of the above groups to high metal concentrations (Cd, Pb, Zn, and Cu). The investigated ponds were formed either during the peak of the ore exploitation (DOWN) or after mining cessation (UP). Currently, the concentrations of dissolved metals (in $\mu\text{g g}^{-1}$) in water reached up to 0.53 for Cd, 7.3 for Pb and up to 47.1 for Zn. All the sediment cores from subsidence ponds were heavily polluted with Cd 6.7–612 $\mu\text{g g}^{-1}$, Pb 0.1–10.2 mg g^{-1} , and Zn 0.5–23.1 mg g^{-1} . Core sediments varied also in respect to pH 5.8–7.1 and concentrations of organic matter (5.7–39.8%). The impact of high metal concentrations was expressed by the occurrence of metal tolerant taxa like diatoms – *Nitzschia amphibia*, *Sellaphora nigri*, and *Surirella brebisonii* var. *kuetzingii*; Cladocera – *Chydorus sphaericus* (dominated in cores from all ponds), and Chironomidae – *Chironomus* and *Cricotopus* especially in the DOWN ponds. Statistical analysis exhibited a negative impact of metals on some taxa of diatoms and Cladocera, but only on *Polypedilum* sp. from Chironomidae. Abundance of such diatoms like *Gomphonema utae*, *Staurosirella pinnata*, *Eunotia bilunaris*, and Cladocera like *Alona*, *Chydorus*, *Graptoleberis*, and *Pleuroxus* decreased with increasing Pb concentration. However, the occurrence or dominance of more sensitive species of diatoms and Cladocera indicates their adaptation to higher metal loads, which was facilitated by neutral pH and slightly alkaline waters. Diatom assemblages were generally resistant to Zn, Pb, Cu, and Cd pollution as indicated by their large similarity to populations from non-contaminated waters. Comparison with reference objects clearly indicates the dominance of *Achnanthes minutissimum*, *Staurosira venter*, and *Fragilaria gracilis* in very diverse assemblages of unpolluted waters. The distribution of the Cladocera and Chironomidae taxa depended on the habitat type. The DOWN ponds with stagnant water and overgrown with macrophytes were more suitable for Cladocerans (14 taxa, higher diversity) than the UP ponds with river water flowing through their centre and with a small share of macrophytes (8 taxa). The Chironominae, mainly *Chironomus* and *Microspectra*, were abundant in cores from the UP ponds with muddy bottom. Inversely, the density of Orthocladinae, especially genus *Cricotopus*, were related to the organic matter content and dominated in cores from the DOWN ponds. The presence of diatoms like *Nitzschia amphibia*, *Sellaphora nigri*, and *Surirella brebisonii* var. *kuetzingii*, Cladocerans *Bosmina longirostris*, *Chydorus sphaericus*, littoral cladocerans, *Alona affinis*, *A. rectangularis* as well as Chironomidae *Chironomus* sp. (UP ponds) and *Psectrotanypus varius* (DOWN ponds) indicate the influence of the water trophy on their distribution.

Keywords : Chironomidae, cladocerans, diatoms, metals, Zn-Pb mine, sediment cores, subsidence ponds

Conference Title : ICHMEE 2024 : International Conference on Heavy Metals in the Environment and Ecosystems

Conference Location : Rome, Italy

Conference Dates : August 22-23, 2024