

Factors Affecting the Fear of Insulin Injection and Finger Punching in Individuals Diagnosed with Diabetes

Authors : Gaye Demirtaş Adli

Abstract : Research: It was conducted to determine the factors affecting the fear of self-injection and self-pricking of fingers of diabetic individuals. The study was conducted as a cross-sectional, relation-seeking, and descriptive study. The study was conducted on 122 patients who had just started insulin therapy. Data were obtained through The Descriptive Patient Form, The Diabetic Self-Injection, and the Fear of Testing Questionnaire Form (D-FISQ). Descriptive statistical methods used in the evaluation of data are the Mann-Whitney U test, Kruskal-Wallis H test, and the Spearman correlation. The factors affecting the scale scores were evaluated with multiple linear regression analysis. The value of $P < 0.05$ was considered statistically significant. Study group: 56.6% of the patients are male patients. Fear of self-injection (injection), fear of self-testing (test), and total fear (total) scores of women were found to be statistically higher than men ($p < 0.001$). Age, gender, and pain experience were important variables that affected patients' fear of injections. With a one-unit increase in age, the injection fear score decreased by 0.13 points, and the mean injection fear score of women was 2.11 points higher than that of men. It was determined that the patient's age, gender, living with whom, and blood donation history were important variables affecting the fear of self-testing. It is seen that the fear test score decreases by 0.18 points with an increase in age by one unit, and the fear test scores of women compared to men are on average 3,358 points, the fear test scores of those living alone are 4,711 points compared to those living with family members, and the fear test scores of those who do not donate blood are 2,572 compared to those who donate blood score, it was determined that those with more pain experience were 3,156 points higher on average than those with less fear of injection. As a result, it was seen that the most important factors affecting the fear of insulin injection and finger punching in individuals with diabetes were age, gender, pain experience, living with whom, and blood donation history.

Keywords : diabetes, needle phobia, fear of injection, insulin injection

Conference Title : ICN 2024 : International Conference on Nursing

Conference Location : Miami, United States

Conference Dates : March 11-12, 2024