

## Effect of Moringa (*Moringa oleifera* LAM) Leaves Extract on Physicochemical and Organoleptic Properties of Fullfat and Lowfat Yoghurt

**Authors :** B. F. Muhammad, A. M. Abdulqadeer

**Abstract :** The current study determined the effect of fortification using Moringa (*Moringa oleifera*) Leaves Extract (MLE) at different inclusion levels (0, 6, 8, and 10% v/v) on physicochemical and sensory properties of fullfat (FFY) and lowfat (LFY) yoghurt. The results revealed significantly higher protein ( $p < 0.01$ ), fat ( $p < 0.001$ ) and pH in FFY compared to LFY. The MLE inclusion significantly ( $p < 0.001$ ) increased fat contents of the yoghurt types. The pH of LFY produced with 6% MLE was significantly the lowest ( $p < 0.001$ ). The concentration of Mg (21.0 mg/100g), Na (63.0 mg/100g), Ca (173.0 mg/100g), P (416.7 mg/100g), Cu (0.59 mg/100g), Co (0.30 mg/100g), Fe (1.13 mg/100g), and Mn (0.059 mg/100g) were significantly ( $p < 0.001$ ) higher in 10% MLE inclusion level for both FFY and LFY. Also, Mg, Na, Cu and Fe showed significant ( $p < 0.001$ ) negative correlation in fullfat and positive in lowfat yoghurt. The sensory assessment revealed that taste, flavour, colour, texture, and overall acceptability of yoghurt produced with 6% MLE (rated as liked very much) was significantly ( $p < 0.001$ ) better than that produced with 8 and 10% (rated liked slightly). It was concluded that fortification of FFY and LFY with 6% MLE produced acceptable yoghurt that has high nutritional value.

**Keywords :** moringa, fortification, yoghurt, bioactive compounds

**Conference Title :** ICASVM 2014 : International Conference on Animal Science and Veterinary Medicine

**Conference Location :** Sydney, Australia

**Conference Dates :** December 15-16, 2014