

Comparison Ileal and Excreta Digestibility of Protein Poultry by-Product Meal in 21 to 28 Days of Age Broiler Chicken

Authors : N. Mahmoudnia, M. Khormali

Abstract : This experiment was conducted to determine the apparent protein digestibility of poultry by- product meal (PBPM) from two industrial poultry slaughter houses on Ross 308 male broiler chickens in independent comparisons. The experiment consisted of seven dietary treatments and three replicates per treatment with three broiler chickens per replicate in a completely randomized design. Dietary treatments consisted of a control corn- soybean diet, and levels 3, 6 and 9% PBPM produced by slaughter house 1 and levels 3, 6 and 9% PBPM produced by slaughter house 2. Chromic oxide was added to the experimental diets as indigestible marker. The apparent protein digestibility of each diet were determined with two methods of sample collection of ileum and excreta in 21-28 d of age. The results this experiment showed that use of PBPM had no significant effect on performance of broiler chicks during period of experiments. The apparent protein digestibility of PBPM groups was significantly higher than control group by excreta sampling procedure ($P < 0.05$). Using of PBPM 2 significantly ($P < 0.05$) decreased the apparent protein digestibility values based on ileum sampling procedure vs control (85.21 vs 90.14).Based results of this experiment,it is possible to use of PBPM 1 in broiler chicken.

Keywords : poultry by-product meal, apparent protein digestibility, independent comparison, broiler chicken

Conference Title : ICVBS 2015 : International Conference on Veterinary and Biomedical Sciences

Conference Location : Prague, Czechia

Conference Dates : March 23-24, 2015