Bifidobacterium lactis Fermented Milk Was Not Effective to Eradication of Helicobacter Pylori Infection: A Prospective, Randomized, Double-Blind, Controlled Study

Authors : R. C. Barbuti, M. N. Oliveira, N. P. Perina, C. Haro, P. Bosch, C. S. Bogsan, J. N. Eisig, T. Navarro-Rodriguez **Abstract :** Background: The management of Helicobacter pylori (H. pylori) eradication is still a matter of discussion, full effectiveness is rarely achieved and it has many adverse effects. Probiotics are believed to have a role in eradicating and possibly preventing H. pylori infection as an adjunctive treatment. The present clinical study was undertaken to see the efficacy of a specially designed fermented milk product containing Bifidobacterium lactis B420 on the eradication of H. pylori infection in a prospective, randomized, double-blind, controlled study in humans. Method: Four test products were specially designed fermented milks, counts of viable cells in all products were 1010 Log CFU. 100 mL-1 for Bifidobacterium lactis-Bifidobacterium species 420, and 1011 Log CFU. 100 mL-1 for Streptococcus thermophiles were administered to subjects infected with H. pylori with a previous diagnosis of functional dyspepsia according to the Rome III criteria in a prospective, randomized, doubleblind, placebo-controlled study in humans. Results: After FM supplementation, not all subjects showed a reduction in H. pylori colonization. Conclusion: Bifidobacterium lactis B420, administered twice a day for 90 days did not show an increase in H. pylori eradication effectiveness in Brazilian patients with functional dyspepsia.

Keywords : antibacterial therapy, Bifidobacteria fermented milk, Helicobacter pylori, probiotics **Conference Title :** ICPFF 2015 : International Conference on Probiotics and Functional Foods **Conference Location :** Miami, United States

Conference Dates : March 09-10, 2015